




Draft Genome Sequence of the Planktic Cyanobacterium *Tychonema bourrellyi*, Isolated from Alpine Lentic Freshwater

Federica Pinto,^a Adrian Tett,^a Federica Armanini,^a Francesco Asnicar,^a Adriano Boscaini,^b Edoardo Pasolli,^a Moreno Zolfo,^a Claudio Donati,^b Nico Salmaso,^b  Nicola Segata^a

Centre for Integrative Biology, University of Trento, Trento, Italy^a; Edmund Mach Foundation, San Michele all'Adige, Italy^b

ABSTRACT We describe here the draft genome sequence of the cyanobacterium *Tychonema bourrellyi*, assembled from a metagenome of a nonaxenic culture. The strain (FEM_GT703) was isolated from a freshwater sample taken from Lake Garda, Italy. The draft genome sequence represents the first assembled *T. bourrellyi* strain.

Tychonema bourrellyi (J. W. G. Lund) (1) is a filamentous cyanobacterium historically found in northern regions, such as northern Europe and Canada (2). *Tychonema* is a cold-stenotherm genus with thin filaments (4 to 6 μm wide) and mainly colonizes pelagic freshwater environments. *T. bourrellyi* is an important cyanobacterial species, since some strains can produce the neurotoxins anatoxin-a (ATX) and homoanatoxin-a (HTX), which can be harmful for human and animal health. Recently, Salmaso et al. (3) found that *T. bourrellyi* is spreading and diversifying in Alpine freshwater lakes, with 49 distinct strains being identified in lakes previously dominated by *Planktothrix* species. However, despite the relevance of this species for human health, no sequenced *T. bourrellyi* strains are currently available.

We reconstructed the draft genome of *T. bourrellyi* from a nonaxenic culture. The culture originated from a water sample collected in summer 2014 by vertical tows from 20 to 30 m to the surface with a plankton net at Lake Garda. A single filament of *Tychonema* was isolated under the microscope and, after being washed repeatedly with Z8 Cellstar medium (Greiner Bio-One GmbH, Germany), was incubated at 20°C at constant-light illumination (3). The *Tychonema* culture was filtered through 0.22- μm filters, and the PowerWater DNA isolation kit (Qiagen) was then used to extract the DNA. The paired-end libraries (Illumina) were prepared and run on an Illumina HiSeq 2500 platform (100-nucleotide [nt]-long reads). Raw reads were assembled using metaSPAdes version 3.10.1 (4), with default parameters, which generated 7,029 contigs larger than 1,000 bp, with a total size of 74 Mbp. From this assembly, the draft genome of *T. bourrellyi* was binned using the manually supervised anvio protocol based on the abundance and tetranucleotide frequency distributions (5), quality controlled by CheckM (6) (95% predicted completeness with no indication of contamination or strain heterogeneity), and scaffolded through SSPACE (7).

The assembled draft genome was confirmed to belong to the *T. bourrellyi* species by PhyloPhlAn analysis (8) and BLASTn searches of the 16S and *rbcLX* genes, which were 98.7% and 100% identical, respectively, to previously reported *T. bourrellyi* genes (3). The genome is 5.08 Mb assembled into 271 scaffolds, with an N_{50} value of 29,997 bp and a GC content of 44.63%. The genome annotation was performed with Prokka version 1.11 (9), which identified 4,493 coding sequences and 49 tRNAs. A candidate of a polyketide synthase family protein gene cluster potentially linked with the neurotoxic

Received 17 October 2017 **Accepted** 1 November 2017 **Published** 22 November 2017

Citation Pinto F, Tett A, Armanini F, Asnicar F, Boscaini A, Pasolli E, Zolfo M, Donati C, Salmaso N, Segata N. 2017. Draft genome sequence of the planktic cyanobacterium *Tychonema bourrellyi*, isolated from Alpine lentic freshwater. *Genome Announc* 5:e01294-17. <https://doi.org/10.1128/genomeA.01294-17>.

Copyright © 2017 Pinto et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Nico Salmaso, nico.salmaso@fmach.it, or Nicola Segata, nicola.segata@unitn.it.

activity was identified in the genome at 40% amino acid identity (over an alignment length of 50%).

To our knowledge, this work represents the first sequencing of a *Tychonema bourrellyi* genome. Further analysis of the anatoxin genes involving more strains of this species are needed to unravel the genetic basis of *T. bourrellyi* toxicity.

Accession number(s). This genome project has been deposited at DDBJ/ENA/GenBank under the accession number [NXIB00000000](#). The version described in this paper is version NXIB02000000.

ACKNOWLEDGMENTS

This work was financially supported by the European Union's Horizon 2020 research and innovation program under Marie Skłodowska-Curie grant agreement 704603 to F. Pinto and Marie Skłodowska-Curie grant agreement 707345 to E. Pasolli and by the European Research Council (ERC-STG project MetaPG) to N. Segata.

REFERENCES

1. Anagnostidis K, Komárek J. 1988. Modern approach to the classification system of cyanophytes. 3. *Oscillatoriales*. Arch Hydrobiol Suppl 80:327–472.
2. Shams S, Capelli C, Cerasino L, Ballot A, Dietrich DR, Sivonen K, Salmaso N. 2015. Anatoxin-a producing *Tychonema* (*Cyanobacteria*) in European waterbodies. Water Res 69:68–79. <https://doi.org/10.1016/j.watres.2014.11.006>.
3. Salmaso N, Cerasino L, Boscaini A, Capelli C. 2016. Planktic *Tychonema* (*Cyanobacteria*) in the large lakes south of the Alps: phylogenetic assessment and toxigenic potential. FEMS Microbiol Ecol 92. <https://doi.org/10.1093/femsec/fiw155>.
4. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new versatile metagenomic assembler. Genome Res 27:824–834. <https://doi.org/10.1101/gr.213959.116>.
5. Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, Delmont TO. 2015. Anvi'o: an advanced analysis and visualization platform for 'omics data. PeerJ 3:e1319. <https://doi.org/10.7717/peerj.1319>.
6. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. Genome Res 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
7. Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. Bioinformatics 27:578–579. <https://doi.org/10.1093/bioinformatics/btq683>.
8. Segata N, Börnigen D, Morgan XC, Huttenhower C. 2013. PhyloPhlAn is a new method for improved phylogenetic and taxonomic placement of microbes. Nat Commun 4:2304. <https://doi.org/10.1038/ncomms3304>.
9. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.