

## Review

# Landscape of microalgae omics and metabolic engineering research for strain improvement: An overview



Evangelia Stavridou<sup>a</sup>, Lefkothhea Karapetsi<sup>a,b</sup>, Georgia Maria Nteve<sup>a</sup>, Georgia Tsintzou<sup>b</sup>, Marianna Chatzikonstantinou<sup>c</sup>, Meropi Tsaousi<sup>c</sup>, Angel Martinez<sup>d</sup>, Pablo Flores<sup>d</sup>, Marián Merino<sup>e</sup>, Luka Dobrovic<sup>f</sup>, José Luis Mullor<sup>e</sup>, Stefan Martens<sup>g</sup>, Leonardo Cerasino<sup>g</sup>, Nico Salmaso<sup>g</sup>, Maslin Osathanunkul<sup>h,i</sup>, Nikolaos E. Labrou<sup>j</sup>, Panagiotis Madesis<sup>a,b,\*</sup>

<sup>a</sup> Institute of Applied Biosciences, Centre for Research and Technology Hellas, 57001 Thessaloniki, Greece

<sup>b</sup> Laboratory of Molecular Biology of Plants, School of Agricultural Sciences, University of Thessaly, 38221 Volos, Greece

<sup>c</sup> Freshline Cosmetics, 1st km Lavriou Ave Koropiou—Markopoulou, GR-19400 Koropi, Greece

<sup>d</sup> Centro Tecnológico Nacional de la Conserva y Alimentación CTNC, Calle Concordia s/n. 30500, Molina de Segura (MURCIA), Spain

<sup>e</sup> Bionos Biotech S.L., Av. Fernando Abril Martorell, 106, 46026 Valencia, Spain

<sup>f</sup> Particula group LTD, Tina Ujevića 9, 51000 Rijeka, Croatia

<sup>g</sup> IASMA Research and Innovation Centre, Fondazione Edmund Mach, via E. Mach, 1, 38010 San Michele all'Adige (TN), Italy

<sup>h</sup> Department of Biology, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand

<sup>i</sup> Research Center in Bioresources for Agriculture, Industry and Medicine, Chiang Mai University, Chiang Mai 50200, Thailand

<sup>j</sup> Laboratory of Enzyme Technology, Department of Biotechnology, School of Applied Biology and Biotechnology, Agricultural University of Athens, 75 Iera Odos Street, GR-11855 Athens, Greece

## ARTICLE INFO

## ABSTRACT

## Keywords:

Genomics  
Transcriptomics  
Genetic engineering metabolism  
Microalgae applications  
Gene-editing  
Genetic transformation

The unique metabolic capabilities and fast growth rates of microalgae render them promising candidates for various industrial applications, such as biofuel production, pharmaceuticals, nutraceuticals, and wastewater treatment. Metabolic engineering is a powerful approach used to enhance the sustainable production of high-value compounds in microalgae, improve their stress tolerance, growth characteristics and suitability for large-scale cultivation. This review provides a snapshot of the current state of knowledge on omics and metabolic engineering research to further enhance our understanding on microalgal metabolism and enable the development of optimized strains with improved productivity and functionality. More specifically, it focuses on the recent breakthroughs in microalgal omics, driven by advancements in genomics technologies, such as improved sequencing platforms and bioinformatics tools, that have enabled the functional characterization of key genes, identification of metabolic pathways, and elucidation of microalgae cell physiology. Conventional and state-of-the-art genetic engineering approaches used in the last decades to manipulate the metabolic pathways of microalgae in a targeted manner, are highlighted in the scope of microalgal optimization. In this review, the different applications of genetic engineering and their impact on microalgae industry are also discussed. Integrating pan-omics data in future research is crucial for predicting novel functional interactions and identifying aspects of metabolic flux, towards enhancing algal strain-engineering techniques.

## 1. Introduction

Microalgae, one of the most ancient living forms on earth, are microscopic mono or multicellular organisms that include the prokaryotic cyanobacteria, and the eukaryotic, green algae and diatoms (Mishra

et al., 2019; Mobin and Alam, 2017). Microalgae are autotrophic organisms having high photosynthetic efficiency, and no intrinsic sensitivity to seasonality, due to the multiple coping mechanisms developed under different light regimes (Perin et al., 2019). They can recycle atmospheric carbon dioxide (CO<sub>2</sub>) (Zhu et al., 2017) and combined with

\* Corresponding author at: Institute of Applied Biosciences, Centre for Research and Technology Hellas, 57001 Thessaloniki, Greece.

E-mail addresses: estavrid@certh.gr (E. Stavridou), lefki8@certh.gr (L. Karapetsi), margodeves@certh.gr (G.M. Nteve), m.chatzikonstantinou@freshline.gr (M. Chatzikonstantinou), angel@ctnc.es (A. Martinez), mmerino@bionos.es (M. Merino), luka.dobrovic@particula-group.com (L. Dobrovic), jlmullor@bionos.es (J.L. Mullor), stefan.martens@fmach.it (S. Martens), maslin.o@cmu.ac.th (M. Osathanunkul), lambrou@aua.gr (N.E. Labrou), pmadesis@uth.gr (P. Madesis).

the use of non-agricultural land for biomass production can minimize the associated environmental and food-security impacts (Levasseur et al., 2020). The ability of microalgae to adapt to various abiotic stress factors, such as light, salinity, temperature, provides an interesting potential for the induction of high-value metabolites production (Gauthier et al., 2020; Little et al., 2021; Mishra et al., 2019; Rastogi et al., 2020; Ren et al., 2021; Suparmaniam et al., 2022).

There is a great diversity of microalgal species that provide energy-rich compounds for sustainable energy production, and other high value bioactive molecules of significant commercial interest, such as proteins, lipids and pigments used in pharmaceutical, nutraceutical, and cosmetics industries (Abu-Ghosh et al., 2021; Chung et al., 2017; Coulobier et al., 2021; De Luca et al., 2021; Martínez-Ruiz et al., 2022; Morais Junior et al., 2020). As such, a possible pipeline to optimize traits of high importance (i.e. yield, nutritional content) for large scale sustainable production may include i) species selection, ii) use of the trait plasticity/evolution through random DNA alteration or controlled DNA manipulation and iii) bioprocess optimization of the developed strain (Torres-Tijji et al., 2020).

Recent developments in genome sequencing, strain development, and genome editing technologies offer great opportunities for boosting microalgae productivity. Genomics, transcriptomics and the use of cutting-edge single-cell technology, as well as comparative data from different microalgae, would be beneficial in developing a comprehensive model of algae genomes (Shi et al., 2021). Algae Omics technologies, often known as "Algomics," provide insights into the evolutionary origins of algae and a comprehensive image of algae structure and system-function (Mishra et al., 2019; Ribeiro et al., 2017). In general, aspects of interest in microalgae research are focusing on omics approaches regarding biofuel and biofertilizer production, heavy metal bioremediation, pharmaceuticals, nutraceuticals, and cosmetics (Ahmad et al., 2022; Ranjbar and Malcata, 2022; Rumin et al., 2020).

The microalgae genomes database could enable the identification and distribution of gene families involved in high value molecules synthesis, among various selected microalgae models. However, only a limited number of whole genome sequences is available and is restricted to microalgae models (Brar et al., 2021; Fu et al., 2019), and therefore a significant and relevant mapping may not be possible. Gene characterization related to high value compounds synthesis using the -omics approach could facilitate the establishment of phyla-, genera-, or species-specific expression patterns, whereas genomics could be used as a quick guide in bioprospecting to discover compounds of interest. Further algal genome research will facilitate the access to functional and important genes, such as genes of industrial and biotechnological interest, for future applications (Abu-Ghosh et al., 2021).


Microalgae hold great potential for genetic engineering applications for enhancing productivity, accelerating growth rates, optimizing biofuel production, and generating valuable compounds. The omics technologies and the availability of numerous molecular tools and methods have significantly contributed to the advancement of algae biotechnology. Over the past five years, there has been a significant expansion in the genetic manipulation toolkit available for industrially important microalgal strains (Fajardo et al., 2020a, 2020b; Grama et al., 2022; Sproles et al., 2021). Numerous genetic transformation methods were initially created and improved using the green algae *Chlamydomonas reinhardtii*, and their use has subsequently been extended to other groups of algae, including red algae (rhodophytes), brown algae (phaeophytes), euglenoids, diatoms, and dinoflagellates. Recent efforts to enhance the quality and quantity of desired products in microalgae have concentrated on the development of targeted genome editing tools, such as the CRISPR/Cas9, TALEN, zinc-finger nuclease (ZFN) and RNA interference (RNAi) technologies, along with the development of high-throughput screening methodologies for fast and robust strain improvement procedures (Liang et al., 2019; Lin et al., 2019; Patel et al., 2019).

The current knowledge on the strategies used to induce and improve the bioproduction of high-added value metabolites in microalgae,

including natural diversity, optimization of growth and productivity in a crosstalk with omics technologies and genetic engineering tools, towards application in various industrial sectors is presented in Fig. 1. This review focuses on the omics combinatorial approach to high-throughput analysis of microalgae for trait improvement. It broadly encompasses genomics, transcriptomics, and genetic engineering tools applied to improve microalgal traits. It first covers a global presentation of genomics, transcriptomics and genetic engineering, to finally summarize and evaluate the application of such technologies in microalgae industry. This review aims to explore the different genetic engineering methods employed in algae and their potential applications in greater detail. The objective of this review is to provide insights into the recent landscape of omics approaches to unravel the intricate metabolic pathways and regulatory networks within microalgae and the use of genetic engineering tools in boosting microalgae strain improvement through targeted enhancement of desirable traits highlighting the challenges that need to be remediated. The different applications of genetic engineering and their impact on microalgae industry are also discussed. It also emphasizes and discusses the potential aspects of using genetic engineering for industrial production of bioactive compounds.

## 2. Genomics

Algae are among the most genetically complex organisms known. Genomic analysis is a powerful tool, providing crucial insights into the genetic complexity of microalgae. Genome sequencing projects for different species of microalgae have been completed, allowing the discovery and annotation of genes involved in numerous biological processes. Genomic resources have been established for several groups of microalgae, notably Chlamydomonas, Nannochloropsis, and diatoms. This availability of genomic data has provided valuable insights into the genetic makeup and molecular mechanisms of these microalgae species (Armbrust et al., 2012; Gong et al., 2020; Merchant et al., 2010). Comparative genomics studies have played a crucial role in investigating the evolutionary connections between different microalgal species, identifying expansions or contractions in gene families, and pinpointing conserved regions shared among diverse microalgal taxa. The genome of *Cyanidioschyzon merolae*, a red extremophile alga, was



**Fig. 1.** Graphical outline on the current knowledge of microalgae strain improvement using the omics resources and genetic engineering strategies. Natural diversity in microalgae and optimization techniques of growth and bioproduction also play an important role in strain improvement towards enhancing stress tolerance and production of high added value compounds.

the first to be published in 2004 (Matsuzaki et al., 2004), and although *Chlamydomonas reinhardtii*, a single-celled flagellate freshwater green alga, is one of the most researched species (Mishra et al., 2019) its genome was published only later in 2007 (Merchant et al., 2010).

Currently, 191 algal genomes are publicly available, representing a diverse spectrum of algae taxa. Among these are 117 green algae (111 Chlorophyta and six Charophyta), 34 Heterokonts (diatoms), 12 Rhodophytes (red algae), ten Dinoflagellates, six Haptophytes, five Cryptophyceae, and a Glaucophyte (Shi et al., 2021) (Table 1). Green algae have the highest representation, accounting for approximately 58% of the sequenced algal genomes, followed by Heterokonts (18%), red algae (6%), the toxin producing Dinoflagellates (5%), and Haptophytes (5%). Together, these five groups represent 92% of the published algal genomes. Green algae are the most extensively studied, constituting 91% (174) of all published algal genomes. Among the sequenced genomes, the green alga *Ostreococcus tauri* (Prasinophyceae) is the smallest known free-living eukaryote, with a genome size of 12.56 Mb (Shi et al., 2021), whereas the biggest known the genome is that of the Dinoflagellate *Lingulodinium polyedrum* having a size of 185 Gbp (Casabianca et al., 2017). Regarding the organelle genomes, *Grateloupia taiwanensis* stands out with the highest gene richness, having 233 protein-coding genes (DePriest et al., 2013). In contrast, *Cyanidioschyzon merolae*, a red microalga, has the least gene-rich nuclear genome, with only 4775

protein-coding genes (Nozaki et al., 2007). Interestingly, up to 20% of the nuclear genes in the green microalgae appear to have originated from the cyanobacterial endosymbiont through a process called endosymbiotic gene transfer (Dagan et al., 2013; Deusch et al., 2008) highlighting the significant impact of endosymbiosis on the evolution and genetic composition of microalgae.

Thus far, and in terms of completed genomes, only a few species have been fully sequenced without any gaps (Table 1). These include the relatively small nuclear genomes of the red alga *Cyanidioschyzon merolae* (16.5 Mbp) (Nozaki et al., 2007) and the Prasinophytes *Micromonas commoda* RCC299 (Worden et al., 2009) and *Ostreococcus lucimarinus* (Palenik et al., 2007). The number of published algal genome sequences continues to rise, driven by advancements in sequencing technologies, reduction in sequencing costs, and improved sequencing quality. The ALG-ALL-CODE project focused on sequencing over 120 genomes of microalgae, obtained from newly isolated and maintained stocks provided by culture collection centers, such as UTEX and NMCA (Ashworth and Ralph, 2018). According to the latest update of the project (March 2021), 22 subtropical microalgal species have been isolated and sequenced from the UAE and 107 additional species were obtained from international algal culture collections, including UTEX ([www.utex.org](http://www.utex.org)) and the Bigelow National Center for Algae and Marine Microbiota (<https://ncma.bigelow.org/>). Notably, the 10KP Genome Sequencing project initiative, launched in 2017, aims at further expanding this dataset by sequencing the genomes of 1000 green algae (chlorophytes and streptophytes) and 3000 protists (photosynthetic and heterotrophic) (Sirohi et al., 2021), with, thus far, having sequenced and assembled over 300 green algae species.

The diversity of transposable elements (TEs), some of which form new classes, is another aspect revealed by algal whole genome sequencing (Tirichine and Bowler, 2011). Transposable elements are repetitive genetic sequences that play a vital role in the genome by being capable of relocating to different positions within the genome. This mobility can cause disruptions in the normal functioning of genes and induce alterations in the overall structure of the genome. The ratios of TEs in algae range from 5% to 40%, compared to 161 terrestrial plants, which normally contain approximately 50% of repeated sequences (Shi et al., 2021).

Studies have shown that horizontal gene transfer (HGT) in green and red algae is associated with various functions such as carbohydrate metabolism, osmolyte management, sulfate scavenging, cell cycle control, energy shuttle, arsenic detoxification, and stress adaptation mechanisms, with many algae exhibiting relatively low levels of HGT gene percentages except for a few specific species, such as Charophyta, Dinophyceae, Haptophyta and Stramenopiles (Heterokonts) (Foflonker et al., 2015; Hirooka et al., 2017). These genomic resources serve as invaluable tools for studying microalgae biology, identifying important genes and pathways, and facilitating genetic engineering and biotechnological applications in the field.

### 3. Transcriptomics and methylomics

Transcriptomics is a powerful approach to unveil gene expression patterns across the entire transcriptome, allowing the investigation of an organism's capabilities and limitations related to various traits such as resistance, acclimatisation, and developmental processes (Page and Lawley, 2022). *De novo* assemblies (Blanc-Mathieu et al., 2014), microbiome investigation (Carney et al., 2014), and other genome sequencing applications (Teng et al., 2019), have become the gold standard method for transcriptome studies (Rismani-Yazdi et al., 2012), with >700 microalgae transcriptomes being available on alga database ([www.algaedb.org/](http://www.algaedb.org/)). The "Algaenaut" is an interactive web resource available to the public, designed for the exploration of microalgae transcriptomes (Ashworth and Ralph, 2018). This resource combines a total of 1375 samples obtained from 69 independent experimental trials, representing ten different microalgae species and

**Table 1**  
Summary of the complete or nearly complete genomes of different microalgae species.

Species	Genome Size (Mb)	References
<i>Auxenochlorella protothecoides</i>	22,92	[Vogler et al., 2018; Xing et al., 2018]
<i>Bathycoccus prasinos</i>	15,07	[Moreau et al., 2012]
<i>Botryococcus braunii</i>	164,2	[Browne et al., 2017]
<i>Chlamydomonas debaryana</i>	120,36	[Yoshimitsu et al., 2018]
<i>Chlamydomonas reinhardtii</i>	120,4	[Salguero et al., 2018]
<i>Chlamydomonas sphaerooides</i>	126	[Hirashima et al., 2016]
<i>Chlorella pyrenoidosa</i>	56,99	[Run et al., 2016]
<i>Chlorella sorokiniana</i>	58,53	[Dawson et al., 1997]
<i>Chlorella variabilis</i>	46,16	[Juneja et al., 2016; Sorigué et al., 2016]
<i>Chlorella vulgaris</i>	39,08	[Zuñiga et al., 2018; Mócsai et al., 2019]
<i>Chloridium sp.</i>	54,31	[Nelson et al., 2017]
<i>Chondrus crispus</i>	104,98	[Collén et al., 2013]
<i>Chromochloris zofingiensis</i>	58	[Liu et al., 2014b; Liu et al., 2019]
<i>Coccomyxa sp.</i>	48,54	[Koechler et al., 2016]
<i>Cyanidioschyzon merolae 10D</i>	17	[Nozaki et al., 2007]
<i>Dunaliella salina</i>	550	[Polle et al., 2017]
<i>Galdieria phlegrea DBV009</i>	11,4	[Qiu et al., 2013]
<i>Galdieria sulphuraria</i>	13,71	[Thangaraj et al., 2011]
<i>Gonium pectorale</i>	148,8	[Hanschen et al., 2016]
<i>Haematococcus pluvialis</i>	669	[Luo et al., 2019]
<i>Helicosporidium sp.</i>	13,2	[Pombert et al., 2014]
<i>Klebsormidium nitens</i>	104,21	[Nagao et al., 2008]
<i>Micromonas pusilla</i>	21,96	[Kujawinski et al., 2017]
<i>Monoraphidium neglectum</i>	69,71	[Jaeger et al., 2017]
<i>Nannochloropsis gaditana</i>	29	[Fernández-Acero et al., 2019; Patelou et al., 2020]
<i>Nannochloropsis oceanica</i>	29,3	[Guo et al., 2019]
<i>Ostreococcus lucimarinus</i>	13,2	[Krumholz et al., 2012]
<i>Ostreococcus tauri</i>	13,03	[Blanc-Mathieu et al., 2014]
<i>Parachlorella kessleri</i>	64,9	[You et al., 2019]
<i>Picochlorum sp.</i>	13,5	[Foflonker et al., 2015]
<i>Porphyra umbilicalis</i>	87,7	[Brawley et al., 2017]
<i>Porphyridium purpureum CCMP1328</i>	19	[Bhattacharya et al., 2013]
<i>Pyropia yezoensis U-51</i>	108	[Wang et al., 2020a, 2020b]
<i>Senedesmus obliquus</i>	107,72	[Wang et al., 2019]
<i>Senedesmus sp.</i>	93,24	[Wang et al., 2019]
<i>Tetraselmis striata</i>	227,95	[Steadman Tyler et al., 2019]
<i>Volvox carteri f.Magariensis</i>	137,68	[Prochnik et al., 2010]

four distinct clades. Additionally, the Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP) (Keeling et al., 2014) has created 678 assembled, functionally annotated, and publicly available transcriptomes of 306 pelagic and endosymbiotic marine eukaryotic species representing >40 phyla.

The utilization of transcriptomic analyses can greatly accelerate advancements in harnessing algae for biofuel production. One example is the unicellular green algae species *Chlamydomonas moewusii*, which produce hydrogen under both light and dark anaerobic conditions (Sirohi et al., 2021). By conducting RNA sequencing (RNA seq) transcriptomic studies, over a specific time frame, valuable insights can be obtained regarding the expression patterns of genes involved in anaerobic fermentation, glycolysis, starch catabolism, and hydrogen evolution in dark anaerobic conditions. Such transcriptomic investigations serve as a foundation for characterizing and understanding the underlying genetic mechanisms and provide valuable information for potential genomic modifications aimed at enhancing specific production capabilities or improving stress tolerance in algae (Table 2). Knowledge on the genes and gene functions in microalgae species, such as the model organisms *Chlamydomonas* and *Nannochloropsis spp.*, has assisted on the application of genetic engineering tools such as, gene editing and RNA interference (RNAi), given the available genetic resources and mutant collections (<https://www.chlamylibrary.org/>) of these species (further examples are presented in detail in section 4). Similarly, transcriptional engineering has also been used to overexpress the endogenous *bzip* transcription factors (TFs) mediated by the promoter of the heat-shock protein HSP20, increased the lipid content in the transformant lines compared to wild strains in *Nannochloropsis salina* (*NsbZIP1*) and *N. oceanica* (*NobZIP1*) (Kwon et al., 2018; Li et al., 2019). The processes underlying stress adaptation and metabolite production in microalgae at the epigenetic level are still unknown despite the crucial role in microalgae adaptation to the variable growth environments. Recent studies have provided evidence of methylation events occurring in algae. The *C. reinhardtii* nuclear genome is methylated at low levels (5.4, 2.6 and 2.5% in CG, CHG and CHH context, respectively). Similarly, *Chlorella sp.* demonstrates even lower levels of CG methylation, ranging from 4% to 5%. These findings shed light on the presence of methylation processes in algae genome (Ferrari et al., 2023). Microalgal genomes are shaped by epigenetic markers, which also control gene expression and may have an impact on how accessible the genome is to TFs or alter the expression of TFs. For instance, in *C. reinhardtii*, various epigenetic alterations altering TF activity have been discovered in relation to histone modifications (Bacova et al., 2020). At present, there is increasing interest in the microalgal epigenome, especially regarding the regulation of gene expression under variable conditions (Bacova et al., 2020; Ferrari et al., 2023). The genome of *C. reinhardtii* comprises of both <sup>6</sup>mA and <sup>5</sup>mC modified bases compared to those of other unicellular organisms (Hattman et al., 1978). After mapping the distribution of the <sup>6</sup>mA modification in >84% of genes of *C. reinhardtii*, the <sup>6</sup>mAs were found to cluster around the transcriptional start sites (TSSs) of active genes (Fu et al., 2015). Future studies are anticipated to unravel the role of epigenetics in microalgal biochemical pathways, which might be used to adjust cellular and physiological balance to boost the production of beneficial metabolites.

### 3.1. Environmental stresses

Transcriptomics have been used in microalgae research to mainly elucidate the mechanisms involved in environmental adaptation and acclimation (Rismani-Yazdi et al., 2012) (Table 2). Environmental stresses, such as nutrient deprivation, increased salinity, and temperature fluctuations, shifting light intensities and toxic compounds including heavy metals and organic toxins, that have a significant impact on microalgae growth and metabolism (Al-Ealayawi and Al-Dulaimy, 2023) and (Tirichine and Bowler, 2011). Nevertheless, different taxa have variable natural tolerance to such environmental

**Table 2**

Summary of the different conditions and abiotic stresses tested on various microalgae species in transcriptomics studies of the last decade from 2014 to 2023.

Conditions	Species	References
Growth conditions	<i>Chlorella sp.</i>	[Koh et al., 2023]
	<i>A. protothecoides</i>	[Steichen et al., 2022]
	<i>Symbiodinium microadriaticum</i>	[Aranda et al., 2016]
	<i>Micromonas sp</i>	[Van Baren et al., 2016]
	<i>A. protothecoides</i>	[Gao et al., 2014]
	<i>Ettlia oleoabundans</i>	[Sturme et al., 2018]
Oil accumulation	<i>Lobosphaera bisecta</i>	[Gao et al., 2020]
	<i>Schizochytrium sp.</i>	[Chang et al., 2021]
Different growth phases	<i>S. obliquus</i>	[da Silva et al., 2021], [Xi et al., 2021]
	<i>Chlorococcum sp</i>	[Lv et al., 2019]
	<i>Messastrum gracile SE-MC4</i>	[Afifudeen et al., 2021]
Day-night cycle	<i>Chlamydomonas sp.</i>	[Willamme et al., 2018]
<b>Stresses</b>		
Temperature stress	<i>A. protothecoides</i>	[Xing et al., 2018]
	<i>Emiliania huxleyi</i>	[Brenner et al., 2013]
	<i>Fragilariaopsis cylindrus</i>	[Jabre et al., 2021]
	<i>Picochlorum sp.</i>	[Barten et al., 2022]
	<i>Symbiodinium sp.</i>	[Gierz et al., 2017]
	<i>Chlorella sp.</i>	[Poong et al., 2018]
Phosphatase stress	<i>Schizochytrium sp.</i>	[Hu et al., 2022]
	<i>C. reinhardtii</i>	[Légeret et al., 2016]
	<i>A. protothecoides</i>	[Park et al., 2018]; [Bachy et al., 2018]
	<i>M. pusilla</i>	[Fang et al., 2015]
	<i>B. braunii</i>	[Yang et al., 2022]
	<i>C. reinhardtii</i>	[Nordin et al., 2020]
Nitrogen deprivation	<i>C. vulgaris</i>	[Zhu et al., 2022]
	<i>C. sorokiniana</i> (1230)	[de Jaeger et al., 2018]
	<i>Neochloris oleoabundans</i>	[Jaeger et al., 2017]
	<i>M. neglectum</i>	[Polle et al., 2020], [Wang et al., 2018a, 2018b]
	<i>D. salina</i>	[Wang et al., 2018a, 2018b]
	<i>C. reinhardtii</i>	[Sun et al., 2014]
Salinity	<i>C. pyrenoidosa</i>	[Abdellaoui et al., 2019]
	<i>C. vulgaris</i>	[de Jaeger et al., 2018]
	<i>N. oleoabundans</i>	[You et al., 2019]
	<i>P. kessleri</i>	[Fan et al., 2016]
	<i>C. pyrenoidosa</i>	[Li et al., 2016a, 2016b]
	<i>C. sorokiniana</i>	[Peng et al., 2016]
CO <sub>2</sub>	<i>Coccomyxa subellipsoidea</i>	[Rademacher et al., 2016]
	<i>C. merolae</i>	[Wang et al., 2022a, 2022b]
	<i>E. huxleyi</i>	[Wei et al., 2019]
	<i>N. oceanica</i>	[Clement et al., 2017]
	<i>Thalassiosira pseudonana</i>	[Huang et al., 2022]
	<i>Dunaliella salina</i>	[Chernobai et al., 2022]
Response to osmotic and oxidative stress	<i>D. salina</i>	[Choi et al., 2023]
	<i>C. reinhardtii</i>	[Rossoni et al., 2019]
Cold acclimation	<i>Galdieria sulphuraria</i>	[Azaman et al., 2020]
	<i>Chlorella sp.</i>	[He et al., 2015]
Light intensity	<i>Monoraphidium sp.</i>	[Breuer et al., 2013]
	<i>S. obliquus</i>	[Carvalho and Malcata, 2005]
UV radiation	<i>Pavlova lutheri</i>	[Mitra et al., 2015]
	<i>N. gaditana</i>	[Ekhtari et al., 2019]
	<i>V. carteri f. Magariensis</i>	[Tardu et al., 2016]
	<i>C. merolae</i>	[Zheng et al., 2020]
	<i>Pb - C. reinhardtii</i>	[Ding et al., 2020]
	<i>Cd - C. sorokiniana</i>	[Tian et al., 2022]
Heavy metals	<i>Cd - Synechocystis sp</i>	[Zhu et al., 2019]
	<i>Cd - D. salina</i>	[Puente-Sánchez et al., 2018]
	<i>Cd - Chlamydomonas acidophila</i>	[Lu et al., 2019]
	<i>Cd - A. protothecoides</i>	[Simon et al., 2008]
	<i>Cd - C. reinhardtii</i>	[Wang et al., 2020a, 2020b]
	<i>Cu - Microcystis aeruginosa</i>	[Willamme et al., 2018]
		(continued on next page)

**Table 2 (continued)**

Conditions	Species	References
Nanoparticles	Cu – <i>C. reinhardtii</i>	[Beauvais-Flück et al., 2019]
	Co - <i>D. salina</i>	[Ling et al., 2021]
	Co – <i>B. braunii</i>	[Cheng et al., 2018]
	Hg – <i>Chromera velia</i>	[Sharaf et al., 2019]
	Hg – <i>C. Reinhardtii</i>	[Beauvais-Flück et al., 2017]
	FeO – <i>O. Tauri</i>	[Genevière et al., 2020]
Wastewater	ZnO – <i>M. Commoda</i>	[Bibi et al., 2021]
	α-Fe2O3 – <i>C. vulgaris</i>	[Wang et al., 2022a, 2022b]
Wastewater	<i>Desmodesmus sp</i>	[Wang et al., 2022a, 2022b]

stressors. The red algae *Galdieria sulphuraria* can thrive in a pH range of 0 to 4 and at temperatures as high as 56 °C (Shi et al., 2021). The halophytic algae *Emiliania huxleyi* creates algal blooms from the equator to the subarctic, and under a wide range of climatic conditions. Whereas the green alga *D. salina* (Polle et al., 2017), *Picochlorum SENEW3* (Foflonker et al., 2015) and *Picochlorum renovo* (Dahlin et al., 2019) exhibit enhanced salt tolerance, under a range of 1,4 to 1,6 M NaCl.

Microalgae cultivation under different abiotic stress conditions may increase the synthesis of high-value metabolites, whilst in the past decades research has focused on the effects of variable conditions on metabolic pathways. Many research groups are focusing on microalgae adaptation in fuel gas settings, primarily comprising CO<sub>2</sub>, Nox, and Sox (Cheng et al., 2016; Listmann et al., 2020; Schaum et al., 2017). The adaptation strategies of microalgae under nutrient stress conditions have also been evaluated unlocking their mechanisms for survival and growth in nutrient-limited environments (Helliwell et al., 2015; Li et al., 2016b). Conventional lipid enhancement strategies for sustainable microalgal biodiesel production include nutrient stress and alterations in cultivation conditions (Singh et al., 2016; Zienkiewicz et al., 2016). One of the leading approaches in deciphering the mechanisms of algal lipid metabolism is the functional analysis of potential candidate genes governing lipid synthesis in algal cells through comparing global gene expression in nutrient-replete and nutrient-deprived cells of different microalgal species (Zienkiewicz et al., 2016). More specifically, nitrogen and silicon limitation along with CO<sub>2</sub> enrichment coupled with low temperature shock were shown to improve lipid content and stimulate lipid productivity in the diatom *Skeletonema costatum* (Gao et al., 2019; Xie et al., 2022).

Transcriptomics approach has also been used to assess and enhance the heat tolerance of *Dunaliella bardawil*, a halophilic green alga strain widely utilized for outdoor production of β-carotene showing that genes encoding for heat shock proteins (HSP) and antioxidant enzymes were up-regulated (Sirohi et al., 2021). Stress tolerance of microalgae to ultraviolet radiation (UVR) and high light intensity has also been investigated in the frame of inducing co-tolerance between abiotic and chemical stress factors, such as herbicides (Deblois et al., 2013; Korkaric et al., 2015), indicating that abiotic stress acclimation can provide evidence on the mechanistic effect of both stressors and the underlying co-tolerance mechanisms.

Moreover, research on salt tolerance mechanisms in microalgae has increased our understanding on osmolyte accumulation and physiological changes (Hu et al., 2020; Kato et al., 2022; X. Li et al., 2018; Meijer et al., 2017; Shetty et al., 2019; X. M. Sun et al., 2018). Transcriptomic analysis of the green microalga *C. sorokiniana* under high salt conditions, have unraveled the upregulation of genes involved in osmoregulation and ion transport processes (Azaman et al., 2020). Similarly, different microalgae strains exhibit distinctive iron uptake mechanisms under iron deficiency, such as diversity in proteins responsible for iron absorption and their regulation under variable iron levels or iron-starvation-induced proteins and iron-reductases through siderophore-mediated iron assimilation (reviewed by Rana and Prajapati, 2021).

Recent genomics analysis of the diatom *T. pseudonana* under iron deficiency has revealed the presence of multiple genes related to iron assimilation and transport, indicating a robust adaptive response to iron scarcity (Al-Ealyawi and Al-Dulaimy, 2023).

Microalgal species play a significant role in environmental pollution monitoring and bioremediation through the phytoremediation of both hazardous metals and nanometals (Abdel-Ghany et al., 2015). Microalgae play an important role in environmental remediation by absorbing, sequestering, and altering toxic metals and nanometals. Their participation in phytoremediation and monitoring renders them valuable tools for tackling the consequences of metal contamination in aquatic environments (Touliabah et al., 2022). The tolerance of microalgae to organic pollutants (Cho et al., 2016; H. Li et al., 2021; X. Wang et al., 2019), high concentrations of heavy metals, and under wastewater conditions (Okurowska et al., 2021; Wang et al., 2018a; Yu et al., 2020) has been examined. Research on algae transcriptome in response to different pollutant stresses is a commonly employed approach to uncover molecular changes and assess biological risks accurately and sensitively (Duan et al., 2019; Wang et al., 2018a; Q. Zhu et al., 2021) that can potentially enhance the bioremediation ability of microalgae.

#### 4. Genetic engineering

The advancement of algae biotechnology greatly benefits from the valuable and comprehensive data repositories provided by omics technologies. The field of microalgae biotechnology is gaining increasing commercial significance across a wide range of industries including food, pharmaceuticals, nutraceuticals, cosmetics, animal feed, energy, and environmental applications, such as water and soil remediation (Singh et al., 2021; Vanhoudt et al., 2018) as the genetic modification of microalgae expands the biotechnological applications. In this context, genetic engineering plays a key role in molecular farming, also known as gene pharming, that involves the production of enzymes, antibodies, immunotoxins, bioactive peptides, and hormones (Hempel et al., 2011; Mayfield, 2013; Rasala and Mayfield, 2011; Rosales-Mendoza, 2016; E. Specht et al., 2010; E. A. Specht and Stephen, 2014; Yan et al., 2016; Yusibov et al., 2016).

Thus far, the process of modifying the alga genome has traditionally relied on known genetic engineering tools (Table 3) followed by selecting desired mutations. However, with the advent of genome editing tools, or gene editing techniques given their focus on the modification of specific targeted genes in microalgae (Jeon et al., 2017), it is possible to achieve precise and specific genomic mutations (Table 3). These editing tools enable the targeting and modification of specific regions in the microalgae genome, utilizing sequence-specific nucleases, such as zinc finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs), and CRISPR-associated protein 9 (Cas9). Among these nucleases, Cas9 and related Cas proteins are preferred in microalgae genetic engineering due to their simplicity, precision, and efficiency (Fayyaz et al., 2020; Jeong et al., 2023). Furthermore, chloroplast transformation techniques have been successfully developed and applied to *C. reinhardtii* and various other microalgal species. In these approaches, antibiotic resistance genes, herbicide resistance genes, and metabolic marker genes have been implemented (Doron et al., 2016; Esland et al., 2018).

The first report on gene editing in microalgae was in *C. reinhardtii*, using zinc finger nucleases (ZFNs) to target the COP3 gene as a proof-of-concept experiment (Sizova et al., 2021). Furthermore, engineered transcription activator-like effector nucleases (TALENs) have also been employed in microalgae (Dubois et al., 2014). Nevertheless, the CRISPR-Cas system has become the preferred choice for genome editing in microalgae due to its versatility and simplicity compared to the ZFNs and TALENs (Greiner et al., 2017; Kurita et al., 2020). Since the initial efforts to use CRISPR-Cas9 in Chlamydomonas, further research studies have incorporated ten different genera of eukaryotic microalgae, leading to the development of new strains with improved productivity,

**Table 3**

Transformation methods and gene mutants used for different algae species and strains for genetic engineering from 1997 to 2022.

Species/Strain	Transformation method	Gene mutants	References
<i>C. reinhardtii</i>	Bioelectric, Glass bead agitation/ Electroporation/ <i>Agrobacterium</i> -mediated	Disruption of the photoreceptor genes <i>COP1/2</i> , <i>COP3</i> (encoding channelrhodopsin 1 [Chr1]), <i>COP4</i> (encoding Chr2), <i>COP5</i> , <i>PHOT</i> , <i>UVR8</i> , <i>VGCC</i> , <i>MAT3</i> , and <i>aCRY</i> and creation of the <i>chr1 chr2</i> and <i>uvr8 phot</i> double mutants; generation of an IR transgene targeting the <i>EXO</i> gene (protein ID 407657); insertion of <i>Paromomycin</i> ( <i>Paro</i> ) gene and a reporter gene ( <i>Renilla reniformis</i> luciferase ( <i>Luc</i> )) Mutants of <i>PPXJ1</i> and <i>FTSY</i> gene; knock-out of <i>cptfsy</i> gene; knock-out mutants of <i>cp29</i> and <i>cp26</i> gene; targeted knockout of <i>CrSPD1</i> ; knock-in of the gene <i>CrFTSY</i> using a combination of ribonucleoprotein (RNP) complex and DNA fragment (antibiotics resistance gene); knockout mutant of the zeaxanthin epoxidase gene ( <i>ZEP</i> ), double deletion knock-out mutants (dZAs), for the <i>ZEP</i> (Zeaxanthin epoxidase) and <i>AGP</i> (ADP glucose pyrophosphorylase) genes	[Greiner et al., 2017; E. J. Kim and Cerutti, 2009; Mini et al., 2018]
<i>C. reinhardtii</i> (CW15, CC-503, CC4349, CC124, CC503, CC-4349, CC-5415, CC-124, UVM4, N-UVM4, CC-4349)	Electroporation/RNP/Cas9		[Akella et al., 2021; Angstenberger et al., 2020; Cazzaniga et al., 2020; Freudenberg et al., 2022; J. H. Kim et al., 2020; Song et al., 2020, 2022]
<i>C. reinhardtii</i> (CC-124, CC-400, UVM11)	Cell penetrating peptide/ RNP/Cas9	Disruption of the genes <i>Maa7</i> and <i>FKB12</i> through a ribonucleoprotein delivery system mediated by the cell penetrating peptide pVEC (LILILRRRIRKQAHASK)	[Kang et al., 2020]
<i>C. reinhardtii</i> (CC-125, CC-5415)	Electroporation/ Glass bead/ RNP/Cas9	Mutants of <i>IFT81</i> (first gDNA), <i>IFT43</i> , <i>MOT17</i> , <i>CDPK13</i> , <i>FAP70</i> , <i>IFT81</i> (second gRNA), and <i>CEP131</i> , through TIM Loss-of-function mutation of <i>FK506-binding protein 12</i> ( <i>FKB12</i> ) locus	[Picariello et al., 2020]
<i>C. reinhardtii</i> (CC-1883)	Electroporation/RNP/Cas12a		[Ferenczi et al., 2021]
<i>Phaeodactylum tricornutum</i> (CCMP2561)	Bacterial conjugation/ Episomal plasmid/ Cas9, Bombardment/Plasmid /Cas9	Mutants of <i>NR</i> gene	[Moosburner et al., 2020]
<i>P. tricornutum</i> (CCMP2561)	Bombardment/Plasmid/Cas9	Mutants of light-harvesting complex ( <i>LHC</i> ) genes belonging to the <i>LHCF</i> gene family	[Sharma et al., 2021]
<i>P. tricornutum</i> (CCMP2561)	Bacterial conjugation/ episomal plasmid/ Cas9	knock-out of cryptochrome <i>CryP</i> ( <i>CryP</i> gene)	[Yang et al., 2022]
<i>P. tricornutum</i>	Bioelectric/ Electroporation/ Bacterial conjugation	Knock-out mutants of <i>PtUMPS</i> ( <i>Phatr3.J11740</i> ) and <i>PtAureo1a</i> ( <i>Phatr3.J8113</i> ) genes; single base pair mutation of the Myb transcription factor (XM_002181623; named <i>Pt_Myb1R_SHAQKYFS</i> )	[Serif et al., 2018; Sharma et al., 2018]
<i>P. tricornutum</i> (CCAP1055/1)	Electroporation/plasmid/ Cas9	knock-out mutants of <i>THIC</i> , <i>SSSP</i> , <i>SSUA/THI5-like</i> (encoding NMT1 domains) and <i>FOLR</i> genes	[Llavero-Pasquina et al., 2022]
<i>T.pseudonana</i>	Bioelectric/ Bacterial conjugation	Insertion of the <i>CEN6-ARSH4-HIS3</i> fragment	[Karas et al., 2015]
<i>T. pseudonana</i> (CCMP1335)	Bombardment/Plasmid/Cas9	Indels mutations of the <i>TpθCA3 THAPS DRAFT_bd1765</i>	[Nawaly et al., 2020]
<i>Nannochloropsis salina</i> (CCMP1776)	Electroporation/plasmid/ Cas9	Knockout and knockdown mutants of cellulose synthase gene ( <i>CesA</i> )	[Jeong et al., 2020]
<i>N. oceanica</i> IMET1	Electroporation/RNP/ Cas12a, CRISPR	Down-regulation of <i>NR</i> (nitrate reductase) gene	[Naduthodi et al., 2021]
<i>N. salina</i> (CCMP1776)	Electroporation/Cas9	Knock-in of the <i>FAD12</i> gene, encoding Δ12-fatty acid desaturase	[Ryu et al., 2021]
<i>N. gaditana</i> (CCMP526)	Electroporation/Cas9	Knock-out of beta-glucan synthase ( <i>BGS</i> ) gene and transglycosylase ( <i>TGS</i> ) gene	[Vogler et al., 2021]
<i>N. oceanica</i> IMET1	Electroporation/episomal plasmid/Cas9	Telomere-deletion mutants and large genome fragment deletion	[Wang et al., 2021]
<i>N. oceanica</i> (CCMP1779)	Electroporation/episomal plasmid/Cas9	Deletion mutants of the <i>NoAUREO2</i> or <i>NoAUREO4</i> genes	[Poliner et al., 2018]
<i>N. gaditana</i>	Electroporation	knock-out mutants of a homolog of fungal Zn(ii)2Cys6-encoding genes	[Ajjawi et al., 2017]
<i>N. limnetica</i>	Electroporation	Insertion of bleomycin-resistant gene ( <i>sh ble</i> ) or hygromycin B-resistant gene ( <i>hprt</i> )	[Y. Chen and Hu, 2019]
<i>N. oceanica</i>	Electroporation	Insertion of β-tubulin promoter, <i>sh ble</i> gene, and <i>fcpA</i> terminator; random mutation of genomic DNA by insertion of a Transposone complex Tn5; overexpression of EPA biosynthesis genes	[Li et al., 2014; Osorio et al., 2019; Poliner et al., 2018]
<i>N. oculata</i>	Electroporation	Insertion of β-tubulin promoter, <i>sh ble</i> gene, and <i>fcpA</i> terminator	[F. Li et al., 2014]
<i>N. oceanica</i> IMET1	Electroporation/plasmid/ CRISPRa	Over-expression of gene <i>g1248</i>	[Wei et al., 2022]
<i>C. pyrenoidosa</i>	Electroporation	Insertion of <i>NptII</i> gene and <i>eGFP</i>	[Run et al., 2016]
<i>C. sorokiniana</i>	Bioelectric	Insertion of the nitrate reductase ( <i>NR</i> ) gene into <i>NR</i> -deficient <i>C. sorokiniana</i> mutants	[Dawson et al., 1997]
<i>C. vulgaris</i>	Electroporation/ Glass bead agitation/ <i>Agrobacterium</i> -mediated,	Insertion of GFP*His6 or <i>aph7</i> gene	[Muñoz et al., 2018]
<i>C. sorokiniana</i> , <i>C. vulgaris</i> FSP-E	Electroporation/Plasmid/ Cas9	Deletion mutant (fatty acid desaturase- <i>fad3</i> gene)	[Lin and Ng, 2020]
<i>C. vulgaris</i> UTEX395	Electroporation/Plasmid, RNP/Cas9, Bombardment/ RNP/Cas9	Null mutation of edited genes of nitrate reductase ( <i>NR</i> ) and adenine Phosphoribosyltransferase ( <i>APT</i> )	[Kim et al., 2021]

(continued on next page)

**Table 3 (continued)**

Species/Strain	Transformation method	Gene mutants	References
<i>C. sorokiniana</i> UTEX 1602	Electroporation/Plasmid/Cas9/CRISPRi/ CRISPRa	Mutants of phosphoenolpyruvate carboxylase ( <i>PEPC</i> ) and glutamine synthetase ( <i>GS</i> ) genes	[Lin et al., 2022]
Other species	Transformation method	Gene mutants	References
<i>Euglena gracilis</i>	Microinjection/RNP/Cas9	Deletion mutants of the partial <i>EgGSL2</i> and <i>crtP1</i> gene	[Chen et al., 2022]
<i>E. gracilis</i>	Electroporation/RNP/Cas9	Deletion mutants of the <i>Bardet-Biedl syndrome</i> ( <i>BBS</i> ) genes	[Ishikawa et al., 2022]
<i>Picochlorum celeri</i>	Electroporation/RNP/Cas9	Loss-of-function mutants of nitrate reductase and carotenoid isomerase genes	[Krishnan et al., 2020]
<i>P. purpureum</i> CCMP 1328	Bombardment/RNP/Cas9	Chlorophyll synthase loss-of-function mutants ( $\Delta chs1$ )	[Jeon et al., 2021]
<i>Tetraselmis sp.</i> <i>KCTC12432BP</i>	Bombardment/RNP/Cas9	Development of <i>AGP</i> gene (ADP-glucose pyrophosphorylase) loss-of-function mutants	[Chang et al., 2020]
<i>B. braunii</i>	Electroporation	Insertion of the <i>AphVIII</i> gene	[Berrios et al., 2016]
<i>C. zofingiensis</i>	Biolistic	Insertion of <i>CHYb</i> ( $\beta$ -carotene hydroxylase) and <i>BKT</i> ( $\beta$ -carotene ketolase) genes	[Liu et al., 2014a, 2014b]
<i>Coccomyxa sp.</i>	Biolistic/ Electroporation	Overexpression of acyl-(acyl-carrier-protein) thioesterase gene, and a type-2 diacylglycerol acyltransferase gene; knock-out mutants of the <i>FTSY</i> gene	[Kasai et al., 2018; Yoshimitsu et al., 2018]
<i>C. subelliptoidea</i>	Electroporation	Insertion of the blue fluorescent protein ( <i>BFP</i> ) gene and <i>hptII</i> gene coding for hygromycin B phosphotransferase	[Kania et al., 2020]
<i>C. merolae</i>	PEG-mediated	double knock-in of HA-tagged cyclin 1 ( <i>CYC1/CML219C</i> ) and FLAG-tagged cyclin-dependent kinase A ( <i>CDKA/CME119C</i> ); Insertion of the antisense strand of the <i>catalase</i> gene; insertion of the acyl-ACP reductase gene of <i>Synechocystis</i> sp. PCC 6803 (sll0209)	[Fujiwara et al., 2017; Ohnuma et al., 2009; Sumiya et al., 2015]
<i>D. salina</i>	Electroporation/ Biolistic/ Glass beads agitation/ <i>Agrobacterium</i> -mediated	Insertion of nitrate reductase ( <i>NR</i> ) gene and bar gene; insertion of GFP as a reporter gene and, kanamycin and hygromycin resistant genes	[Li et al., 2007; Srinivasan and Gothandam, 2016]
<i>Fistulifera solaris</i>	Biolistic	Overexpression of the endogenous glycerol kinase ( <i>GK</i> ) gene	[Muto et al., 2015]
<i>Gonium pectorale</i>	Biolistic	Insertion of <i>aphVIII</i> gene and luciferase gene	[Lerche and Hallmann, 2009]
<i>H. pluvialis</i>	Biolistic	Mutation and up-regulation of the <i>pds</i> gene; insertion of <i>tub/aadA</i> , <i>tub/hyr</i> and <i>tub/ble</i>	[Steinbrenner and Sandmann, 2006; Yuan et al., 2019]
<i>M. neglectum</i>	Electroporation	Insertion of <i>aphVII</i> " gene	[Jaeger et al., 2017]
<i>O. tauri</i>	Electroporation/ PEG-based	Insertion of the aminoglycoside 30-phosphotransferase type I ( <i>APHI</i> ) coding gene of the bacterial transposon Tn903 for G418 resistance and the firefly luciferase CDS and expression assay by using four native <i>O. tauri</i> promoter sequences for histone 2 A ( <i>pOLK2</i> ), histone 3 ( <i>pOLK3</i> ), ubiquitin ( <i>pOLK4</i> ) and thioredoxin ( <i>pOLK5</i> ); Genomic transformation protocol	[Sanchez et al., 2019; van Ooijen et al., 2012]
<i>P. kessleri</i>	Biolistic/ <i>Agrobacterium</i> -mediated	Insertion of hygromycin phosphotransferase ( <i>hpt</i> ) gene and a <i>GUS</i> gene under the CaMV promoter	[Rathod et al., 2013]
<i>S. obliquus</i>	Electroporation	Insertion of CAT gene	[Guo et al., 2013]
<i>S. microadriaticum</i>	Silicon carbide whiskers	Insertion of neomycin phosphotransferase gene ( <i>nptII</i> ) and hygromycin B phosphotransferase gene ( <i>hpt</i> )	[Ten Lohuis and Miller, 1998]
<i>V. carteri f. magariensis</i>	Biolistic	Indel mutants of <i>glsA</i> , <i>regA</i> and <i>invA</i> genes	[Ortega-Escalante et al., 2019]

nutrient content, and stress tolerance (Ryool Jeong et al., 2023).

Another powerful method is the use of RNA interference (RNAi) to selectively inhibit the expression of specific genes in organisms. Microalgae can undergo gene silencing or knockdown through the introduction or expression of artificial microRNAs or by using RNA interference (RNAi) techniques (Cerutti et al., 2011; Kim and Cerutti, 2009; Molnar et al., 2009). In algae, RNAi has been employed to suppress the activity of target genes by introducing small RNA molecules that can specifically recognize and degrade the mRNA transcripts of those genes. Antisense technology and RNAi have been successfully applied in the model microalga *C. reinhardtii* as reported in various studies (Kim and Cerutti, 2009; Moellering and Benning, 2010; Molnar et al., 2009). Characterization of the microRNA (miRNA) system and its potential for gene regulation has been performed in several microalgae species including *C. reinhardtii*, the red alga *Porphyra yezoensis*, and the diatoms *T. pseudonana* and *Phaeodactylum tricornutum* (Huang et al., 2011; Molnar et al., 2009; Norden-krichmar et al., 2011; Zhao et al., 2007). A successful application of RNAi in algae is the suppression of genes involved in lipid degradation, resulting in increased lipid accumulation within cells, which is important for sustainable biofuel production from algae (Aratboni et al., 2019).

#### 4.1. DNA elements required for transformation

##### 4.1.1. Selectable marker genes and reporter genes

Various endogenous selective markers have been used in gene editing in microalgae along with the co-targeted gene of interest, such as the nitrate reductase, adenine phosphoribosyl transferase, peptidylprolyl isomerase, tryptophan synthase beta subunit, orotidine 5'-phosphate decarboxylase, protoporphyrinogen IX oxidase, and spermidine synthase (Freudenberg et al., 2022; Jiang and Weeks, 2017; Serif et al., 2018). These markers have been employed to facilitate gene modifications without the need for transgene integration, thereby improving the efficiency of the intended gene modifications in microalgae. In addition to the endogenous selective markers, antibiotic and herbicide resistance-based pre-selection methods have also been used and have been refined to achieve high selection efficiency up to 90% (Greiner et al., 2017; J. H. Kim et al., 2020; Picariello et al., 2020). These markers are derived from bacteria and, when introduced into microalgae, confer resistance to specific antibiotics or herbicides (León-Bañares et al., 2004).

Moreover, reporter genes, such as luciferase or mVenus have been successfully inserted at desired gene sites, enabling easy selection of mutant strains (Freudenberg et al., 2022; J. H. Kim et al., 2020). Reporter genes typically encode enzymes or fluorescent proteins that generate a visible or detectable signal, allowing the monitoring of the activity and expression of the gene of interest. Among the widely used reporter genes are the green fluorescent protein (GFP) and its variants,

such as blue (BFP), cyan (CFP), orange (OFP), red (RFP), and yellow (YFP) (Day and Davidson, 2009; Rasala et al., 2013). Reporter genes play a crucial role in genetic studies and biotechnological applications, by enabling the visualization and quantification of gene expression in real time and protein localization.

#### 4.1.2. Regulatory sequences

Regulatory sequences play a crucial role in gene expression as they control the time and site of gene expression, especially for both selectable marker and reporter genes (Sproles et al., 2021). In microalgae, it is often necessary to use endogenous regulatory sequences to ensure gene expression, as heterologous regulatory sequences may not be recognized or may result in minimal or no expression (Potvin and Zhang, 2010). As such, endogenous regulatory sequences are often replaced with different ones to modify gene expression patterns or increase expression rates. This can involve replacing the promoter region or using larger “regulatory regions” or “promoter regions” when the exact position of essential DNA elements is not known. Constitutive regulatory sequences, such as the small chain of ribulose bisphosphate carboxylase (RbcS2), heat shock protein 70 A (HSP70A), or  $\beta$ -tubulin, are commonly used when continuous expression is desired, such as for selectable marker genes (Potvin and Zhang, 2010; Strenkert et al., 2013). In contrast, inducible regulatory sequences are useful when controlling gene expression and turning it on or off at specific times or under specific conditions. These inducible regulatory sequences can be designed to respond to specific chemicals or light, allowing precise control over the gene expression.

In *C. reinhardtii*, >60 working regulatory sequences (promoters, UTRs, terminators, tags, reporters, antibiotic resistance genes, and introns) have been identified and used for gene expression (Crozet et al., 2018). Some of the most implemented regulatory sequences are small chain of ribulose bisphosphate carboxylase /oxygenase small subunit (RbcS2) (Stevens et al., 1996),  $\beta$ -tubulin (Blankenship and Kindle, 1992; Hallmann and Sumper, 1996), photosystem I complex proteins (psaD) (Fischer and Rochaix, 2001), cytochrome c6 (CYC6) (Quinn et al., 2003), nopaline synthase (NOS) (Díaz-Santos et al., 2013), and NAD(P)H nitrate reductase (nia1) (Llamas et al., 2002). For chloroplast transformation, regulatory sequences from genes, such as chlL, petD, rbcL, atpA, atpB, and rrn16 have been found to be suitable (Kim et al., 2015).

#### 4.1.3. UTRs and introns

The 5' and 3' untranslated regions (UTRs) of mRNA can impact the gene expression in microalgae. More specifically, the 3' UTRs affect the mRNA stability and polyadenylation. In *C. reinhardtii*, an effective polyadenylation signal such as UGUAA in the 3' UTR of genes can support expression (Bell et al., 2016). In the case of 3' UTRs, specific UTRs from genes such as RBCS2 or PSAD have been preferred in studies (Fischer and Rochaix, 2001; Lumbrales et al., 1998). On the other hand, the 5' UTRs contain introns and regulatory elements that could influence the transcriptional regulation (Specht et al., 2010). These regulatory elements within the intron sequences can affect gene expression by interacting with transcription factors (TF) or other regulatory proteins. The exact position and function of regulatory elements within the 5' UTRs may not always be known; therefore, larger promoter regions provide a better chance in capturing these important elements (Scranton et al., 2016).

Spliceosomal introns are a common feature in eukaryotic genomes, including microalgae, and hold important roles in gene expression regulation. While the number of introns per gene can vary greatly, it has been observed that the majority of microalgal genes contain at least one intron (Baier et al., 2018). Introns harbor various regulatory elements such as enhancers or silencers, transcription factor binding sites, and alternative splicing signals, which can influence the transcriptional activity of genes and affect the abundance, processing, stability, and translation efficiency of the resulting mRNA isoforms. Therefore, introns can be crucial for achieving an adequate expression of a gene of interest

in microalgae. Insertion of introns from the target species at unnatural positions within heterologous or homologous cDNAs is an alternative strategy for enhancing gene expression in microalgae (Baier et al., 2018; Eichler-Stahlberg et al., 2009; Hallmann et al., 1999; Lumbrales et al., 1998). The insertion of the intron of the ribulose-1,5-bisphosphate carboxylase/oxygenase small subunit 2 (rbcS2i1) throughout codon-optimized coding sequences (CDS) has been shown to generate optimized algal transgenes which have been expressed reliably in *C. reinhardtii* (Baier et al., 2018). Also, Baier et al. (2020) have identified several native and non-native introns in the green microalga *C. reinhardtii* with the ability to elicit strong Intron-Mediated Enhancement (IME) and were broadly applicable in different expression constructs.

#### 4.1.4. De novo DNA synthesis

*De novo* DNA synthesis involves the chemical synthesis of DNA in the laboratory without the need for DNA or RNA template. This process allows for the development of custom DNA sequences based on electronically available or designed sequences. Until recently, the upper limit for the length of chemically synthesized single-stranded DNA was around 200-300 nucleotides (Johnson and Widlanski, 2003; Palluk et al., 2018). But in 2021, first synthesis of synthesis of long (>10kb) DNA was reported (Hoose et al., 2023). In microalgae, large-scale oligonucleotide synthesis was applied for the creation of libraries of variant 5' UTRs of two *C. reinhardtii* genes, psbD and psaA, with the aim of identifying key regulators of chloroplast gene expression and developing tools for increased recombinant protein accumulation in algal plastids (Specht and Mayfield, 2013). In addition to endogenous regulatory sequences, synthetic regulatory sequences have also been developed that can drive robust gene expression in the nucleus of microalgae (Scranton et al., 2016). These synthetic regulatory sequences offer flexibility and control in gene expression for various biotechnological applications. Furthermore, DNA synthesis technology has enabled the synthesis of metabolic ORFeomes of two cyanobacteria strains of *Prochlorococcus marinus* (MED4, NATL1A) and *Chlamydomonas*, allowing for carrying out high throughput experiments with high levels of success rate (Fu et al., 2019). DNA synthesis is an indispensable tool of genetic engineering, allowing for the development of custom-designed DNA sequences (DNA molecules of desired composition and complexity) (Hoose et al., 2023) but its future application is highly affected by the cost and affordability (Patwari et al., 2023).

#### 4.1.5. Vector construction

Vectors used for transformation of microalgae often incorporate elements from both eukaryotic microalgae and bacterial vectors. The bacterial vector backbone, such as pBluescript, allows for a vector construction in *Escherichia coli* strains like DH5 $\alpha$ . The minimal requirements for propagation in *E. coli* include an origin of replication (e.g., pUC ori), an antibiotic resistance gene (e.g., AmpR), and a multiple cloning site for easy cloning (Rosales-Mendoza, 2016). One approach to simplify vector construction is the use of the  $\lambda$ -phage based gateway system, which utilizes recombination sequences for the transfer of DNA fragments between vectors *in vitro* (Hartley et al., 2000; Oey et al., 2014). Another strategy is the cassette multiplication technique, which enables the production of vectors with multiple gene copies, potentially leading to increased gene dosage and expression rates (Hallmann and Wodniok, 2006). Additionally, a modular cloning toolkit based on Golden Gate cloning with standard syntax has been developed for *C. reinhardtii*, containing various genetic parts for maximum modularity (Crozet et al., 2018). Transformation vectors have been designed to direct engineered gene products to different subcellular compartments, such as the chloroplast (Beth et al., 2014). Additionally, vectors that facilitate efficient secretion of recombinant proteins have been developed, which simplifies the purification process., for *C. reinhardtii* *P. tricornutum* (Vanier et al., 2017).

#### 4.2. Transformation methods

In parallel with selecting the most appropriate DNA elements required for transformation, various techniques have been developed to facilitate the uptake of foreign DNA into the cell and subsequently into the nucleus, chloroplast, or mitochondria of the algae, for ensuring higher yields of the targeted metabolite (lipids, antioxidants, high-value bioactive compounds, and proteins). Nevertheless, in contrast to the plant systems, the transformation of microalgae continues to face challenges in achieving high efficiency, with the exception of *Chlamydomonas* (Kumar et al., 2020).

The primary challenge in accomplishing successful microalgae transformation is overcoming the cell wall to allow the entry of DNA, and as such, most protocols include a physical method to penetrate the cell wall (Ortiz-Matamoros et al., 2018). Integration of exogenous genes into the genome typically occurs at random sites through a process called recombination, which does not depend on homology or specific DNA sequences, albeit regions of microhomology may play a role. Technical limitations in the generation of transgenic lines in microalgae.

The most commonly used transformation techniques employed in algal genetic engineering are presented below and examples can also be found in Table 3.

Electroporation involves the use of high-intensity electric pulses to create temporary pores in the cell membrane allowing foreign DNA to enter the cell walls and membrane's phospholipid bilayer (Rathod et al., 2017). This technique has been used successfully to transform many algae species, including *C. reinhardtii*, *. salina*, *Cyanidioschyzon merolae*, *Nannochloropsis* sp., *C. vulgaris*, *Chlorella ellipsoidea*, and *Chlorella saccharophila* (Z. Chen and Lee, 2019; Garoma and Shackelford, 2014; Ladygin, 2003). In some cases, enzymatic pretreatment is required to disrupt the cell wall; however, this method has been successful only in some cases given the limited available information regarding the diverse composition of cell walls in different microalgae species. Nevertheless, this method is particularly effective for nuclear gene transformation in microalgae compared to other systems (Fajardo et al., 2020a, 2020b).

Microinjection method has been used to directly inject foreign DNA into the nucleus of the algal cell using a fine-tipped glass needle. This technique is highly precise and can be used to introduce DNA into specific cells within a population (Qin et al., 2012); yet it is labor-intensive and time-consuming.

Microparticle bombardment (biolistics) is a versatile method that enables transformation of various cell types regardless of their cell wall thickness or rigidity, as well as plastid transformation. This method was successfully used in microalgae to transform chloroplast genome and in studies for the production of different compounds (Gangl et al., 2015). This technique has been successfully applied in various microalgae species. These include *C. reinhardtii*, *V. carteri*, *Gonium*, *Pandorina morum*, *Eudorina elegans*, *H. pluvialis*, *Euglena gracilis*, *Porphyridium* sp., 'Chlorella' *kessleri*, *C. sorokiniana* and *D. salina* (Lapidot et al., 2002; Siddiqui et al., 2020). Diatoms such as *P. tricornutum*, *Cylindrotheca fusiformis*, *Navicula saprophila*, and *Cyclotella cryptica* can also undergo transformation through particle bombardment (Velmurugan and Deka, 2018). Nevertheless, the transformation efficiency is relatively low (Ng et al., 2017).

Agitation with glass beads is a popular transformation method due to its simplicity and reproducibility, especially for species that lack rigid cell wall, such as *D. salina* (Feng et al., 2009), however low transformation efficiencies and reduced cell recovery are some of the drawbacks (Ortiz-Matamoros et al., 2018). This method has also been applied successfully to *C. reinhardtii* (Wannathong et al., 2016) and *Symbiodinium* spp. (Ortiz-Matamoros et al., 2015a, 2015b). The protocol involves an enzymatic pre-treatment for the degradation of the eukaryotic cell wall and vigorous agitation of algal cells/DNA suspension with glass beads and the membrane fusion agent polyethylene glycol (PEG) (Ortiz-Matamoros et al., 2018).

Similar to higher plants, microalgae have also been successfully

transformed using *Agrobacterium tumefaciens*. *Agrobacterium* possesses a tumor-inducing (Ti) plasmid that integrates into the genome of infected cells. *Agrobacterium*-mediated transformation has been used in microalgae species, such as *Chlamydomonas reinhardtii*, *H. pluvialis*, *D. bardawil*, *Parachlorella kessleri*, *Schizochytrium* sp and *Symbiodinium* spp (Anila et al., 2011; R. Cheng et al., 2012; Kathiresan et al., 2009; Kathiresan and Sarada, 2009; Kumar et al., 2004; Ortiz-Matamoros et al., 2015a, 2015b; Rathod et al., 2013).

Interestingly, transformation methods, such as electroporation, microinjection, biolistics and microfluidic techniques, used as physical delivery techniques coupled with the use of lipid nanoparticles, polymers, nanogels, inorganic nanoparticles, and cell-derived vesicles as synthetic carriers have been, thus far, successfully employed for enhancing the efficiency of genome editing using the Cas9 ribonucleoprotein (RNP) (Kang et al., 2020; Lu et al., 2021).

As interest grows in developing transgenic microalgae for biotechnological applications and research, overcoming issues related to transgene efficiency, integration, or stability are common is a major challenge. Selecting the most appropriate transformation strategy is crucial for ensuring a successful transformation using approaches specifically tailored to the diverse cellular characteristics of microalga. The main challenge lies in permeabilizing the cell walls and membranes to introduce DNA without causing significant damage that could harm or kill the cells. Other important factors for enhancing the transformation efficiency include the optimization of the transformation conditions, the type of genetic material to be transferred and that the cells can resume normal cell division after genetic modification.

#### 5. Applications of genetic engineering and omics tools in microalgae

The field of microalgae biotechnology is gaining increasing commercial significance across a wide range of industries including food, pharmaceuticals, nutraceuticals, cosmetics, animal feed, energy and can also be employed in environmental applications, such as water and soil remediation (Singh et al., 2021; Vanhoudt et al., 2018). Whilst applications of genetically engineered microalgae have also been found in other scientific fields, such as in neuroscience, where the light-dependent ion channels of microalgae allow for the manipulation of the neuronal activity (Ishizuka et al., 2006). The genetic modification of microalgae offers expanded possibilities for biotechnological applications in various contexts through enabling the production of compounds and introduction of traits that are not naturally present in algae. Various studies refer to molecular farming, molecular pharming, biopharming, or gene pharming, that involves the production of enzymes, antibodies, immunotoxins, bioactive peptides, hormones, and vaccines (Dyo and Purton, 2018; Rasala and Mayfield, 2015; Taunt et al., 2018).

##### 5.1. Enhanced productivity of high-added value products and improved growth rates

Genetic engineering techniques can be employed to enhance the productivity of algae by manipulating their metabolic pathways. One approach is to overexpress specific enzymes that play a crucial role in lipid biosynthesis (Mulgund, 2022). By increasing the expression of these key enzymes, the lipid content of algae can be augmented. This is particularly advantageous in the production of biofuels given that lipids are a valuable raw material. Through genetic modifications, algae can be engineered to have higher lipid content, and thus be more efficient and economically viable for biofuel production. Ultimately, the enhanced lipid content contributes to the development of sustainable and renewable energy sources (Ng et al., 2017).

Through genetic engineering, the bioproduction of pharmaceutical and nutraceutical compounds in algae can be optimized rendering them a more viable and sustainable source for the development of novel therapies (Garduño-González et al., 2022; Qiao et al., 2020).

Overexpressing key enzymes in biosynthetic pathways can enhance the algae's capacity towards higher yields of the desired compounds, involving manipulation of genes for precursor synthesis, enzyme regulation, and other pertinent metabolic processes (Garduño-González et al., 2022; Qiao et al., 2020).

Genetic engineering techniques offer a means to enhance the growth rates of algae by manipulating their photosynthetic machinery. Additionally, the overall productivity and growth rates of algae can be significantly enhanced by optimizing the photosynthetic machinery through overexpressing essential genes involved in photosynthesis (Hitchcock et al., 2022).

## 5.2. Bioremediation

Algae have shown great potential for environmental remediation based on their ability to remove pollutants, such as heavy metals and organic compounds from the environment. Genetic engineering techniques can be employed to enhance this capability further. By over-expressing key enzymes involved in the degradation of pollutants, algae can be genetically modified to increase their efficiency in bioremediation processes. These enzymes play a crucial role in the breakdown and metabolism of specific pollutants, facilitating their removal or transformation into less harmful forms. Through genetic engineering, the expression of these enzymes can be upregulated, enabling the algae to effectively target and degrade pollutants in the environment (Martínez-Ruiz et al., 2022; Zeraatkar et al., 2016). This approach offers a promising avenue for the development of sustainable and eco-friendly strategies to address pollution challenges.

## 6. Impact of genetic engineering and omics data on microalgae industry

The integration of omics data and metabolic engineering allows researchers to design and implement precision modifications in microalgae strains for specific biotechnological applications, ranging from biofuel production and pharmaceuticals to environmental remediation. Overall, genetic engineering of algae involves the manipulation of the genetic material to achieve specific characteristics with notable implications for the algae industry, offering both advantages and potential drawbacks. While it can increase productivity, enhance stress tolerance, and improve the quality of algae products, it necessitates a careful consideration of potential risks and ethical implications. More specifically, one of the positive effects is the improvement on growth rate and overall productivity resulting in higher biomass yields, biofuels, or other high-added value microalgae derived products. It is also used effectively to optimize the composition of lipids, proteins, or other compounds with commercial or industrial value. Additionally, the development of resilient microalgae strains may not only contribute to the effective cultivation in adverse conditions but also in producing stress-induced high-added value products. Negative aspects of genetic engineering include the release of transgenic strains in the environment that could affect the natural diversity and possibly disrupt the balance of ecosystems, the production of algae-derived products that may not be suitable for human consumption. Therefore, there is a need for careful consideration of the benefits and risks of genetic engineering of algae, and for the development of regulations and guidelines that ensure its safe and sustainable use. Comprehensive risk assessments, ethical frameworks, and robust regulatory mechanisms are essential to guide the responsible deployment of genetically modified microalga.

The CRISPR/Cas technology is a potent genetic engineering tool with numerous potential applications in microalgae. It offers the opportunity to introduce specific genetic modifications that can enhance algae's productivity, improve biofuel yields, increase disease resistance, and promote carbon sequestration. However, there are still obstacles and limitations that must be addressed, including the development of efficient delivery methods, the potential for off-target effects, and safety

concerns. Through ongoing research and development, CRISPR/Cas has the capacity to revolutionize algae biotechnology, offering sustainable solutions to a range of environmental challenges. It is essential to carefully weigh the benefits and risks associated with genetic engineering of algae to ensure its safe and sustainable use.

Despite the significant advancements in the field, the current knowledge on the metabolism of industrially relevant microalgae strains remains limited. A substantial portion of genes lack significant sequence similarity to known entities, making it challenging to assign them specific metabolic functions (Vieler et al., 2012). Additionally, uncertainties persist regarding protein localization, compartmentalization of metabolite pools, and trafficking of metabolites between organelles, further complicating the development of comprehensive metabolic models. These limitations must be addressed given that a deeper understanding of microalgae and their metabolism is crucial for guiding the design of strain engineering approaches. In the absence of high-quality metabolic models, genetic engineering strategies are often constrained to educated guesses based on knowledge obtained from other organisms.

The advancement of genetic tools and omic technologies has significantly accelerated research in a select few microalgae strains, leading to the accumulation of a large database of transcriptomic responses to various stimuli (Südfeld et al., 2021; Zienkiewicz et al., 2016). To enable metabolic engineering, it is crucial to have an advanced toolbox for genetic manipulation. However, developing genome editing methods specifically tailored for microalgae species is a time-consuming process that requires careful attention. Currently, the genetic toolbox for industrially relevant microalgae is relatively limited compared to other microbial systems, but efforts are underway to expand and enhance these tools.

## 7. Conclusions

Microalgal omics hold considerable promise due to the ongoing breakthroughs in genomics technologies, such as better sequencing platforms, bioinformatics tools, and data processing techniques. Transcriptional and metabolic engineering strategies have played vital roles in advancing microalgae research. These approaches have facilitated the functional characterization of key genes and regulators, the identification of metabolic pathways, and the elucidation of microalgae cell physiology (Bajhaiya et al., 2017). Moreover, the use of knowledge on epigenetic and metabolic cellular regulating processes may facilitate the confirmation of regulatory protein functions and their impact on diverse metabolic pathways and will contribute to the development of robust microalgal strains (Bacova et al., 2020). The rapid advancement of omics technologies in the field of microalgae have laid a strong foundation for genetic engineering. Extensive research conducted over several decades, especially in the model algae such as Chlamydomonas, has elucidated gene expression patterns, biochemical processes, and metabolic pathways in microalgae (Jeong et al., 2023). Overall, the review provides a comprehensive overview of the current state of microalgae research, emphasizing the synergistic integration of omics and metabolic engineering approaches to drive advancements in strain improvement for various practical and sustainable applications. Future research is required with a biology systems approach by integrating the pan-omics data of microalgae research to enable us to predict novel functional interactions between molecular mediators at multiple levels and identify points of metabolic flux for improving algal strain-engineering techniques.

## Funding information

This project has received funding from the European Union's Horizon 2020, under the Marie Skłodowska-Curie grant agreement No 778263 – AlgaeCeuticals, The Structure of Research, Innovation and Excellence (DEKA) of the University of Thessaly for the New Professors and the

scholarship to new PhD candidates (grant numbers 5600.03.08.04 and 5600.03.06.03) and the Marie Skłodowska-Curie grant agreement No 101086437 – AlgaeNet4AV.

### CRediT authorship contribution statement

**Evangelia Stavridou:** Writing – review & editing, Writing – original draft. **Lefkothhea Karapetsi:** Writing – review & editing. **Georgia Maria Nteve:** Writing – review & editing. **Georgia Tsintzou:** Writing – review & editing. **Marianna Chatzikonstantinou:** Writing – review & editing. **Meropi Tsaousi:** Writing – review & editing. **Angel Martinez:** Writing – review & editing. **Pablo Flores:** Writing – review & editing. **Marián Merino:** Writing – review & editing. **Luka Dobrovic:** Writing – review & editing. **José Luis Mullor:** Writing – review & editing. **Stefan Martens:** Writing – review & editing. **Leonardo Cerasino:** Writing – review & editing. **Nico Salmaso:** Writing – review & editing. **Maslin Osathanunkul:** Writing – review & editing. **Nikolaos E. Labrou:** Writing – review & editing, Writing – original draft. **Panagiotis Madesis:** Writing – review & editing, Writing – original draft, Funding acquisition.

### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### Data availability

No data was used for the research described in the article.

### References

- Abdel-Ghany, S.E., Golovkin, M., Reddy, A.S.N., 2015. Engineering of plants for the production of commercially important products: approaches and accomplishments. In: *In Plant Biology and Biotechnology: Volume II: Plant Genomics and Biotechnology*, Vol. II. [https://doi.org/10.1007/978-81-322-2283-5\\_28](https://doi.org/10.1007/978-81-322-2283-5_28).
- Abdellaoui, N., Jeong, M., Tae, K., Choi, J., 2019. Transcriptome analysis of gene expression in *Chlorella vulgaris* under salt stress. *World J. Microbiol. Biotechnol.* 35 (9), 1–11. <https://doi.org/10.1007/s11274-019-2718-6>.
- Abu-Ghosh, S., Dubinsky, Z., Verdelho, V., Iluz, D., 2021. Unconventional high-value products from microalgae: A review. *Bioresour. Technol.* 329 (February), 124895. <https://doi.org/10.1016/j.biortech.2021.124895>.
- Afifudeen, C.L.W., Aziz, A., Lian, L., Takahashi, K., Toda, T., Effendi, M., Wahid, A., San, T., 2021. Phytochemistry transcriptome-wide study in the green microalga *Messastrum gracile* SE-MC4 identifies prominent roles of photosynthetic integral membrane protein genes during exponential growth stage. *Phytochemistry* 192 (August), 112936. <https://doi.org/10.1016/j.phytochem.2021.112936>.
- Ahmad, A., Banat, F., Alsafar, H., Hasan, S.W., 2022. Recent breakthroughs in integrated biomolecular and biotechnological approaches for enhanced lipid and carotenoid production from microalgae. *Phytochem. Rev.* 5 <https://doi.org/10.1007/s11101-022-09804-5>.
- Ajaiji, I., Verruto, J., Aqui, M., Soriaga, L.B., Coppersmith, J., Kwok, K., Peach, L., Orchard, E., Kalb, R., Xu, W., Carlson, T.J., Francis, K., Konigsfeld, K., Bartalis, J., Schultz, A., Lambert, W., Schwartz, A.S., Brown, R., Moellerling, E.R., 2017. Lipid production in *Nannochloropsis gaditana* is doubled by decreasing expression of a single transcriptional regulator. *Nat. Biotechnol.* 35 (7), 647–652. <https://doi.org/10.1038/nbt.3865>.
- Akella, S., Ma, X., Bacova, R., Harmer, Z.P., Kolackova, M., Wen, X., Wright, D.A., Spalding, M.H., Weeks, D.P., Cerutti, H., 2021. Co-targeting strategy for precise, scarless gene editing with CRISPR/Cas9 and donor ssODNs in *Chlamydomonas*. *Plant Physiol.* 187 (4), 2637–2655. <https://doi.org/10.1093/plphys/kia418>.
- Al-Eyalawi, Z.A., Al-Dulaimy, A.F.Z., 2023. Marine algae and applications to plant nutrition: a review. *IOP Conf. Series: Earth Environ. Sci.* 1158 (4), 042004. <https://doi.org/10.1088/1755-1315/1158/4/042004>.
- Angstenberger, M., De Signori, F., Vecchi, V., Dall’Osto, L., Bassi, R., 2020. Cell synchronization enhances nuclear transformation and genome editing via Cas9 enabling homologous recombination in *Chlamydomonas reinhardtii*. *ACS Synth. Biol.* 9 (10), 2840–2850. <https://doi.org/10.1021/acssynbio.0c00390>.
- Anila, N., Chandrashekhar, A., Ravishankar, G.A., Sarada, R., 2011. Establishment of *Agrobacterium tumefaciens* – mediated genetic transformation in *Dunaliella bardawil*. *Eur. J. Phycol.* 0262 <https://doi.org/10.1080/09670262.2010.550386>.
- Aranda, M., Li, Y., Liew, Y.J., Baumgarten, S., Sim, O., Wilson, M.C., Piel, J., Ashoor, H., 2016. Genomes of Coral Dinoflagellate Symbionts Highlight Evolutionary Adaptations Conducive to a Symbiotic Lifestyle. *Nature Publishing Group*, November, pp. 1–15. <https://doi.org/10.1038/srep39734>.
- Aratboni, A., Cell, M., Aratboni, H.A., Rafiei, N., Granados, R.G., Alemzadeh, A., 2019. Biomass and lipid induction strategies in microalgae for biofuel production and other applications. *Microb. Cell Factories* 1–17. <https://doi.org/10.1186/s12934-019-1228-4>.
- Armburst, E.V., Armbrust, E.V., Berges, J.A., Bowler, C., Green, B.R., Martinez, D., Putnam, N.H., Zhou, S., Allen, A.E., Apt, K.E., Bechner, M., Brzezinski, M.A., Chaal, B.K., Chiovitti, A., Glavina, T., Goodstein, D., Hadi, M.Z., 2012. The genome of the diatom, *79* (2004). <https://doi.org/10.1126/science.1101156>.
- Ashworth, J., Ralph, P.J., 2018. An exploratory public transcriptomics compendium for eukaryotic microalgae. *BioRxiv* 403063. <https://www.biorxiv.org/content/10.1101/403063v3.abstract>.
- Azaman, S.N.A., Wong, D.C.J., Tan, S.W., Yusoff, F.M., Nagao, N., Yeap, S.K., 2020. De novo transcriptome analysis of *Chlorella sorokiniana*: effect of glucose assimilation, and moderate light intensity. *Sci. Rep.* 10 (1), 1–12. <https://doi.org/10.1038/s41598-020-74410-4>.
- Bachy, C., Charlesworth, C.J., Chan, A.M., Finke, J.F., Wong, C., Wei, C., Sudek, S., Coleman, M.L., Suttle, C.A., Worden, A.Z., 2018. Transcriptional responses of the marine green alga *Micromonas pusilla* and an infecting prasinovirus under different phosphorus conditions, 20, pp. 2898–2912. <https://doi.org/10.1111/1462-2920.14273>.
- Bacova, R., Kolackova, M., Klejdus, B., Adam, V., Huska, D., 2020. Epigenetic mechanisms leading to genetic flexibility during abiotic stress responses in microalgae: A review. *Algal Res.* 50 (July 2019), 101999. <https://doi.org/10.1016/j.algal.2020.101999>.
- Baier, T., Wichmann, J., Kruse, O., Lauersen, K.J., 2018. Intron-containing algal transgenes mediate efficient recombinant gene expression in the green microalga *Chlamydomonas reinhardtii*, 46 (13), 6909–6919. <https://doi.org/10.1093/nar/gky532>.
- Baier, T., Jacobebbinghaus, N., Einhaus, A., Lauersen, K.J., Kruse, O., 2020. Introns mediate post-transcriptional enhancement of nuclear gene expression in the green microalga *Chlamydomonas reinhardtii*. *PLoS Genet.* 16 (7), 1–21. <https://doi.org/10.1371/journal.pgen.1008944>.
- Bajhaiya, A.K., Ziehe Moreira, J., Pittman, J.K., 2017. Transcriptional engineering of microalgae: prospects for high-value chemicals. *Trends Biotechnol.* 35 (2), 95–99. <https://doi.org/10.1016/j.tibtech.2016.06.001>.
- Barten, R., Van Workum, D.J.M., De Bakker, E., Risso, J., Kleisman, M., Navalho, S., Smit, S., Wijffels, R.H., Nijveen, H., Barbosa, M.J., 2022. Genetic mechanisms underlying increased microalgal thermostolerance, maximal growth rate, and yield on light following adaptive laboratory evolution. *BMC Biol.* 1–14. <https://doi.org/10.1186/s12915-022-01431-y>.
- Beauvais-Flück, R., Slaveykova, V.I., Cosio, C., 2017. Cellular toxicity pathways of inorganic and methyl mercury in the green microalga *Chlamydomonas reinhardtii*. *Sci. Rep.* 7 (1), 1–12. <https://doi.org/10.1038/s41598-017-08515-8>.
- Beauvais-Flück, R., Slaveykova, V.I., Cosio, C., 2019. Comparative study of Cu uptake and early transcriptome responses in the green microalga *Chlamydomonas reinhardtii* and the macrophyte *Elodea nuttallii*. *Environ. Pollut.* 250, 331–337. <https://doi.org/10.1016/j.envpol.2019.04.032>.
- Bell, S.A., Shen, C., Brown, A., Hunt, A.G., 2016. Experimental genome-wide determination of RNA polyadenylation in *Chlamydomonas reinhardtii*. *PLoS One* 11 (1). <https://doi.org/10.1371/journal.pone.0146107>.
- Benner, I., Diner, R.E., Lefebvre, S.C., Li, D., Komada, T., Carpenter, E.J., Stillman, J.H., Benner, I., Stillman, J.H., 2013. *Emiliania huxleyi* increases calcification but not expression of calcification-related genes in long-term exposure to elevated temperature and p CO<sub>2</sub>.
- Berrios, H., Zapata, M., Rivas, M., 2016. A method for genetic transformation of *Botryococcus braunii* using a cellulase pretreatment. *J. Appl. Phycol.* 28 (1), 201–208. <https://doi.org/10.1007/s10811-015-0596-3>.
- Beth, A., Rasala, B.A., Chao, S., Pier, M., Barrera, D.J., Mayfield, S.P., 2014. Enhanced Genetic Tools for Engineering Multigene Traits into Green Algae. <https://doi.org/10.1371/journal.pone.0094028>.
- Bhattacharya, D., Price, D., Chan, C., et al., 2013. Genome of the red alga *Porphyridium purpureum*. *Nat Commun* 4, 1941. <https://doi.org/10.1038/ncomms2931>.
- Bibi, M., Zhu, X., Munir, M., Angelidakis, I., 2021. Bioavailability and effect of α-Fe2O<sub>3</sub> nanoparticles on growth, fatty acid composition and morphological indices of *Chlorella vulgaris*. *Chemosphere* 282 (March). <https://doi.org/10.1016/j.chemosphere.2021.131044>.
- Blanc-Mathieu, R., Verhelst, B., Derelle, E., Rombauts, S., Bouget, F.Y., Carré, I., Château, A., Eyre-Walker, A., Grimsley, N., Moreau, H., Piégay, B., Rivals, E., Schackwitz, W., Van de Peer, Y., Piganeau, G., 2014. An improved genome of the model marine alga *Ostreococcus tauri* unfolds by assessing Illumina de novo assemblies. *BMC Genomics* 15 (1). <https://doi.org/10.1186/1471-2164-15-1103>.
- Blankenship, J.E., Kindle, K.L., 1992. Expression of chimeric genes by the light-regulated cabII-1 promoter in *Chlamydomonas reinhardtii*: a cabII-1/nit1 gene functions as a dominant selectable marker in a nit1- nit2 strain. *Mol. Cell. Biol.* 12 (11), 5268–5279. <https://doi.org/10.1128/mcb.12.11.5268>.
- Brar, A., Kumar, M., Soni, T., Vivekanand, V., Pareek, N., 2021. Insights into the genetic and metabolic engineering approaches to enhance the competence of microalgae as biofuel resource: a review. *Bioresour. Technol.* 339 (July), 125597. <https://doi.org/10.1016/j.biortech.2021.125597>.
- Brawley, S.H., Blouin, N.A., Ficko-Blean, E., Wheeler, G.L., Lohr, M., Goodson, H.V., Jenkins, J.W., Blaby-Haas, C.E., Helliwell, K.E., Chan, C.X., Marriage, T.N., Bhattacharya, D., Klein, A.S., Badis, Y., Brodie, J., Cao, Y., Collén, J., Dittami, S.M., Gachon, C.M.M., Prochnik, S.E., 2017. Insights into the red algae and eukaryotic evolution from the genome of *Porphyra umbilicalis* (Bangiophyceae, Rhodophyta). *Proc. Natl. Acad. Sci. USA* 114 (31), E6361–E6370. <https://doi.org/10.1073/pnas.1703088114>.

- Breuer, G., Lamers, P.P., Martens, D.E., Draaisma, R.B., Wijffels, R.H., 2013. Bioresource technology effect of light intensity, pH, and temperature on triacylglycerol (TAG) accumulation induced by nitrogen starvation in *Scenedesmus obliquus*. *Bioresour. Technol.* 143, 1–9. <https://doi.org/10.1016/j.biortech.2013.05.105>.
- Browne, D.R., Jenkins, J., Schmutz, J., Shu, S., Barry, K., Grimwood, J., Chiniquy, J., Sharma, A., Niehaus, T.D., Weiss, T.L., et al., 2017. Draft nuclear genome sequence of the liquid hydrocarbon-accumulating green microalga *Botryococcus braunii* Race B (Showa). *Genome Announc.* 5, 17–19. <https://doi.org/10.1128/genomeA.00215-17>.
- Carney, L.T., Reinsch, S.S., Lane, P.D., Solberg, O.D., Jansen, L.S., Williams, K.P., Trent, J.D., Lane, T.W., 2014. Microbiome analysis of a microalgal mass culture growing in municipal wastewater in a prototype OMEGA photobioreactor. *Algal Res.* 4 (1), 52–61. <https://doi.org/10.1016/j.algal.2013.11.006>.
- Carvalho, A.P., Malcata, F.X., 2005. Optimization of omega-3 fatty acid production by microalgae: crossover effects of CO<sub>2</sub> and light intensity under batch and continuous cultivation modes. *Mar Biotechnol (NY)* 7 (4), 381–388. <https://doi.org/10.1007/s10126-004-4047-4>.
- Casabianca, S., Cornetti, L., Capellacci, S., Vernesi, C., Penna, A., 2017. Genome complexity of harmful microalgae. *Harmful Algae* 63, 7–12. <https://doi.org/10.1016/j.hal.2017.01.003>.
- Cazzaniga, S., Kim, M., Bellamoli, F., Jeong, J., Lee, S., Perozeni, F., Pompa, A., Jin, E.S., Ballottari, M., 2020. Photosystem II antenna complexes CP26 and CP29 are essential for nonphotochemical quenching in *Chlamydomonas reinhardtii*. *Plant Cell Environ.* 43 (2), 496–509. <https://doi.org/10.1111/pce.13680>.
- Cerutti, H., Ma, X., Msanne, J., Repas, T., 2011. RNA-mediated silencing in algae: biological roles and tools for analysis of gene function. *Eukaryot. Cell* 10 (9), 1164–1172. <https://doi.org/10.1128/EC.05106-11>.
- Chang, K.S., Kim, J., Park, H., Hong, S.J., Lee, C.G., Jin, E.S., 2020. Enhanced lipid productivity in AGP knockout marine microalga *Tetraselmis sp.* using a DNA-free CRISPR-Cas9 RNP method. *Bioresour. Technol.* 303 (January), 122932 <https://doi.org/10.1016/j.biortech.2020.122932>.
- Chang, M., Zhang, T., Li, L., Lou, F., Ma, M., Liu, R., Jin, Q., Wang, X., 2021. Choreography of multiple omics reveals the mechanism of lipid turnover in *Schizochytrium sp.* S31. *Algal Res.* 54 (December 2020), 102182 <https://doi.org/10.1016/j.algal.2021.102182>.
- Chen, Y., Hu, H., 2019. High efficiency transformation by electroporation of the freshwater alga *Nannochloropsis limnetica*. *World J. Microbiol. Biotechnol.* 35 (8), 1–10. <https://doi.org/10.1007/s11274-019-2695-9>.
- Chen, Z., Lee, W.G., 2019. Electroporation for microalgal biofuels: A review. *Sustain. Energy Fuels* 3 (11), 2954–2967. <https://doi.org/10.1039/c9se00087a>.
- Chen, Z., Zhu, J., Chen, Z., Du, M., Yao, R., Fu, W., Lei, A., Wang, J., 2022. High-throughput sequencing revealed low-efficacy genome editing using Cas9 RNPs electroporation and single-celled microinjection provided an alternative to deliver CRISPR reagents into *Euglena gracilis*. *Plant Biotechnol. J.* 20 (11), 2048–2050. <https://doi.org/10.1111/pbi.13915>.
- Cheng, R., Ma, R., Li, K., Rong, H., Lin, X., Wang, Z., 2012. *Agrobacterium tumefaciens* mediated transformation of marine microalgae *Schizochytrium*. *Microbiol. Res.* 167, 179–186. <https://doi.org/10.1016/j.micres.2011.05.003>.
- Cheng, J., Li, K., Yang, Z., Lu, H., Zhou, J., Cen, K., 2016. Gradient domestication of *Haematococcus pluvialis* mutant with 15% CO<sub>2</sub> to promote biomass growth and astaxanthin yield. *Bioresour. Technol.* 216, 340–344. <https://doi.org/10.1016/j.biortech.2016.05.095>.
- Cheng, P., Zhou, C., Wang, Y., Xu, Z., Xu, J., Zhou, D., Zhang, Y., Wu, H., Zhang, X., Liu, T., Tang, M., Yang, Q., Yan, X., Fan, J., 2018. Biotechnology for biofuels comparative transcriptome analyses of oleaginous *Botryococcus braunii* race A reveal significant differences in gene expression upon cobalt enrichment. *Biotechnol. Biofuels* 1–19. <https://doi.org/10.1186/s13068-018-1331-5>.
- Chernobai, N., Kadnikova, N., Vozovik, K., Rozanov, L., 2022. Study of Osmotic Behavior of *Asystasia longa* and *Dunalieilla salina* Cells as an Important Stage in Developing their Cryopreservation Protocol, 59. <https://doi.org/10.3390/iecp2021-11939>.
- Cho, K., Lee, C.H., Ko, K., Lee, Y.J., Kim, K.N., Kim, M.K., Chung, Y.H., Kim, D., Yeo, I.K., Oda, T., 2016. Use of phenol-induced oxidative stress acclimation to stimulate cell growth and biodiesel production by the oceanic microalga *Dunalieilla salina*. *Algal Res.* 17, 61–66. <https://doi.org/10.1016/j.ALGAL.2016.04.023>.
- Chung, Y.S., Lee, J.W., Chung, C.H., 2017. Molecular challenges in microalgae towards cost-effective production of quality biodiesels. *Renew. Sust. Energ. Rev.* 74 (March 2016), 139–144. <https://doi.org/10.1016/j.rser.2017.02.048>.
- Clement, R., Lignon, S., Mansuelle, P., Jensen, E., Pophillat, M., Lebrun, R., Denis, Y., Puppo, C., Maberly, S.C., Gontero, B., 2017. Responses of the marine diatom *Thalassiosira pseudonana* to changes in CO<sub>2</sub> concentration: a proteomic approach. *Nat. Publ. Group* 1–12. <https://doi.org/10.1038/srep42333>. July 2016.
- Collén, J., Porcel, B., Carré, W., Ball, S.G., Chaparro, C., Tomon, T., Barbevion, T., Michel, G., Noel, B., Valentin, K., Elias, M., Artiguenave, François, Arun, A., Aury, J. M., Barbosa-Neto, J.F., Bothwell, J.H., Bouget, F.Y., Brillet, L., Cabello-Hurtado, F., Boyen, C., 2013. Genome structure and metabolic features in the red seaweed *Chondrus crispus* shed light on evolution of the Archaeplastida. *Proc. Natl. Acad. Sci. USA* 110 (13), 5247–5252. <https://doi.org/10.1073/pnas.1221259110>.
- Coulombier, N., Jauffrais, T., Lebouvier, N., 2021. Antioxidant compounds from microalgae: A review. *Mar. Drugs* 19 (10). <https://doi.org/10.3390/md19100549>.
- Crozet, P., Navarro, F.J., Willmund, F., Mehrshahi, P., Bakowski, K., Lauersen, K.J., Pérez-Pérez, M.E., Auroy, P., Gorchs Rovira, A., Sauret-Gueto, S., Niemeyer, J., Spaniol, B., Theis, J., Trösch, R., Westrich, L.D., Vavitsas, K., Baier, T., Hübner, W., De Carpenter, F., Lemaire, S.D., 2018. Birth of a photosynthetic chassis: A MoClo toolkit enabling synthetic biology in the microalga *Chlamydomonas reinhardtii*. *ACS Synth. Biol.* 7 (9), 2074–2086. <https://doi.org/10.1021/acssynbio.8b00251>.
- da Silva, M.E.T., Leal, M.A., de Resende, M.O., Martins, M.A., dos Coimbra, J.S.R., 2021. *Scenedesmus obliquus* protein concentrate: A sustainable alternative emulsifier for the food industry. *Algal Res.* 59 (March) <https://doi.org/10.1016/j.algal.2021.102468>.
- Dagan, T., Roettger, M., Stucken, K., Landan, G., Koch, R., Major, P., Gould, S.B., Goremykin, V.V., Rippka, R., De Marsac, N.T., Gugger, M., Lockhart, P.J., Allen, J.F., Brune, I., Maus, I., Pühler, A., Martin, W.F., 2013. Genomes of stigonematalean cyanobacteria (subsection V) and the evolution of oxygenic photosynthesis from prokaryotes to plastids. *Genome Biol. Evol.* 5 (1), 31–44. <https://doi.org/10.1093/gbe/evs117>.
- Dahlin, L.R., Gerritsen, A.T., Henard, C.A., Van Wychen, S., Linger, J.G., Kunde, Y., Hovde, B.T., Starkenburg, S.R., Posewitz, M.C., Guarneri, M.T., 2019. Development of a high-productivity, halophilic, thermotolerant microalga *Picochlorum renovum*. *Commun. Biol.* 2 (1), 1–9. <https://doi.org/10.1038/s42003-019-0620-2>.
- Dawson, H.N., Burlingame, R., Cannons, A.C., 1997. Stable transformation of *Chlorella*: rescue of nitrate reductase-deficient mutants with the nitrate reductase gene. *Curr. Microbiol.* 35 (6), 356–362. <https://doi.org/10.1007/s002849900268>.
- Day, R.N., Davidson, M.W., 2009. The fluorescent protein palette: tools for cellular imaging. *Chem. Soc. Rev.* 38 (10), 2887–2921. <https://doi.org/10.1039/b901966a>.
- de Jaeger, L., Carreres, B.M., Springer, J., Schaap, P.J., Eggink, G., Martins Dos Santos, V. A.P., Wijffels, R.H., Martens, D.E., 2018. *Neochloris oleoabundans* is worth its salt: transcriptomic analysis under salt and nitrogen stress. *PLoS One* 13 (4), 1–21. <https://doi.org/10.1371/journal.pone.0194834>.
- De Luca, M., Pappalardo, I., Limongi, A.R., Viviano, E., Radice, R.P., Todisco, S., Martelli, G., Infantino, V., Vassallo, A., 2021. Lipids from microalgae for cosmetic applications. *Cosmetics* 8 (2). <https://doi.org/10.3390/cosmetics8020052>.
- Deblöis, C.P., Dufresne, K., Juneau, P., 2013. Response to variable light intensity in photoacclimated algae and cyanobacteria exposed to atrazine. *Aquat. Toxicol.* 126, 77–84. <https://doi.org/10.1016/j.aquatox.2012.09.005>.
- DePriest, M.S., Bhattacharya, D., López-Bautista, J.M., 2013. The plastid genome of the red macroalga *Grateloupia taiwanensis* (Halymeniales). *PLoS One* 8 (7), 1–7. <https://doi.org/10.1371/journal.pone.0068246>.
- Deusch, O., Landan, G., Roettger, M., Gruenheit, N., Kowallik, K.V., Allen, J.F., Martin, W., Dagan, T., 2008. Genes of cyanobacterial origin in plant nuclear genomes point to a heterocyst-forming plastid ancestor. *Mol. Biol. Evol.* 25 (4), 748–761. <https://doi.org/10.1093/molbev/msn022>.
- Díaz-Santos, Encarnación, De la Vega, Marta, Vila, Marta, Vigara, Javier, León, R., 2013. Efficiency of different heterologous promoters in the unicellular. *Biotechnol. Prog.* 29 (2), 319–328.
- Ding, N., Wang, L., Kang, Y., Luo, K., Zeng, D., Bon, Y., Qiuyun, M., Lixuan, Z., Jiwen, Z., Feng, L., Zeng, Q.Z.A.L., Jiang, A.J.L.A.F., 2020. The comparison of transcriptomic response of green microalga *Chlorella sorokiniana* exposure to environmentally relevant concentration of cadmium (II) and 4-n-nonylphenol. *Environ. Geochem. Health* 42 (9), 2881–2894. <https://doi.org/10.1007/s10653-020-00526-1>.
- Doron, L., Segal, N., Shapira, M., 2016. Transgene expression in microalgae — from tools to applications technical approaches used for. *Front. Plant Sci.* 7 (April), 1–24. <https://doi.org/10.3389/fpls.2016.00505>.
- Duan, L., Chen, Q., Duan, S., 2019. Transcriptional analysis of *Chlorella pyrenoidosa* exposed to bisphenol A. *Int. J. Environ. Res. Public Health* 16, 1374. <https://doi.org/10.3390/ijerph16081374>.
- Dubois, G., Perez-michaut, C., Amato, A., Falciatore, A., Juillerat, A., Beurdeley, M., Voytas, D.F., Cavarec, L., Duchateau, P., 2014. *Phaeodactylum tricornutum* for biotechnology. *May*, pp. 1–7. <https://doi.org/10.1038/ncomms4831>.
- Dyo, Y.M., Purton, S., 2018. The algal chloroplast as a synthetic biology platform for production of therapeutic proteins. *Microbiology (United Kingdom)* 113–121. <https://doi.org/10.1099/mic.0.000599>.
- Eichler-Stahlberg, A., Weisheit, W., Ruecker, O., Heitzer, M., 2009. Strategies to facilitate transgene expression in *Chlamydomonas reinhardtii*. *Planta* 229 (4), 873–883. <https://doi.org/10.1007/s00425-008-0879-x>.
- Ekhtari, S., Razeghi, J., Hasanpur, K., Kianianmomeni, A., 2019. Different regulations of cell-type transcription by UV-B in multicellular green alga *Volvox carteri*. *Plant Signal. Behav.* 14 (11) <https://doi.org/10.1080/15592324.2019.1657339>.
- Esland, L., Larrea-Alvarez, M., Purton, S., 2018. Selectable markers and reporter genes for engineering the chloroplasts of *Chlamydomonas reinhardtii*. *Biology* 7 (4). <https://doi.org/10.3390/biology7040046>.
- Fajardo, C., De Donato, M., Carrasco, R., Martínez-Rodríguez, G., Mancera, J.M., Fernández-Aceró, F.J., 2020a. Advances and challenges in genetic engineering of microalgae. *Rev. Aquac.* 12 (1), 365–381. <https://doi.org/10.1111/raq.12322>.
- Fajardo, C., Carrasco-Reinado, R., Escobar-Niño, A., Fernández-Aceró, F.J., 2020b. Microalgae: Molecular biology, proteomics and applications. In: Prime Archives in Molecular Biology, July. <https://doi.org/10.37247/pamb.1.2020.45>.
- Fan, J., Xu, H., Li, Y., 2016. Transcriptome-based global analysis of gene expression in response to carbon dioxide deprivation in the green alga *Chlorella pyrenoidosa*. *ALGAL* 16, 12–19. <https://doi.org/10.1016/j.algal.2016.02.032>.
- Fang, L., Sun, D., Xu, Z., He, J., Qi, S., Chen, X., Chew, W., Liu, J., 2015. Transcriptomic analysis of a moderately growing subisolate *Botryococcus braunii* 779 (Chlorophyta) in response to nitrogen deprivation. *Biotechnol. Biofuels* 1–21. <https://doi.org/10.1186/s13068-015-0307-y>.
- Fayyaz, M., Chew, K.W., Ling, T.C., Ng, I., 2020. Genetic engineering of microalgae for enhanced biorefinery capabilities. *Biotechnol. Adv.* 43 (May), 107554 <https://doi.org/10.1016/j.biotechadv.2020.107554>.
- Feng, S., Xue, L., Liu, H., Lu, P., 2009. Improvement of efficiency of genetic transformation for Dunaliella salina by glass beads method. *Mol. Biol. Rep.* 36, 1433–1439. <https://doi.org/10.1007/s11033-008-9333-1>.
- FERENZI, A., Chew, Y.P., KROLL, E., von KOPPENFELS, C., HUDSON, A., MOLNAR, A., 2021. Mechanistic and genetic basis of single-strand templated repair at Cas12a-induced

- DNA breaks in *Chlamydomonas reinhardtii*. Nat. Commun. 12 (1), 1–12. <https://doi.org/10.1038/s41467-021-27004-1>.
- Fernández-Acero, F.J., Amil-Ruiz, F., Durán-Peña, M.J., Carrasco, R., Fajardo, C., Guarnizo, P., Fuentes-Almagro, C., Vallejo, R.A., 2019. Valorisation of the microalgae *Nannochloropsis gaditana* biomass by proteomic approach in the context of circular economy. J. Proteome 193 (August 2018), 239–242. <https://doi.org/10.1016/j.jprot.2018.10.015>.
- Ferrari, M., Muto, A., Bruno, L., Cozza, R., 2023. DNA methylation in algae and its impact on abiotic stress responses. Plants 12 (2). <https://doi.org/10.3390/plants12020241>.
- Fischer, N., Rochaix, J.D., 2001. The flanking regions of PsaD drive efficient gene expression in the nucleus of the green alga *Chlamydomonas reinhardtii*. Mol. Gen. Genomics. 265 (5), 888–894. <https://doi.org/10.1007/s004380100485>.
- Foflonker, F., Price, D.C., Qiu, H., Palenik, B., Wang, S., Bhattacharya, D., 2015. Genome of the halotolerant green alga *Picoclorum sp.* reveals strategies for thriving under fluctuating environmental conditions. Environ. Microbiol. 17 (2), 412–426. <https://doi.org/10.1111/1462-2920.12541>.
- Freudenberg, R.A., Wittemeier, L., Einhaus, A., Baier, T., Kruse, O., 2022. The spermidine synthase gene SPDI: A novel auxotrophic marker for *Chlamydomonas reinhardtii* designed by enhanced CRISPR/Cas9 gene editing. Cells 11 (5), 1–17. <https://doi.org/10.3390/cells11050837>.
- Fu, Y., Luo, G.Z., Chen, K., Deng, X., Yu, M., Han, D., Hao, Z., Liu, J., Lu, X., Doré, L.C., Weng, X., Ji, Q., Mets, L., He, C., 2015. N6-methyldeoxyadenosine marks active transcription start sites in *Chlamydomonas*. Cell 161 (4), 879–892. <https://doi.org/10.1016/j.cell.2015.04.010>.
- Fu, W., Nelson, D.R., Mystikou, A., Daakour, S., Salehi-Ashtiani, K., 2019. Advances in microalgal research and engineering development. Curr. Opin. Biotechnol. 59, 157–164. <https://doi.org/10.1016/j.cobio.2019.05.013>.
- Fujiiwara, T., Ohnuma, M., Kuroiwa, T., Ohbayashi, R., Hirooka, S., Miyagishima, S.Y., 2017. Development of a double nuclear gene-targeting method by two-step transformation based on a newly established chloramphenicol-selection system in the red alga *Cyanidioschyzon merolae*. Front. Plant Sci. 8 (March), 1–10. <https://doi.org/10.3389/fpls.2017.00343>.
- Gangl, D., Zedler, J.A., Rajakumar, P.D., Martinez, E.M.R., Riseley, A., Włodarczyk, A., Robinson, C., 2015. Biotechnological exploitation of microalgae. J. Exp. Bot. 66 (22), 6975–6990. <https://doi.org/10.1093/jxb/erv426>.
- Gao, C., Wang, Y., Shen, Y., Yan, D., He, X., Dai, J., Wu, Q., 2014. Oil accumulation mechanisms of the oleaginous microalga *Chlorella protothecoides* revealed through its genome, transcriptomes, and proteomes. BMC Genomics 1–14. <https://doi.org/10.1186/1471-2164-15-582>.
- Gao, G., Wu, M., Fu, Q., Li, X., Xu, J., 2019. A two-stage model with nitrogen and silicon limitation enhances lipid productivity and biodiesel features of the marine bloom-forming diatom *Skeletonema costatum*. Bioresour. Technol. 289, 121717 <https://doi.org/10.1016/j.biortech.2019.121717>.
- Gao, B., Huang, L., Lei, X., Meng, G., Liu, J., Zhang, C., 2020. Evaluation and transcriptome analysis of the novel oleaginous microalga *Lobosphaera bisecta* (trebouxiophyceae, chlorophyta) for arachidonic acid production. Mar. Drugs 18 (5). <https://doi.org/10.3390/md18050229>.
- Garduño-González, K.A., Peña-Benavides, S.A., Araújo, R.G., Castillo-Zacarías, C., Melchor-Martínez, E.M., Oyervides-Muñoz, M.A., Sosa-Hernández, J.E., Purton, S., Iqbal, H.M.N., Parra-Saldívar, R., 2022. Current challenges for modern vaccines and perspectives for novel treatment alternatives. J. Drug Deliv. Sci. Technol. 70 (March) <https://doi.org/10.1016/j.jddst.2022.103222>.
- Garoma, T., Shackelford, T., 2014. Bioresource technology electroporation of *Chlorella vulgaris* to enhance biomethane production. Bioresour. Technol. 169, 778–783. <https://doi.org/10.1016/j.biortech.2014.07.001>.
- Gauthier, M.R., Senhorin, G.N.A., Scott, J.A., 2020. Microalgae under environmental stress as a source of antioxidants. Algal Res. 52 (October), 102104 <https://doi.org/10.1016/j.algal.2020.102104>.
- Genevière, A.M., Derelle, E., Escande, M.L., Grimsley, N., Klopp, C., Ménager, C., Michel, A., Moreau, H., 2020. Responses to iron oxide and zinc oxide nanoparticles in echinoderm embryos and microalgae: uptake, growth, morphology, and transcriptomic analysis. Nanotoxicology 14 (10), 1342–1361. <https://doi.org/10.1080/17435390.2020.1827074>.
- Gierz, S.L., Forêt, S., Leggat, W., 2017. Transcriptomic analysis of thermally stressed symbiodinium reveals differential expression of stress and metabolism genes, 8 (February), 1–20. <https://doi.org/10.3389/fpls.2017.00271>.
- Gong, Y., Kang, N.K., Kim, Y.U., Wang, Z., Wei, L., Xin, Y., Shen, C., Wang, Q., You, W., Lim, J.M., Jeong, S.W., Park, Y. Il, Oh, H.M., Pan, K., Poliner, E., Yang, G., Li-Beisson, Y., Li, Y., Hu, Q., Xu, J., 2020. The NanDeSyn database for *Nannochloropsis* systems and synthetic biology. Plant J. 104 (6), 1736–1745. <https://doi.org/10.1111/tpj.15025>.
- Grama, S.B., Liu, Z., Li, J., 2022. Emerging trends in genetic engineering of microalgae for commercial applications. Mar. Drugs 20 (5), 1–24. <https://doi.org/10.3390/md20050285>.
- Greiner, A., Keltzborn, S., Evers, H., Kreimer, G., Sizova, I., Hegemann, P., 2017. Targeting of photoreceptor genes in *Chlamydomonas reinhardtii* via zinc-finger nucleases and CRISPR / Cas9, 29 (October), 2498–2518. <https://doi.org/10.1105/tpc.17.00659>.
- Guo, S.L., Zhao, X.Q., Tang, Y., Wan, C., Alam, M.A., Ho, S.H., Bai, F.W., Chang, J.S., 2013. Establishment of an efficient genetic transformation system in *Scenedesmus obliquus*. J. Biotechnol. 163 (1), 61–68. <https://doi.org/10.1016/j.jbiotec.2012.10.020>.
- Guo, L., Liang, S., Zhang, Z., Liu, H., Wang, S., Pan, K., Xu, J., Ren, X., Pei, S., Yang, G., 2019. Genome assembly of *Nannochloropsis oceanica* provides evidence of host nucleus overthrow by the symbiont nucleus during speciation. Commun. Biol. 2 (1), 1–12. <https://doi.org/10.1038/s42003-019-0500-9>.
- Hallmann, A., Sumper, M., 1996. The *Chlorella hexose* / H symporter is a useful selectable marker and biochemical reagent when expressed in *Volvox*, 93 (January), 669–673. <https://doi.org/10.1073/pnas.93.2.669>.
- Hallmann, A., Wodniok, S., 2006. Swapped green algal promoters: AphVIII-based gene constructs with *Chlamydomonas* flanking sequences work as dominant selectable markers in *Volvox* and vice versa. Plant Cell Rep. 25 (6), 582–591. <https://doi.org/10.1007/s00299-006-0121-x>.
- Hallmann, A., Rappel, A., I, L. B., 1999. Genetic engineering of the multicellular green alga *Volvox*: a modified and multiplied bacterial antibiotic resistance gene as a dominant selectable marker, 17 (October 1998), 99–109.
- Hanschen, E.R., Marriage, T.N., Ferris, P.J., Hamaji, T., Toyoda, A., Fujiyama, A., Neme, R., Noguchi, H., Minakuchi, Y., Suzuki, M., Kawai-Toyooka, H., Smith, D.R., Sparks, H., Anderson, J., Bakarić, R., Luria, V., Karger, A., Kirschner, M.W., Durand, P.M., Olson, B.J.S.C., 2016. The *Gonium pectorale* genome demonstrates co-option of cell cycle regulation during the evolution of multicellularity. Nat. Commun. 7, 1–10. <https://doi.org/10.1038/ncomms11370>.
- Hartley, J.L., Temple, G.F., Brasch, M.A., 2000. DNA cloning using in vitro site-specific recombination. Genome Res. 1788–1795 <https://doi.org/10.1101/gr.143000.that>.
- Hattman, S., Kenny, C., Berger, L., Pratt, K., 1978. Comparative study of DNA methylation in three unicellular eucaryotes. J. Bacteriol. 135 (3), 1156–1157. <https://doi.org/10.1128/jb.135.3.1156-1157.1978>.
- He, Q., Yang, H., Wu, L., Hu, C., 2015. Effect of light intensity on physiological changes, carbon allocation and neutral lipid accumulation in oleaginous microalgae. Bioresour. Technol. 191, 219–228. <https://doi.org/10.1016/j.biortech.2015.05.021>.
- Helliwell, K.E., Collins, S., Kazamia, E., Purton, S., Wheeler, G.L., Smith, A.G., 2015. Fundamental shift in vitamin B12 eco-physiology of a model alga demonstrated by experimental evolution. ISME J. 9 (6), 1446–1455. <https://doi.org/10.1038/ismej.2014.230>.
- Hempel, F., Lau, J., Klingl, A., Maier, U.G., 2011. Algae as protein factories : expression of a human antibody and the respective antigen in the diatom *Phaeodactylum tricornutum*. PLoS One 6 (12). <https://doi.org/10.1371/journal.pone.0028424>.
- Hirashima, T., Tajima, N., Sato, N., 2016. Draft genome sequences of four species of *Chlamydomonas* containing phosphatidylcholine. Genome Announc. 4 (5), 1–2. <https://doi.org/10.1128/genomeA.01070-16>.
- Hirooka, S., Hirose, Y., Kaneko, Y., Higuchi, S., Fujiwara, T., Onuma, R., Era, A., Ohbayashi, R., Uzuaka, A., Nozaki, H., Yoshikawa, H., Miyagishima, S.Y., 2017. Acidophilic green algal genome provides insights into adaptation to an acidic environment. Proc. Natl. Acad. Sci. USA 114 (39), E8304–E8313. <https://doi.org/10.1073/pnas.1707072114>.
- Hitchcock, A., Hunter, C.N., Sobotka, R., Leister, D., 2022. Redesigning the photosynthetic light reactions to enhance photosynthesis – the PhotoRedesign consortium. Plant J. 23–34 <https://doi.org/10.1111/tpj.15552>.
- Hoose, A., Vellacott, R., Storch, M., Freemont, P.S., Ryadnov, M.G., 2023. DNA synthesis technologies to close the gene writing gap. Nat. Rev. Chem. 7 (3), 144–161. <https://doi.org/10.1038/s41570-022-00456-9>.
- Hu, L., He, J., Dong, M., Tang, X., Jiang, P., Lei, A., Wang, J., 2020. Divergent metabolic and transcriptomic responses of *Synechocystis* sp. PCC 6803 to salt stress after adaptive laboratory evolution. Algal Res. 47 (February), 101856 <https://doi.org/10.1016/j.algal.2020.101856>.
- Hu, X., Luo, Y., Man, Y., Tang, X., Bi, Z., Ren, L., 2022. Lipidomic and transcriptomic analysis reveals the self-regulation mechanism of *Schizochytrium* sp. in response to temperature stresses. Algal Res. 64 (December 2021), 102664 <https://doi.org/10.1016/j.algal.2022.102664>.
- Huang, A., He, L., Wang, G., 2011. Identification and characterization of microRNAs from *Phaeodactylum tricornutum* by high- throughput sequencing and bioinformatics analysis. BMC Genomics 12, 337. <https://doi.org/10.1186/1471-2164-12-337>.
- Huang, B., Qu, G., He, Y., Zhang, J., Fan, J., Tang, T., 2022. Study on high-CO<sub>2</sub> tolerant *Dunaliella salina* and its mechanism via transcriptomic analysis. Front. Bioeng. Biotechnol. 10 (December), 1–12. <https://doi.org/10.3389/fbioe.2022.1086357>.
- Ishikawa, M., Nomura, T., Tamaki, S., Ozasa, K., Suzuki, T., Toyooka, K., Hirota, K., Yamada, K., Suzuki, K., Mochida, K., 2022. CRISPR/Cas9-mediated generation of non-motile mutants to improve the harvesting efficiency of mass-cultivated *Euglena gracilis*. Plant Biotechnol. J. 20 (11), 2042–2044. <https://doi.org/10.1111/pbi.13904>.
- Ishizuka, T., Kakuda, M., Araki, R., Yawo, H., 2006. Kinetic evaluation of photosensitivity in genetically engineered neurons expressing green algae light-gated channels. Neurosci. Res. 54, 85–94. <https://doi.org/10.1016/j.neures.2005.10.009>.
- Jabre, L.J., Allen, A.E., McCain, J.S.P., Mccrow, J.P., Tenenbaum, N., Spacken, J.L., 2021. Molecular underpinnings and biogeochemical consequences of enhanced diatom growth in a warming Southern Ocean. Proc. Natl. Acad. Sci. USA 118 (30), 1–9. <https://doi.org/10.1073/pnas.2107238118>.
- Jaeger, D., Hübner, W., Huser, T., Mussgnug, J.H., Kruse, O., 2017. Nuclear transformation and functional gene expression in the oleaginous microalga *Monoraphidium neglectum*. J. Biotechnol. 249, 10–15. <https://doi.org/10.1016/j.jbiotec.2017.03.011>.
- Jeon, S., Lim, J.M., Lee, H.G., Shin, S.E., Kang, N.K., Park, Y.II., 2017. Biotechnology for biofuels current status and perspectives of genome editing technology for microalgae. Biotechnol. Biofuels 1–18. <https://doi.org/10.1186/s13068-017-0957-z>.
- Jeon, M.S., Han, S. Il, Jeon, M., Choi, Y.E., 2021. Enhancement of phycoerythrin productivity in *Porphyridium purpureum* using the clustered regularly interspaced short palindromic repeats/CRISPR-associated protein 9 ribonucleoprotein system.

- Bioresour. Technol. 330 (February), 124974 <https://doi.org/10.1016/j.biotech.2021.124974>.
- Jiang, S.W., Hwangbo, K., Lim, J.M., Nam, S.W., Lee, B.S., Jeong, B.R., Chang, Y.K., Jeong, W.J., Park, Y.I., 2020. Genetic impairment of cellulose biosynthesis increases cell wall fragility and improves lipid extractability from oleaginous alga *Nannochloropsis salina*. Microorganisms 8 (8), 1–14. <https://doi.org/10.3390/microorganisms8081195>.
- Jeong, B., Ryoo, Jang, J., Jin, E.S., 2023. Genome engineering via gene editing technologies in microalgae. Bioresour. Technol. 373 (January), 128701 <https://doi.org/10.1016/j.biotech.2023.128701>.
- Jiang, W.Z., Weeks, D.P., 2017. A gene-within-a-gene Cas9 / sgRNA hybrid construct enables gene editing and gene replacement strategies in *Chlamydomonas reinhardtii*. Algal Res. 26 (April), 474–480. <https://doi.org/10.1016/j.algal.2017.04.001>.
- Johnson, D.C., Widlanski, T.S., 2003. Overview of the synthesis of nucleoside phosphates and polyphosphates. Curr. Protoc. Nucleic Acid Chem. 15 (1), 1–31. <https://doi.org/10.1002/047142700.nc1301s15>.
- Juneja, A., Chaplen, F.W.R., Murthy, G.S., 2016. Genome scale metabolic reconstruction of *Chlorella variabilis* for exploring its metabolic potential for biofuels. Bioresour. Technol. 213, 103–110. <https://doi.org/10.1016/j.biotech.2016.02.118>.
- Kang, S., Jeon, S., Kim, S., Chang, Y.K., Kim, Y.C., 2020. Development of a pVEC peptide – based ribonucleaseprotein (RNP) delivery system for genome editing using CRISPR / Cas9 in *Chlamydomonas reinhardtii*. Sci. Rep. 1–11 <https://doi.org/10.1038/s41598-020-78968-x>.
- Kania, K., Zienkiewicz, M., Drożak, A., 2020. Stable transformation of unicellular green alga *Coccomyxa subellipsoidea* C-169 via electroporation. Protoplasma 257 (2), 607–611. <https://doi.org/10.1007/s00709-019-01447-2>.
- Karas, B.J., Diner, R.E., Lefebvre, S.C., McQuaid, J., Phillips, A.P.R., Noddings, C.M., Brunson, J.K., Valas, R.E., Deerinck, T.J., Jablanovic, J., Gillard, J.T.F., Beeri, K., Ellisman, M.H., Glass, J.L., Hutchison, C.A., Smith, H.O., Venter, J.C., Allen, A.E., Dupont, C.L., Weyman, P.D., 2015. Designer diatom episomes delivered by bacterial conjugation. Nat. Commun. 6 <https://doi.org/10.1038/ncomms7925>.
- Kasai, Y., Tsukahara, T., Ikeda, F., Ide, Y., Harayama, S., 2018. Metabolic engineering using iterative self-cloning to improve lipid productivity in *Coccomyxa*. Sci. Rep. 8 (1), 1–11. <https://doi.org/10.1038/s41598-018-30254-7>.
- Kathiresan, S., Sarada, R., 2009. Towards genetic improvement of commercially important microalgae *Haematococcus pluvialis* for biotech applications. J. Phycol. 553–558 <https://doi.org/10.1007/s10811-009-9414-0>.
- Kathiresan, S., Chandrashekhar, A., Ravishankar, G.A., Sarada, R., 2009. Agrobacterium-mediated plant transformation: the biology behind the “Gene-Jockeying” tool. J. Phycol. 45 (3), 642–649. <https://doi.org/10.1111/j.1529-8817.2009.00688.x>.
- Kato, Y., Inabe, K., Hidese, R., Kondo, A., Hasutsuma, T., 2022. Bioresource technology metabolomics-based engineering for biofuel and bio-based chemical production in microalgae and cyanobacteria: a review. Bioresour. Technol. 344 (PA), 126196 <https://doi.org/10.1016/j.biotech.2021.126196>.
- Keeling, P.J., Burki, F., Wilcox, H.M., Allam, B., Allen, E.E., Amaral-Zettler, L.A., Armbrust, E.V., Archibald, J.M., Bharti, A.K., Bell, C.J., Beszteri, B., Bidle, K.D., Cameron, C.T., Campbell, L., Caron, D.A., Cattolico, R.A., Collier, J.L., Coyne, K., Davy, S.K., Worden, A.Z., 2014. The marine microbial eukaryote transcriptome sequencing project (MMETSP): illuminating the functional diversity of eukaryotic life in the oceans through transcriptome sequencing. PLoS Biol. 12 (6) <https://doi.org/10.1371/journal.pbio.1001889>.
- Kim, E.J., Cerutti, H., 2009. Targeted gene silencing by RNA interference in *Chlamydomonas*. In: Methods in Cell Biology, , First edit93. Elsevier. [https://doi.org/10.1016/S0091-679X\(08\)93005-3](https://doi.org/10.1016/S0091-679X(08)93005-3). Issue 08.
- Kim, E., Ma, X., Cerutti, H., 2015. Bioresource technology gene silencing in microalgae: mechanisms and biological roles. Bioresour. Technol. 184, 23–32. <https://doi.org/10.1016/j.biotech.2014.10.119>.
- Kim, J.H., Kim, N., Moon, H., Lee, S., Jeong, S.Y., Diaz-Pulido, G., Edwards, M.S., Kang, J.H., Kang, E.J., Oh, H.J., Hwang, J.D., Kim, I.N., 2020. Global warming offsets the ecophysiological stress of ocean acidification on temperate crustose coralline algae. Mar. Pollut. Bull. 157 (May), 111324 <https://doi.org/10.1016/j.marpolbul.2020.111324>.
- Kim, J., Chang, K.S., Lee, S., Jin, E., 2021. Establishment of a genome editing tool using CRISPR-Cas9 in *Chlorella vulgaris* UTEX395. Int. J. Mol. Sci. 22 (2), 1–11. <https://doi.org/10.3390/ijms22020480>.
- Koechler, S., Bertin, P.N., Plewniak, F., Baltenweck, R., Casiot, C., Heipieper, H.J., Bouchez, O., Arsène-Ploetz, F., Hugueney, P., Halter, D., 2016. Arsenite response in *Coccomyxa sp*. Carn explored by transcriptomic and non-targeted metabolomic approaches. Environ. Microbiol. 18 (4), 1289–1300. <https://doi.org/10.1111/1462-2920.13227>.
- Koh, H.G., Cho, J.M., Jeon, S., Chang, Y.K., Lee, B., Kang, N.K., 2023. Transcriptional insights into *Chlorella sp*. ABC-001: a comparative study of carbon fixation and lipid synthesis under different CO<sub>2</sub> conditions. Biotechnol. Biofuels Bioprod. 16 (1), 1–16. <https://doi.org/10.1186/s13068-023-02358-4>.
- Korkaric, M., Xiao, M., Behra, R., Eggen, R.I.L., 2015. Acclimation of *Chlamydomonas reinhardtii* to ultraviolet radiation and its impact on chemical toxicity. Aquat. Toxicol. 167, 209–219. <https://doi.org/10.1016/j.aquatox.2015.08.008>.
- Krishnan, A., Cano, M., Burch, T.A., Weissman, J.C., Posewitz, M.C., 2020. Genome editing using Cas9-RNA ribonucleaseprotein complexes in the high-productivity marine alga *Picochlorum celeri*. Algal Res. 49 (May), 101944 <https://doi.org/10.1016/j.algal.2020.101944>.
- Krumholz, E.W., Yang, H., Weisenhorn, P., Henry, C.S., Libourel, I.G., 2012. Genome-wide metabolic network reconstruction of the picoalga *Ostreococcus*. J. Exp. Bot. 63 (6), 2353–2362.
- Kujawinski, E.B., Longnecker, K., Alexander, H., Dyhrman, S.T., Fiore, C.L., Haley, S.T., Johnson, W.M., 2017. Phosphorus availability regulates intracellular nucleotides in marine eukaryotic phytoplankton. Limnol. Oceanogr. Lett. 2 (4), 119–129. <https://doi.org/10.1002/lol.20043>.
- Kumar, S.V., Misquitta, R.W., Reddy, V.S., Rao, B.J., Rajam, M.V., 2004. Genetic transformation of the green alga – *Chlamydomonas reinhardtii* by *Agrobacterium tumefaciens*. Plant Sci. 166 (3), 731–738. <https://doi.org/10.1016/j.plantsci.2003.11.012>.
- Kumar, G., Shekh, A., Jakhu, S., Sharma, Y., Kapoor, R., Sharma, T.R., 2020. Bioengineering of microalgae: recent advances, perspectives, and regulatory challenges for industrial application. Front. Bioeng. Biotechnol. 8, 914.
- Kurita, T., Moroi, K., Iwai, M., 2020. Genes to Cells Efficient and multiplexable genome editing using Platinum TALENs in oleaginous microalgae. *Nannochloropsis oceanica*. Genes Cells 695–702. <https://doi.org/10.1111/gtc.12805>. August.
- Kwon, S., Kang, N.K., Koh, H.G., Shin, S.E., Lee, B., Jeong, B.R., Chang, Y.K., 2018. Enhancement of biomass and lipid productivity by overexpression of a bZIP transcription factor in *Nannochloropsis salina*. Biotechnol. Bioeng. 115 (2), 331–340.
- Ladygin, V.G., 2003. The transformation of the unicellular alga *Chlamydomonas reinhardtii* by electroporation. Microbiology 72 (5), 585–591. <https://doi.org/10.1023/A:1026051402366>.
- Lapidot, M., Raveh, D., Sivan, A., Arad, S.M., Shapira, M., 2002. Stable chloroplast transformation of the unicellular red alga. *Porphyridium* Species 1 129 (May), 7–12. <https://doi.org/10.1104/pp.011023.Growth>.
- Légeret, B., Nguyen, H.M., Auroy, P., Beisson, F., Peltier, G., Blanc, G., 2016. Lipidomic and transcriptomic analyses of *Chlamydomonas reinhardtii* under heat stress unveil a direct route for the conversion of membrane lipids into storage lipids. Plant Cell Environ. 834–847. <https://doi.org/10.1111/pce.12656>.
- León-Bañares, R., González-Ballester, D., Galván, A., Fernández, E., 2004. Transgenic microalgae as green cell-factories. Trends Biotechnol. 22 (1), 45–52. <https://doi.org/10.1016/j.tibtech.2003.11.003>.
- Lerche, K., Hallmann, A., 2009. Stable nuclear transformation of *Gonium pectorale*. BMC Biotechnol. 9, 64. <https://doi.org/10.1186/1472-6750-9-64>.
- Levasseur, W., Perré, P., Pozzobon, V., 2020. A review of high value-added molecules production by microalgae in light of the classification. Biotechnol. Adv. 41 (January), 107545 <https://doi.org/10.1016/j.biotechadv.2020.107545>.
- Li, J., Xue, L., Yan, H., Wang, L., Liu, L., Lu, Y., Xie, H., 2007. The nitrate reductase gene-switch: A system for regulated expression in transformed cells of *Dunaliella salina*. Gene 403 (1–2), 132–142. <https://doi.org/10.1016/j.gene.2007.08.001>.
- Li, F., Gao, D., Hu, H., 2014. High-efficiency nuclear transformation of the oleaginous marine *Nannochloropsis* species using PCR product. Biosci. Biotechnol. Biochem. 78 (5), 812–817. <https://doi.org/10.1080/09168451.2014.905184>.
- Li, L., Zhang, G., Wang, Q., 2016a. De novo transcriptomic analysis of *Chlorella sorokiniana* reveals differential genes expression in photosynthetic carbon fixation and lipid production. BMC Microbiol. 16 (1), 1–12. <https://doi.org/10.1186/s12866-016-0839-8>.
- Li, T., Xu, G., Rong, J., Chen, H., He, C., Giordano, M., Wang, Q., 2016b. The acclimation of *Chlorella* to high-level nitrite for potential application in biological NO<sub>x</sub> removal from industrial flue gases. J. Plant Physiol. 195, 73–79. <https://doi.org/10.1016/j.jplph.2016.03.006>.
- Li, X., Yuan, Y., Cheng, D., Gao, J., Kong, L., Zhao, Q., Wei, W., Sun, Y., 2018. Exploring stress tolerance mechanism of evolved freshwater strain *Chlorella sp*. S30 under 30 g/L salt. Bioresour. Technol. 250 (November 2017), 495–504. <https://doi.org/10.1016/j.biotech.2017.11.072>.
- Li, D.W., Balamurugan, S., Yang, Y.F., Zheng, J.W., Huang, D., Zou, L.G., Yang, W.D., Liu, J.S., Guan, Y., Li, H.Y., 2019. Transcriptional regulation of microalgae for concurrent lipid overproduction and secretion. Sci. Adv. 5 (1), eaau3795.
- Li, H., Tan, J., Sun, T., Wang, Y., Meng, F., 2021. Acclimation of *Isochrysis galbana Parke* (Isochrysidaceae) for enhancing its tolerance and biodegradation to high-level phenol in seawater. Ecotoxicol. Environ. Saf. 207, 111571 <https://doi.org/10.1016/j.jecoenv.2020.111571>.
- Liang, M.H., Wang, L., Wang, Q., Zhu, J., Jiang, J.G., 2019. High-value bioproducts from microalgae: strategies and progress. Crit. Rev. Food Sci. Nutr. 59 (15), 2423–2441. <https://doi.org/10.1080/10408398.2018.1455030>.
- Lin, W.R., Ng, I.S., 2020. Development of CRISPR/Cas9 system in *Chlorella vulgaris* FSP-E to enhance lipid accumulation. Enzym. Microb. Technol. 133 (August 2019), 109458 <https://doi.org/10.1016/j.enzmictec.2019.109458>.
- Lin, W.R., Tan, S.I., Hsiang, C.C., Sung, P.K., Ng, I.S., 2019. Challenges and opportunity of recent genome editing and multi-omics in cyanobacteria and microalgae for biorefinery. Bioresour. Technol. 291 (June), 121932 <https://doi.org/10.1016/j.biotech.2019.121932>.
- Lin, J.Y., Lin, W.R., Ng, I.S., 2022. CRISPRa/i with adaptive single guide assisted regulation DNA (ASGARD) mediated control of *Chlorella sorokiniana* to enhance lipid and protein production. Biotechnol. J. 17 (10) <https://doi.org/10.1002/biot.202100514>.
- Ling, N., Li, W., Xu, G., Qi, Z., Ji, C., Liu, X., Cui, D., Sun, Y., 2021. Transcriptomic sequencing reveals the response of *Dunaliella salina* to copper stress via the increased photosynthesis and carbon mechanism. Mol. Omics 17 (5), 769–782. <https://doi.org/10.1039/dimo00125f>.
- Listmann, L., Hattich, G.S.I., Matthiessen, B., Reusch, T.B.H., 2020. Eco-evolutionary interaction in competing phytoplankton: nutrient driven genotype sorting likely explains dominance shift and species responses to CO<sub>2</sub>. Front. Mar. Sci. 7 (July) <https://doi.org/10.3389/fmars.2020.00634>.
- Little, S.M., Senhorinha, G.N.A., Saleh, M., Basiliko, N., Scott, J.A., 2021. Antibacterial compounds in green microalgae from extreme environments: a review. Algae 36 (1), 61–72. <https://doi.org/10.4490/algae.2021.36.6>.
- Liu, J., Sun, Z., Gerken, H., Huang, J., Jiang, Y., Chen, F., 2014a. Genetic engineering of the green alga *Chlorella zofingiensis*: a modified norflurazon-resistant phytene

- desaturase gene as a dominant selectable marker. *Appl. Microbiol. Biotechnol.* 98 (11), 5069–5079. <https://doi.org/10.1007/s00253-014-5593-y>.
- Liu, J., Sun, Z., Gerken, H., Liu, Z., Jiang, Y., Chen, F., 2014b. *Chlorella zofingiensis* as an alternative microalgal producer of astaxanthin: biology and industrial potential. *Mar. Drugs* 12 (6), 3487–3515. <https://doi.org/10.3390/MD12063487>.
- Liu, J., Sun, Z., Mao, X., Gerken, H., Wang, X., Yang, W., 2019. Multiomics analysis reveals a distinct mechanism of oleaginousness in the emerging model alga *Chromochloris zofingiensis*. *The Plant Journal* 98 (6), 1060–1077. <https://doi.org/10.1111/tpj.14302>.
- Llamas, A., Igéno, M.I., Galván, A., Fernández, E., 2002. Nitrate signalling on the nitrate reductase gene promoter depends directly on the activity of the nitrate transport systems in *Chlamydomonas*. *Plant J.* 30 (3), 261–271. <https://doi.org/10.1046/j.1365-313X.2002.01281.x>.
- Llavero-Pasquina, M., Geisler, K., Holzer, A., Mehrshahi, P., Mendoza-Ochoa, I., Newsad, S., Matthew Davey, M., Smith, A., 2022. Thiamine metabolism genes in diatoms are not regulated by thiamine despite the presence of predicted riboswitches. *New Phytol.* 235, 1853–1867. <https://doi.org/10.1111/nph.18296>.
- Lu, J.J., Ma, Y.L., Xing, G.L., Li, W.L., Kong, X.X., Li, J.Y., Wang, L.J., Yuan, H.L., Yang, J. S., 2019. Revelation of microalgae's lipid production and resistance mechanism to ultra-high cd stress by integrated transcriptome and physiochemical analyses. *Environ. Pollut.* 250, 186–195. <https://doi.org/10.1016/j.envpol.2019.04.018>.
- Lu, Y., Zhang, X., Gu, X., Lin, H., Melis, A., 2021. Critical reviews in biotechnology engineering microalgae: transition from empirical design to programmable cells. *Crit. Rev. Biotechnol.* 41 (8), 1233–1256. <https://doi.org/10.1080/07388551.2021.1917507>.
- Lumbreiras, V., Stevens, D.R., Purton, S., 1998. Efficient foreign gene expression in *Chlamydomonas reinhardtii* mediated by an endogenous intron. *Plant J.* 14 (4), 441–447. <https://doi.org/10.1046/j.1365-313X.1998.00145.x>.
- Luo, Q., Bian, C., Tao, M., Huang, Y., Zheng, Y., Lv, Y., Li, J., Wang, C., You, X., Jia, B., Xu, J., Li, J., Li, Z., Shi, Q., Hu, Z., 2019. Genome and transcriptome sequencing of the Astaxanthin-producing green microalga, *Haematococcus pluvialis*. *Genom. Biol. Evolut.* 11 (1), 166–173. <https://doi.org/10.1093/gbe/evy263>.
- Lv, J., Zhao, F., Feng, J., Liu, Q., Nan, F., Liu, X., Xie, S., 2019. Bioresource technology transcriptomic analysis reveals the mechanism on the response of *Chlorococcum* sp. GD to glucose concentration in mixotrophic cultivation. *Bioresour. Technol.* 288 (March), 121568 <https://doi.org/10.1016/j.biortech.2019.121568>.
- Martínez-Ruiz, M., Martínez-González, C.A., Kim, D.H., Santiesteban-Romero, B., Reyes-Pardo, H., Villaseñor-Zepeda, K.R., Meléndez-Sánchez, E.R., Ramírez-Gamboa, D., Díaz-Zamorano, A.L., Sosa-Hernández, J.E., Coronado-Apodaca, K.G., Gámez-Méndez, A.M., Iqbal, H.M.N., Parra-Saldivar, R., 2022. Microalgae bioactive compounds to topical applications products—A review. *Molecules* 27 (11), 1–22. <https://doi.org/10.3390/molecules27113512>.
- Matsuzaki, M., Misumi, O., Shin-I, T., Maruyama, S., Takahara, M., Miyagishima, S.Y., Mori, T., Nishida, K., Yagisawa, F., Nishida, K., Yoshida, Y., Nishimura, Y., Nakao, S., Kobayashi, T., Momoyama, Y., Higashiyama, T., Minoda, A., Sano, M., Nomoto, H., Kuroiwa, T., 2004. Genome sequence of the ultrasmall unicellular red alga *Cyanidioschyzon merolae* 10D. *Nature* 428 (6983). <https://doi.org/10.1038/nature02398>.
- Mayfield, D. J. B. and S. P., 2013. *Handbook of Microalgal Culture High-value Recombinant Protein Production in microalgae*.
- Meijer, H.J.G., van Himbergen, J.A.J., Musgrave, A., Munnik, T., 2017. Acclimation to salt modifies the activation of several osmotic stress-activated lipid signalling pathways in *Chlamydomonas*. *Phytochemistry* 135, 64–72. <https://doi.org/10.1016/j.phytochem.2016.12.014>.
- Merchant, S.S., Prochnik, S.E., Vallon, O., Harris, E.H., Karpowicz, J., Witman, G.B., Terry, A., Salamov, A., Fritz-laylin, L.K., Maréchal-drouard, L., Marshall, W.F., Qu, L., Nelson, D.R., Sanderfoot, A., Spalding, M.H., Kapitonov, V.V., Ren, Q., Cardol, P., Cerutti, H., Grossman, A.R., 2010. The chlamydomonas genome reveals the evolution of key animal and plant functions. *Science* (80-. ). 318, 245–251. <https://doi.org/10.1126/science.1143609>.
- Mini, P., Demurtas, O.C., Valentini, S., Pallara, P., Aprea, G., Ferrante, P., Giuliano, G., 2018. Agrobacterium-mediated and electroporation-mediated transformation of *Chlamydomonas reinhardtii*: A comparative study. *BMC Biotechnol.* 18 (1), 1–12. <https://doi.org/10.1186/s12896-018-0416-3>.
- Mishra, A., Medhi, K., Malaviya, P., Thakur, I.S., 2019. Omics approaches for microalgal applications: prospects and challenges. *Bioresour. Technol.* 291 (July), 121890 <https://doi.org/10.1016/j.biortech.2019.121890>.
- Mitra, M., Kumar, S., George, B., Shah, F., Mishra, S., 2015. A euryhaline *Nannochloropsis gaditana* with potential for nutraceutical (EPA) and biodiesel production. *ALGAL* 8, 161–167. <https://doi.org/10.1016/j.algal.2015.02.006>.
- Mobin, S., Alam, F., 2017. Some promising microalgal species for commercial applications: a review. *Energy Procedia* 110 (December 2016), 510–517. <https://doi.org/10.1016/j.egypro.2017.03.177>.
- Mócsai, R., Figl, R., Troschl, C., Strasser, R., Svehla, E., Windwarder, M., Thader, A., Altmann, F., 2019. N-glycans of the microalga *Chlorella vulgaris* are of the oligomannosidic type but highly methylated. *Sci. Rep.* 9 (1), 2–9. <https://doi.org/10.1038/s41598-018-36884-1>.
- Moellering, E.R., Benning, C., 2010. RNA interference silencing of a major lipid droplet protein affects lipid droplet size in *Chlamydomonas reinhardtii*. *Eukaryot. Cell* 9 (1), 97–106. <https://doi.org/10.1128/EC.00203-09>.
- Molnar, A., Bassett, A., Thuenemann, E., Schwach, F., Karkare, S., Ossowski, S., Weigel, D., Baulcombe, D., 2009. Highly specific gene silencing by artificial microRNAs in the unicellular alga *Chlamydomonas reinhardtii*, pp. 165–174. <https://doi.org/10.1111/j.1365-313X.2008.03767.x>.
- Moosburner, M.A., Gholami, P., McCarthy, J.K., Tan, M., Bielinski, V.A., Allen, A.E., 2020. Multiplexed knockouts in the model diatom *Phaeodactylum* by episomal delivery of a selectable Cas9. *Front. Microbiol.* 11 (January), 1–13. <https://doi.org/10.3389/fmicb.2020.00005>.
- Morais Junior, W.G., Gorgich, M., Corrêa, P.S., Martins, A.A., Mata, T.M., Caetano, N.S., 2020. Microalgae for biotechnological applications: cultivation, harvesting and biomass processing. *Aquaculture* 528 (May), 735562. <https://doi.org/10.1016/j.aquaculture.2020.735562>.
- Moreau, H., Verhelst, B., Couloux, A., Derelle, E., Rombauts, S., Grimsley, N., Van Bel, M., Poulain, J., Katinka, M., Hohmann-Marriott, M.F., Piganeau, G., Rouzé, P., Da Silva, C., Wincker, P., Van de Peer, Y., Vandepoele, K., 2012. Gene functionalities and genome structure in *Bathycoccus prasinus* reflect cellular specializations at the base of the green lineage. *Genome Biol.* 13 (8), R74. <https://doi.org/10.1186/gb-2012-13-8-r74>.
- Mulgund, A., 2022. Increasing lipid accumulation in microalgae through environmental manipulation, metabolic and genetic engineering : a review in the energy NEXUS framework. *Energy Nexus* 5 (February), 100054. <https://doi.org/10.1016/j.nexus.2022.100054>.
- Muñoz, C.F., de Jaeger, L., Sturme, M.H.J., Lip, K.Y.F., Olijslager, J.W.J., Springer, J., Wolbert, E.J.H., Martens, D.E., Eggink, G., Weusthuis, R.A., Wijffels, R.H., 2018. Improved DNA/protein delivery in microalgae – A simple and reliable method for the prediction of optimal electroporation settings. *Algal Res.* 33 (March), 448–455. <https://doi.org/10.1016/j.algal.2018.06.021>.
- Muto, M., Tanaka, M., Liang, Y., Yoshino, T., Matsumoto, M., Tanaka, T., 2015. Enhancement of glycerol metabolism in the oleaginous marine diatom *Fistulifera solaris* JPCC DA0580 to improve triacylglycerol productivity. *Bio technol. Biofuels* 8 (1), 1–7. <https://doi.org/10.1186/s13068-014-0184-9>.
- Naduthodi, M.I.S., Südfeld, C., Avitzianis, E.K., Trevisan, N., van Lith, E., Alcaide Sancho, J., D'Adamo, S., Barbosa, M., van der Oost, J., 2021. Comprehensive genome engineering toolbox for microalgae *Nannochloropsis oceanica* based on CRISPR-Cas systems. *ACS Synth. Biol.* 10 (12), 3369–3378. <https://doi.org/10.1021/acssynbio.1c00329>.
- Nagao, M., Matsui, K., Uemura, M., 2008. *Klebsormidium flaccidum*, a charophycean green alga, exhibits cold acclimation that is closely associated with compatible solute accumulation and ultrastructural changes. *Plant Cell Environ.* 31 (6), 872–885. <https://doi.org/10.1111/j.1365-3040.2008.01804.x>.
- Nawaly, H., Tsuji, Y., Matsuda, Y., 2020. Rapid and precise genome editing in a marine diatom, *Thalassiosira pseudonana* by Cas9 nickase (D10A). *Algal Res.* 47 (August 2019), 101855. <https://doi.org/10.1016/j.algal.2020.101855>.
- Nelson, D.R., Khraiwesh, B., Fu, W., Alseehk, S., Jaiswal, A., Chaiboonchoe, A., Hazzouri, K.M., O'Connor, M.J., Butterfoss, G.L., Drou, N., Rowe, J.D., Harb, J., Fernie, A.R., Gunsalus, K.C., Salehi-Ashtiani, K., 2017. The genome and phenome of the green alga *Chlorodidium* sp. UTEX 3007 reveal adaptive traits for desert acclimation. *Elife* 6, 1–24. <https://doi.org/10.7554/elife.25783>.
- Ng, I.S., Tan, S.I., Kao, P.H., Chang, Y.K., Chang, J.S., 2017. Recent developments on genetic engineering of microalgae for biofuels and bio-based chemicals. *Biotechnol. J.* 12 (10) <https://doi.org/10.1002/biot.20160044>.
- Norden-krichmar, T.M., Allen, A.E., Gaasterland, T., Hildebrand, M., 2011. Characterization of the small RNA transcriptome of the diatom, *Thalassiosira pseudonana*. *PLoS One* 6 (8). <https://doi.org/10.1371/journal.pone.0022870>.
- Nordin, N., Yusof, N., Maeda, T., Mustapha, N.A., Mohd Yusoff, M.Z., Raja Khairuddin, R. F., 2020. Mechanism of carbon partitioning towards starch and triacylglycerol in *Chlorella vulgaris* under nitrogen stress through whole-transcriptome analysis. *Biomass Bioenergy* 138 (September 2019), 105600. <https://doi.org/10.1016/j.biombioe.2020.105600>.
- Nozaki, H., Takano, H., Misumi, O., Terasawa, K., Matsuzaki, M., Maruyama, S., Nishida, K., Yagisawa, F., Yoshida, Y., Fujiwara, T., Takio, S., Tamura, K., Chung, S. J., Nakamura, S., Kuroiwa, H., Tanaka, K., Sato, N., Kuroiwa, T., 2007. A 100%-complete sequence reveals unusually simple genomic features in the hot-spring red alga *Cyanidioschyzon merolae*. *BMC Biol.* 5, 1–8. <https://doi.org/10.1186/1741-7007-5-28>.
- Oey, M., Ross, I.L., Hankamer, B., 2014. Gateway-assisted vector construction to facilitate expression of foreign proteins in the chloroplast of single celled algae. *PLoS One* 3–6. <https://doi.org/10.1371/journal.pone.0086841>.
- Ohnuma, M., Misumi, O., Fujiwara, T., Watanabe, S., Tanaka, K., Kuroiwa, T., 2009. Transient gene suppression in a red alga, *Cyanidioschyzon merolae* 10D. *Protoplasma* 236 (1–4), 107–112. <https://doi.org/10.1007/s00709-009-0056-5>.
- Okurowska, K., Karunakaran, E., Al-Fartootsy, A., Couto, N., Pandhal, J., 2021. Adapting the algal microbiome for growth on domestic landfill leachate. *Bioresour. Technol.* 319, 124246. <https://doi.org/10.1016/j.biortech.2020.124246>.
- Ortega-Escalante, J., Jasper, R., Miller, M.S., 2019. CRISPR/Cas9 mutagenesis in *Volvox carteri*. *Plant J.* 97, 661–672. <https://doi.org/10.1111/tpj.14149>.
- Ortiz-Matamoros, M.F., Islas-flores, T., Voigt, B., Menzel, D., 2015a. Heterologous DNA uptake in cultured *Symbiodinium* sp aided by *Agrobacterium tumefaciens*. *PLoS One* 1–16. <https://doi.org/10.1371/journal.pone.0132693>.
- Ortiz-Matamoros, M.F., Villanueva, M.A., Islas-Flores, T., 2015b. Transient transformation of cultured photosynthetic dinoflagellates (*Symbiodinium* spp.) with plant-targeted vectors. *Cienc. Mar.* 41 (1), 21–32. <https://doi.org/10.7773/cm.v41i1.2449>.
- Ortiz-Matamoros, M.F., Villanueva, M.A., Islas-Flores, T., 2018. Genetic transformation of cell-walled plant and algal cells: delivering DNA through the cell wall. *Brief. Funct. Genom.* 17 (1), 26–33. <https://doi.org/10.1093/bfgp/elx014>.
- Osorio, H., Jara, C., Fuenzalida, K., Rey-Jurado, E., Vásquez, M., 2019. High-efficiency nuclear transformation of the microalgae *Nannochloropsis oceanica* using Tn5 Transposome for the generation of altered lipid accumulation phenotypes. *Biotechnol. Biofuels* 12 (1), 1–12. <https://doi.org/10.1186/s13068-019-1475-y>.

- Page, T.M., Lawley, J.W., 2022. The next generation is here: a review of transcriptomic approaches in marine ecology. *Front. Mar. Sci.* 9 (May), 1–7. <https://doi.org/10.3389/fmars.2022.757921>.
- Palenik, B., Grimwood, J., Aerts, A., Rouze, P., Podell, S., Gaasterland, T., Napoli, C., Gendler, K., Manuell, A., Tai, V., Vallon, O., Rokhsar, D., Van De Peer, Y., 2007. The tiny eukaryote *Ostreococcus* provides genomic insights into the paradox of plankton speciation. *Proc. Natl. Acad. Sci. USA* 104, 7705–7710. <https://doi.org/10.1073/pnas.0611046104>.
- Palluk, S., Arlow, D.H., De Rond, T., Barthel, S., Kang, J.S., Bector, R., Baghdassarian, H.M., Truong, A.N., Kim, P.W., Singh, A.K., Hillson, N.J., Keasling, J.D., 2018. De novo DNA synthesis using polymerase nucleotide conjugates. *Nat. Biotechnol.* 36, 645–650. <https://doi.org/10.1038/nbt.4173>.
- Park, S., Kyndt, J., Chougule, K., Park, J., Brown, J.K., 2018. Low-phosphate-selected *Auxenochlorella protothecoides* redirects phosphate to essential pathways while producing more biomass. *PLoS One* 13, e0198953. <https://doi.org/10.1371/journal.pone.0198953>.
- Patel, V.K., Soni, N., Prasad, V., Sapre, A., Dasgupta, S., Bhadra, B., 2019. CRISPR–Cas9 System for Genome Engineering of Photosynthetic Microalgae. *Mol. Biotechnol.* 61 (8), 541–561. <https://doi.org/10.1007/s12033-019-00185-3>.
- Patelou, M., Infante, C., Dardelle, F., Randewig, D., Kouri, E.D., Uvdardi, M.K., Tsiplakou, E., Mantecón, L., Flemetakis, E., 2020. Transcriptomic and metabolomic adaptation of *Nannochloropsis gaditana* grown under different light regimes. *Algal Res.* 45 (December 2019), 101735. <https://doi.org/10.1016/j.algal.2019.101735>.
- Patwari, P., Pruckner, F., Fabris, M., 2023. Biosensors in microalgae: a roadmap for new opportunities in synthetic biology and biotechnology. *Biotechnol. Adv.* 68 (July), 108221. <https://doi.org/10.1016/j.biotechadv.2023.108221>.
- Peng, H., Wei, D., Chen, G., Chen, F., 2016. Biotechnology for biofuels transcriptome analysis reveals global regulation in response to CO<sub>2</sub> supplementation in oleaginous microalga *Cocomyxa subelliptoidea* C - 169. *Biotechnol. Biofuels* 1–17. <https://doi.org/10.1186/s13068-016-0571-5>.
- Perin, G., Bellan, A., Bernardi, A., Bezzo, F., Morosinotto, T., 2019. The potential of quantitative models to improve microalgae photosynthetic efficiency. *Physiol. Plant.* 166 (1), 380–391. <https://doi.org/10.1111/plp.12915>.
- Picariello, T., Hou, Y., Kubo, T., McNeill, N.A., Yanagisawa, H.A., Oda, T., Witman, G.B., 2020. TIM, a targeted insertional mutagenesis method utilizing CRISPR/Cas9 in *Chlamydomonas reinhardtii*. *PLoS One* 15 (5), 1–15. <https://doi.org/10.1371/journal.pone.0232594>.
- Poliner, E., Pulman, J.A., Zienkiewicz, K., Childs, K., Benning, C., Farré, E.M., 2018. A toolkit for *Nannochloropsis oceanica* CCMP1779 enables gene stacking and genetic engineering of the eicosapentaenoic acid pathway for enhanced long-chain polyunsaturated fatty acid production. *Plant Biotechnol. J.* 16 (1), 298–309. <https://doi.org/10.1111/pbi.12772>.
- Polle, J.E.W., Barry, K., Cushman, J., Schmutz, J., Tran, D., Hathwaik, L.T., Yim, W.C., Jenkins, J., McKie-Krisberg, Z., Prochnik, S., Lindquist, E., Dockter, R.B., Adam, C., Molina, H., Bunkenberg, J., Jin, E., 2017. Draft nuclear genome sequence of the halophilic and beta-carotene-accumulating green alga *Dunaliella Salina* strain CCAP19/18. *Genome Announc.* 5, 17–19. <https://doi.org/10.1128/genomeA.01105-17>.
- Polle, J.E.W., Calhoun, S., McKie-Krisberg, Z., Prochnik, S., Neofotis, P., Yim, W.C., Hathwaik, L.T., Jenkins, J., Molina, H., Bunkenberg, J., Grigoriev, I.V., Barry, K., Schmutz, J., Jin, E.S., Cushman, J.C., Magnusson, J.K., 2020. Genomic adaptations of the green alga *Dunaliella salina* to life under high salinity. *Algal Res.* 50 (March), 101990. <https://doi.org/10.1016/j.algal.2020.101990>.
- Pombert, J.F., Blouin, N.A., Lane, C., Boucias, D., Keeling, P.J., 2014. A lack of parasitic reduction in the obligate parasitic green alga *Helicosporidium*. *PLoS Genet.* 10 (5) <https://doi.org/10.1371/journal.pgen.1004355>.
- Poong, S.W., Lee, K.K., Lim, P.E., Pai, T.W., Wong, C.Y., Phang, S.M., Chen, C.M., Yang, C.H., Liu, C.C., 2018. RNA-seq-mediated transcriptomic analysis of heat stress response in a polar *Chlorella* sp. (Trebouxiophyceae, Chlorophyta). *J. Appl. Phycol.* 30 (6), 3103–3119. <https://doi.org/10.1007/s10811-018-1455-9>.
- Potvin, G., Zhang, Z., 2010. Strategies for high-level recombinant protein expression in transgenic microalgae: a review. *Biotechnol. Adv.* 28 (6), 910–918. <https://doi.org/10.1016/j.biotechadv.2010.08.006>.
- Prochnik, S.E., Umen, J., Nedelcu, A.M., Hallmann, A., Miller, S.M., Nishii, I., Ferris, P., Kuo, A., Mitros, T., Fritz-Laylin, L.K., Hellsten, U., Chapman, J., Simakov, O., Rensing, S.A., Terry, A., Pangilinan, J., Kapitonov, V., Jurka, J., Salamov, A., Rokhsar, D.S., 2010. Genomic analysis of organismal complexity in the multicellular green alga *Volvox carteri*. *Science* 329 (5988), 223–226. <https://doi.org/10.1126/science.1188800>.
- Puente-Sánchez, F., Díaz, S., Penacho, V., Aguilera, A., Olsson, S., 2018. Basis of genetic adaptation to heavy metal stress in the acidophilic green alga *Chlamydomonas acidophila*. *Aquat. Toxicol.* 200 (February), 62–72. <https://doi.org/10.1016/j.aquatox.2018.04.020>.
- Qiao, Y., Yang, F., Xie, T., Du, Z., Zhong, D., Qi, Y., Li, Y., Li, W., Lu, Z., Rao, J., Sun, Y., Zhou, M., 2020. Engineered algae: A novel oxygen-generating system for effective treatment of hypoxic cancer. *Sci. Adv.* 6 <https://doi.org/10.1126/sciadv.aba5996>.
- Qin, S., Lin, H., Jiang, P., 2012. Advances in genetic engineering of marine algae. *Biotechnol. Adv.* 30 (6) <https://doi.org/10.1016/j.biotechadv.2012.05.004>.
- Qiu, H., Price, D.C., Weber, A.P.M., Reeb, V., Chan Yang, E., Lee, J.M., Kim, S.Y., Yoon, H.S., Bhattacharya, D., 2013. Adaptation through horizontal gene transfer in the cryptoendolithic red alga *Galdieria phlegrea*. *Curr. Biol.* 23 (19), R865–R866. <https://doi.org/10.1016/j.cub.2013.08.046>.
- Quinn, J.M., Kropat, J., Merchant, S., 2003. Copper response element and Crr1-dependent Ni 2+ responsive promoter for induced, reversible gene expression in *Chlamydomonas reinhardtii*. *Eukaryot. Cell* 2 (5), 995–1002. <https://doi.org/10.1128/EC.2.5.995-1002.2003>.
- Rademacher, N., Kern, R., Fujiwara, T., Mettler-altmann, T., Miyashima, S., 2016. Photosynthetic glycolate oxidase is essential for the survival of the red alga *Cyanidioschyzon merolae* under ambient CO<sub>2</sub> conditions. *J. Exp. Bot.* 67 (10), 3165–3175. <https://doi.org/10.1093/jxb/erw118>.
- Rana, M.S., Prajapati, S.K., 2021. Resolving the dilemma of iron bioavailability to microalgae for commercial sustenance. *Algal Res.* 59 (August), 102458. <https://doi.org/10.1016/j.algal.2021.102458>.
- Ranjbar, S., Malcata, F.X., 2022. Is genetic engineering a route to enhance microalgae-mediated bioremediation of heavy metal-containing effluents? *Molecules* 27 (5). <https://doi.org/10.3390/molecules27051473>.
- Rasala, B.A., Mayfield, S.P., 2011. Therapeutics for the production of human protein therapeutics. *Bioeng. Bugs* 1018. <https://doi.org/10.4161/bbug.2.1.13423>.
- Rasala, B.A., Mayfield, S.P., 2015. Photosynthetic biomanufacturing in green algae: production of recombinant proteins for industrial, nutritional, and medical uses. *Photosynth. Res.* 123 (3), 227–239. <https://doi.org/10.1007/s11120-014-9994-7>.
- Rasala, B.A., Barrera, D.J., Ng, J., Plucinak, T.M., Rosenberg, J.N., Weeks, D.P., Oyler, A., Peterson, T.C., Haerizadeh, F., Mayfield, S.P., 2013. Expanding the spectral palette of fluorescent proteins for the green microalga *Chlamydomonas reinhardtii*. *Plant J.* 545–556. <https://doi.org/10.1111/tpj.12165>.
- Rastogi, R.P., Madamwar, D., Nakamoto, H., Incharoensakdi, A., 2020. Resilience and self-regulation processes of microalgae under UV radiation stress. *J. Photochem. Photobiol. C Photchem. Rev.* 43, 100322. <https://doi.org/10.1016/j.jphotchemrev.2019.100322>.
- Rathod, J.P., Prakash, G., Pandit, R., Lali, A.M., 2013. *Agrobacterium*-mediated transformation of promising oil-bearing marine algae *Parachlorella kessleri*. *Photosynth. Res.* 141–146. <https://doi.org/10.1007/s11120-013-9930-2>.
- Rathod, J.P., Gade, R.M., Rathod, D.R., Dudhare, M., 2017. A review on molecular tools of microalgal genetic transformation and their application for overexpression of different genes. *Int. J. Curr. Microbiol. App. Sci.* 6 (12), 3191–3207.
- Ren, Y., Sun, H., Deng, J., Huang, J., Chen, F., 2021. Carotenoid production from microalgae: Biosynthesis, salinity responses and novel biotechnologies. *Marine Drugs* 19 (12). <https://doi.org/10.3390/md19120713>.
- Ribeiro, Á.M., Foote, A.D., Kupczok, A., Frazão, B., Limborg, M.T., Piñeiro, R., Abalde, S., Rocha, S., da Fonseca, R.R., 2017. Marine genomics: news and views. *Mar. Genomics* 31, 1–8. <https://doi.org/10.1016/j.margen.2016.09.002>.
- Rismani-Yazdi, H., Haznedaroglu, B.Z., Hsin, C., Peccia, J., 2012. Transcriptomic analysis of the oleaginous microalga *Neochloris oleoabundans* reveals metabolic insights into triacylglyceride accumulation. *Biotechnol. Biofuels* 5. <https://doi.org/10.1186/1754-6834-5-74>.
- Rosas-Mendoza, S., 2016. Algae-based pharmaceuticals. In: *In Algae-Based Biopharmaceuticals*. <https://doi.org/10.1007/978-3-319-32232-2>.
- Rossoni, A.W., Scho, G., Lee, H.J., Rupp, R.L., Flachbart, S., Mettler-altmann, T., Weber, A.P.M., Eisenhut, M., 2019. Cold acclimation of the Thermoacidophilic red alga *Galdieria sulphuraria*: changes in gene expression and involvement of horizontally acquired genes. *Plant Cell Physiol.* 60 (December 2018), 702–712. <https://doi.org/10.1093/pcp/pcy240>.
- Rumin, J., Nicolau, E., de Oliveira, R.G., Fuentes-Grinewald, C., Picot, L., 2020. Analysis of scientific research driving microalgae market opportunities in Europe. *Marine Drugs* 18 (5). <https://doi.org/10.3390/md18050264>.
- Run, C., Fang, L., Fan, J., Fan, C., Luo, Y., Hu, Z., Li, Y., 2016. Stable nuclear transformation of the industrial alga *Chlorella pyrenoidosa*. *Algal Res.* 17, 196–201. <https://doi.org/10.1016/j.algal.2016.05.002>.
- Ryu, A.J., Ryool, B., Kang, N.K., Jeon, S., Sohn, M.G., Yun, H.J., Lim, J.M., Jeong, S.W., Park, Y. Il, Jeong, W.J., Park, S., Chang, Y.K., Jeong, K.J., 2021. Safe-harboring based novel genetic toolkit for *Nannochloropsis salina* CCMP1776: efficient overexpression of transgene via CRISPR/Cas9-mediated Knock-in at the transcriptional hotspot. *Bioresour. Technol.* 340 (July), 125676. <https://doi.org/10.1016/j.biortech.2021.125676>.
- Salguero, D.A.M., Fernández-Niño, M., Serrano-Bermúdez, L.M., Melo, D.O.P., Winck, F. V., Caldana, C., Barrios, A.F.G., 2018. Development of a *Chlamydomonas reinhardtii* metabolic network dynamic model to describe distinct phenotypes occurring at different CO<sub>2</sub> levels. *PeerJ* 6, e5528. <https://doi.org/10.7717/peerj.5528>.
- Sanchez, F., Geffroy, S., Norest, M., Yau, S., Moreau, H., Grimsley, N., 2019. Simplified transformation of *Ostreococcus tauri* using polyethylene glycol. *Genes* 10 (5), 1–16. <https://doi.org/10.3390/genes10050399>.
- Schaum, C.E., Barton, S., Bestion, E., Buckling, A., Garcia-Carreras, B., Lopez, P., Lowe, G., Pawar, S., Smirnoff, N., Trimmer, M., Yvon-Durocher, G., 2017. Adaptation of phytoplankton to a decade of experimental warming linked to increased photosynthesis. *Nat. Ecol. Evolut.* 1 (4), 1–7. <https://doi.org/10.1038/s41559-017-0094>.
- Scranton, M.A., Ostrand, J.T., Georgianna, D.R., Lofgren, S.M., Li, D., Ellis, R.C., Carruthers, D.N., Dräger, A., Masica, D.L., May, S.P., 2016. Synthetic promoters capable of driving robust nuclear gene expression in the green alga *Chlamydomonas reinhardtii*. *Algal Res.* 15, 135–142. <https://doi.org/10.1016/j.algal.2016.02.011>.
- Serif, M., Dubois, G., Finoux, A.L., Teste, M.A., Jallet, D., Daboussi, F., 2018. One-step generation of multiple gene knock-outs in the diatom *Phaeodactylum tricornutum* by DNA-free genome editing. *Nat. Commun.* 9 (1), 1–10. <https://doi.org/10.1038/s41467-018-06378-9>.
- Sharaf, A., De Michele, R., Sharma, A., Fakhari, S., Obornik, M., 2019. Transcriptomic analysis reveals the roles of detoxification systems in response to mercury in *Chromera velia*. *Biomolecules* 9 (11), 1–25. <https://doi.org/10.3390/biom9110647>.
- Sharma, A.K., Nymark, M., Sparstad, T., Bones, A.M., Winge, P., 2018. Transgene-free genome editing in marine algae by bacterial conjugation – comparison with biolistic CRISPR/Cas9 transformation. *Sci. Rep.* 8 (1), 1–11. <https://doi.org/10.1038/s41598-018-32342-0>.

- Sharma, A.K., Nymark, M., Flo, S., Sparstad, T., Bones, A.M., Winge, P., 2021. Simultaneous knockout of multiple LHCF genes using single sgRNAs and engineering of a high-fidelity Cas9 for precise genome editing in marine algae. *Plant Biotechnol. J.* 19 (8), 1658–1669. <https://doi.org/10.1111/pbi.13582>.
- Shetty, P., Gitau, M.M., Maróti, G., 2019. Salinity stress responses and adaptation mechanisms in eukaryotic green microalgae. *Cells* 8 (12), 1–16. <https://doi.org/10.3390/cells8121657>.
- Shi, C., Liu, X., Han, K., Peng, L., Li, L., Ge, Q., Fan, G., 2021. A database and comprehensive analysis of the algae genomes. *BioRxiv*. <https://doi.org/10.1101/2021.10.30.466624>.
- Siddiqui, A., Wei, Z., Boehm, M., Ahmad, N., 2020. Engineering microalgae through chloroplast transformation to produce high-value industrial products. *Biotechnol. Appl. Biochem.* 67 (1), 30–40. <https://doi.org/10.1002/bab.1823>.
- Simon, D.F., Descombes, P., Zerges, W., Wilkinson, K.J., 2008. Global expression profiling of *Chlamydomonas reinhardtii* exposed to trace levels of free cadmium. *Environ. Toxicol. Chem.* 27 (8), 1668–1675. <https://doi.org/10.1897/07-649.1>.
- Singh, P., Kumari, S., Guldhe, A., Misra, R., Rawat, I., Bux, F., 2016. Trends and novel strategies for enhancing lipid accumulation and quality in microalgae. *Renew. Sust. Energ. Rev.* 55, 1–16. <https://doi.org/10.1016/j.rser.2015.11.001>.
- Singh, A., Sharma, R., Pant, D., Malaviya, P., 2021. Science of the Total environment engineered algal biochar for contaminant remediation and electrochemical applications. *Sci. Total Environ.* 774, 145676 <https://doi.org/10.1016/j.scitotenv.2021.145676>.
- Sirohi, R., Joun, J., Choi, H.I., Gaur, V.K., Sim, S.J., 2021. Algal glycobiotechnology: omics approaches for strain improvement. *Microb. Cell Factories* 20 (1). <https://doi.org/10.1186/s12934-021-01656-6>.
- Sizova, I., Kelterborn, S., Verbenko, V., Kateriya, S., 2021. Chlamydomonas POLQ is necessary for CRISPR / Cas9-mediated gene targeting. *G3 Genes Genom. Genet.* 11 <https://doi.org/10.1093/g3journal/jkab114>.
- Song, I., Kim, J., Baek, K., Choi, Y., Shin, B.C., Jin, E.S., 2020. The generation of metabolic changes for the production of high-purity zeaxanthin mediated by CRISPR-Cas9 in *Chlamydomonas reinhardtii*. *Microb. Cell Factories* 19 (1), 1–9. <https://doi.org/10.1186/s12934-020-01480-4>.
- Song, I., Kim, S., Kim, J., Oh, H., Jang, J., Jeong, S.J., Baek, K., Shin, W.S., Sim, S.J., Jin, E.S., 2022. Macular pigment-enriched oil production from genome-edited microalgae. *Microb. Cell Factories* 21 (1), 1–12. <https://doi.org/10.1186/s12934-021-01736-7>.
- Sorrigué, D., Légeret, B., Cuiné, S., Morales, P., Mirabella, B., Guédeney, G., Li-Beisson, Y., Jetter, R., Peltier, G., Beisson, F., 2016. Microalgae synthesize hydrocarbons from long-chain fatty acids via a light-dependent pathway. *Plant Physiol.* 171 (4) <https://doi.org/10.1104/pp.16.00462>.
- Specht, E.A., Mayfield, S.P., 2013. Synthetic oligonucleotide libraries reveal novel regulatory elements in *Chlamydomonas* chloroplast mRNAs. *ACS Synth. Biol.* 2 (1), 34–46. <https://doi.org/10.1021/sb300069k>.
- Specht, E.A., Stephen, P., 2014. Algae-based oral recombinant vaccines. *Front. Microbiol.* 5 (February), 1–7. <https://doi.org/10.3389/fmicb.2014.00060>.
- Specht, E., Miyake-Stoner, S., Mayfield, S., 2010. Micro-algae come of age as a platform for recombinant protein production. *Biotechnol. Lett.* 32 (10), 1373–1383. <https://doi.org/10.1007/s10529-010-0326-5>.
- Sproles, A.E., Fields, F.J., Smalley, T.N., Le, C.H., Badary, A., Mayfield, S.P., 2021. Recent advancements in the genetic engineering of microalgae. *Algal Res.* 53 (December 2020), 102158 <https://doi.org/10.1016/j.algal.2020.102158>.
- Srinivasan, R., Gothandam, K.M., 2016. Synergistic action of D-glucose and acetosyringone on *Agrobacterium* strains for efficient *Dunaliella* transformation. *PLoS One* 11 (6), 1–13. <https://doi.org/10.1371/journal.pone.0158322>.
- Steadman Tyler, C.R., Hovde, B.T., Daligault, H.E., Zhang, X.L., Kunde, Y., Marrone, B.L., Twary, S.N., Starkenburg, S.R., 2019. High-quality draft genome sequence of the green alga *Tetraselmis striata* (Chlorophyta) generated from PacBio sequencing. *Microbiol. Resour. Announc.* 8 (43), 10–1128. <https://doi.org/10.1128/MRA.00780-19>.
- Steichen, S.A., Berim, A., Gang, D.R., Brown, J.K., 2022. Plant stress *Auxenochlorella protothecoides* populations adapted to low phosphate conditions accumulated more non-phosphorus glycerolipids and biomass than wild type progenitors. *Plant Stress* 6 (August), 100115. <https://doi.org/10.1016/j.stress.2022.100115>.
- Steinbrenner, J., Sandmann, G., 2006. Transformation of the green alga *Haematococcus pluvialis* with phytoene desaturase for accelerated astaxanthin biosynthesis. *Appl. Environ. Microbiol.* 72 (12), 7477–7484. <https://doi.org/10.1128/AEM.01461-06>.
- Stevens, D.R., Rochaix, J.D., Purton, S., 1996. The bacterial phleomycin resistance gene ble as a dominant selectable marker in *Chlamydomonas*. *Mol. Gen. Genet.* 251 (1), 23–30. <https://doi.org/10.1007/s004380050135>.
- Strenkert, D., Schmollinger, S., Schroda, M., 2013. Heat shock factor 1 counteracts epigenetic silencing of nuclear transgenes in *Chlamydomonas reinhardtii*. *Nucleic Acids Res.* 41 (10), 5273–5289. <https://doi.org/10.1093/nar/gkt224>.
- Sturme, M.H.J., Gong, Y., Miguel, J., Klok, A.J., Eggink, G., Wang, D., Xu, J., Wij, R.H., 2018. Transcriptome analysis reveals the genetic foundation for the dynamics of starch and lipid production in *Ettlia oleobundans*. *Algal Res.* 33 (December 2017), 142–155. <https://doi.org/10.1016/j.algal.2018.05.004>.
- Südfeld, C., Hub, M., Figueiredo, D., Naduthodi, M.I.S., Van Der Oost, J., Wijffels, H., Barbosa, M.J., Adamo, S.D., 2021. High-throughput insertional mutagenesis reveals novel targets for enhancing lipid accumulation in *Nannochloropsis oceanica*. *Metab. Eng.* 66 (March), 239–258. <https://doi.org/10.1016/j.ymben.2021.04.012>.
- Sumiya, N., Kawase, Y., Hayakawa, J., Matsuda, M., Nakamura, M., Era, A., Tanaka, K., Kondo, A., Hasunuma, T., Imamura, S., Miyagishima, S.Y., 2015. Expression of cyanobacterial acyl-ACP reductase elevates the triacylglycerol level in the red alga *Cyanidioschyzon merolae*. *Plant Cell Physiol.* 56 (10), 1962–1980. <https://doi.org/10.1093/pcp/pcv120>.
- Sun, X., Wang, W., Shen, J., Xu, N., 2014. Transcriptome sequencing of the marine microalga, *Chlorella pyrenoidosa* (Chlorophyta), and analysis of carbonic anhydrase expression under salt stress. *Bot. Mar.* 57 (5), 403–412. <https://doi.org/10.1515/bot-2014-0013>.
- Sun, X.M., Ren, L.J., Zhao, Q.Y., Ji, X.J., Huang, H., 2018. Microalgae for the production of lipid and carotenoids: A review with focus on stress regulation and adaptation. *Biotechnol. Biofuels* 11 (1), 1–16. <https://doi.org/10.1186/s13068-018-1275-9>.
- Suparmanian, U., Lam, M.K., Lim, J.W., Yusup, S., Tan, I.S., Lau, S.Y., Kodgire, P., Kachhwaha, S.S., 2022. Influence of environmental stress on microalgae growth and lipid profile: a systematic review. *Phytochem. Rev.* 7 <https://doi.org/10.1007/s11101-022-09810-7>.
- Tardu, M., Dikbas, U.M., Baris, I., Kavaklı, I.H., 2016. RNA-seq analysis of the transcriptional response to blue and red light in the extremophilic red alga, *Cyanidioschyzon merolae*. *Funct. Integrat. Genom.* 657–669 <https://doi.org/10.1007/s10142-016-0521-0>.
- Taunt, H.N., Stoffels, L., Purton, S., 2018. Green biologics: the algal chloroplast as a platform for making biopharmaceuticals. *Bioengineered* 5979. <https://doi.org/10.1080/21655979.2017.1377867>.
- Ten Lohuis, M.R., Miller, D.J., 1998. Genetic transformation of dinoflagellates (*Amphidinium* and *Symbiodinium*): expression of GUS in microalgae using heterologous promoter constructs. *Plant J.* 13 (3), 427–435. <https://doi.org/10.1046/j.1365-313X.1998.00040.x>.
- Teng, S.Y., Loy, A.C.M., Leong, W.D., How, B.S., Chin, B.L.F., Máša, V., 2019. Catalytic thermal degradation of *Chlorella vulgaris*: evolving deep neural networks for optimization. *Bioresour. Technol.* 292 (June), 121971 <https://doi.org/10.1016/j.biortech.2019.121971>.
- Thangaraj, B., Jolley, C.C., Sarrou, I., Bultema, J.B., Greyslak, J., Whitelegge, J.P., Lin, S., Kouril, R., Subramanyam, R., Boekema, E.J., Fromme, P., 2011. Efficient light harvesting in a dark, hot, acidic environment: the structure and function of PSI-LHCI from *Galdiera sulphuraria*. *Biophys. J.* 100 (1), 135–143.
- Tian, Q., Wang, J., Cui, L., Zeng, W., Qiu, G., Hu, Q., Peng, A., Zhang, D., Shen, L., 2022. Longitudinal physiological and transcriptomic analyses reveal the short term and long term response of *Synechocystis* sp. PCC6803 to cadmium stress. *Chemosphere* 303, 134727.
- Tirichine, L., Bowler, C., 2011. Decoding algal genomes: tracing back the history of photosynthetic life on earth. *Plant J.* 66 (1), 45–57. <https://doi.org/10.1111/j.1365-313X.2011.04540.x>.
- Torres-Tijj, Y., Fields, F.J., Mayfield, S.P., 2020. Microalgae as a future food source. *Biotechnol. Adv.* 41 (August 2019) <https://doi.org/10.1016/j.biotechadv.2020.107536>.
- Toulibah, H.E.S., El-Sheekh, M.M., Ismail, M.M., El-Kassas, H., 2022. A review of microalgae-and Cyanobacteria-based biodegradation of organic pollutants. *Molecules* 27 (3). <https://doi.org/10.3390/molecules27031141>.
- Van Baren, M.J., Bachy, C., Reistetter, E.N., Purvine, S.O., Grimwood, J., Sudek, S., Yu, H., Poirier, C., Deerinck, T.J., Kuo, A., Grigoriev, I.V., Wong, C., Smith, R.D., Callister, S.J., Wei, C., Schmutz, J., Worden, A.Z., 2016. Evidence-based green algal genomics reveals marine diversity and ancestral characteristics of land plants. *BMC Genomics* 1–22. <https://doi.org/10.1186/s12864-016-2585-6>.
- van Ooijen, G., Knox, K., Kis, K., Bouget, F.Y., Millar, A.J., 2012. Genomic transformation of the picoeukaryote *Ostreococcus tauri*. *J. Vis. Exp.* 65, 3–7. <https://doi.org/10.3791/4074>.
- Vanhoudt, N., Vandenhove, H., Leys, N., Janssen, P., 2018. Chemosphere Potential of higher plants , algae, and cyanobacteria for remediation of radioactively contaminated waters. *Chemosphere* 207, 239–254. <https://doi.org/10.1016/j.chemosphere.2018.05.034>.
- Vanier, G., Pierre-louis, L., Loutelier-bou, C., Vanier, J., Tchi-song, P.C., Remy-jouet, I., Mathieu-rivet, E., Bardor, M., 2017. Heterologous expression of the N -acetylglucosaminyltransferase I dictates a reinvestigation of the N -glycosylation pathway in *Chlamydomonas reinhardtii*. *Sci. Rep.* 1–12. <https://doi.org/10.1038/s41598-017-10698-z>.
- Velmurugan, N., Deka, D., 2018. Transformation techniques for metabolic engineering of diatoms and haptophytes : current state and prospects. *Appl. Microbiol. Biotechnol.* 102, 4255–4267. <https://doi.org/10.1007/s00253-018-8925-5>.
- Vieler, A., Wu, G., Tsai, C., Bullard, B., Cornish, A.J., Harvey, C., Reca, I., Thornburg, C., Achawanantakun, R., Buehl, C.J., Campbell, M.S., Cavalier, D., Childs, K.L., Clark, T.J., Deshpande, R., Erickson, E., Handee, W., Kong, Q., Li, X., Benning, C., 2012. Genome, functional gene annotation , and nuclear transformation of the heterokont oleaginous Alga *Nannochloropsis oceanica* CCMP1779. *PLoS Genet.* 8 (11) <https://doi.org/10.1371/journal.pgen.1003064>.
- Vogler, B.W., Starkeburg, S.R., Sudasinghe, N., Schambach, J.Y., Rollin, J.A., Pattathil, S., Barry, A.N., 2018. Characterization of plant carbon substrate utilization by *Auxenochlorella protothecoides*. *Algal Res.* 34 (February), 37–48. <https://doi.org/10.1016/j.algal.2018.07.001>.
- Vogler, B.W., Ashford, A., Posewitz, M.C., 2021. CRISPR/Cas9 disruption of glucan synthase in *Nannochloropsis gaditana* attenuates accumulation of β-1,3-glucose oligomers. *Algal Res.* 58 (June), 102385 <https://doi.org/10.1016/j.algal.2021.102385>.
- Wang, L., Wang, H., Chen, X., Zhuang, Y., Yu, Z., Zhou, T.J., 2018a. Acclimation process of cultivating *Chlorella vulgaris* in toxic excess sludge extract and its response mechanism. *Sci. Total Environ.* 628–629, 858–869. <https://doi.org/10.1016/j.scitotenv.2018.02.020>.
- Wang, N., Qian, Z., Luo, M., Fan, S., Zhang, X., Zhang, L., 2018b. Identification of salt stress responding genes using transcriptome analysis in green alga *Chlamydomonas reinhardtii*. *Int. J. Mol. Sci.* 19 (11) <https://doi.org/10.3390/ijms19113359>.
- Wang, X., Luo, S.W., Luo, W., Yang, W.D., Liu, J.S., Li, H.Y., 2019. Adaptive evolution of microalgal strains empowered by fulvic acid for enhanced polyunsaturated fatty acid

- production. *Bioresour. Technol.* 277 (November 2018), 204–210. <https://doi.org/10.1016/j.biortech.2018.12.116>.
- Wang, D., Yu, X., Xu, K., Bi, G., Cao, M., Zelzion, E., Fu, C., Sun, P., Liu, Y., Kong, F., Du, G., Tang, X., Yang, R., Wang, J., Tang, L., Wang, L., Zhao, Y., Ge, Y., Zhuang, Y., Mao, Y., 2020a. *Pyropia yezoensis* genome reveals diverse mechanisms of carbon acquisition in the intertidal environment. *Nat. Commun.* 11 (1), 1–11. <https://doi.org/10.1038/s41467-020-17689-1>.
- Wang, T., Hu, Y., Zhu, M., Yin, S., 2020b. Integrated transcriptome and physiology analysis of *Microcystis aeruginosa* after exposure to copper sulfate. *J. Oceanol. Limnol.* 38 (1), 102–113. <https://doi.org/10.1007/s00343-019-8357-9>.
- Wang, Q., Gong, Y., He, Y., Xin, Y., Lv, N., Du, X., Li, Y., Jeong, B., ryool, Xu, J., 2021. Genome engineering of *Nannochloropsis* with hundred-Kilobase fragment deletions by Cas9 cleavages. *Plant J.* 106, 1148–1162. <https://doi.org/10.1111/tpj.15227>.
- Wang, C., Wang, J., Li, L., Wang, Y., Lin, S., 2022a. P-limitation promotes carbon accumulation and sinking of *Emiliania huxleyi* through transcriptomic reprogramming. *Front. Mar. Sci.* 9 (May), 1–11. <https://doi.org/10.3389/fmars.2022.860222>.
- Wang, J., Zhang, Q., Chen, J., Zhou, J., Li, J., Wei, Y., Regmi, B., Bu, D., 2022b. Combined transcriptome and metabolome analysis of a new species of microalgae from the Tibetan plateau and its response to sewage treatment. *Water (Switzerland)* 14 (21). <https://doi.org/10.3390/w14213391>.
- Wannathong, T., Waterhouse, J.C., Young, R.E., Economou, C.K., Purton, S., 2016. New tools for chloroplast genetic engineering allow the synthesis of human growth hormone in the green alga *Chlamydomonas reinhardtii*. *Appl. Microbiol. Biotechnol.* 100, 5467–5477. <https://doi.org/10.1007/s00253-016-7354-6>.
- Wei, L., El Hajjam, M., Shen, C., You, W., Lu, Y., Li, J., Jing, X., Hu, Q., 2019. Biotechnology for Biofuels transcriptomic and proteomic responses to very low CO<sub>2</sub> suggest multiple carbon concentrating mechanisms in *Nannochloropsis oceanica*. *Biotechnol. Biofuels* 1–21. <https://doi.org/10.1186/s13068-019-1506-8>.
- Wei, L., Jiang, Z., Liu, B., 2022. A CRISPR/dCas9-based transcription activated system developed in marine microalga *Nannochloropsis oceanica*. *Aquaculture* 546 (June 2022), 737064. <https://doi.org/10.1016/j.aquaculture.2021.737064>.
- Williamme, R., Bogaert, K.A., Remacle, F., Remacle, C., 2018. Surprisal analysis of the transcriptomic response of the green microalga *Chlamydomonas* to the addition of acetate during day/night cycles. *Chem. Phys.* 514, 154–163. <https://doi.org/10.1016/j.chemphys.2018.04.015>.
- Worden, A.Z., Lee, J., Mock, T., Rouzé, P., Simmons, M.P., Aerts, A.L., Allen, A.E., Cuvelier, M.L., Derelle, E., Everett, M.V., Foulon, E., Grimwood, J., Gundlach, H., Henrissat, B., Napoli, C., Badger, J.H., Coutinho, P.M., Demir, E., Dubchak, I., Poliakov, A., 2009. The marine *Piceoukaryotes Micromonas*. *Science* 325.
- Xi, Y., Yin, L., Chi, Z., you, & Luo, G., 2021. Characterization and RNA-seq transcriptomic analysis of a *Scenedesmus obliquus* mutant with enhanced photosynthesis efficiency and lipid productivity. *Sci. Rep.* 11 (1), 1–12. <https://doi.org/10.1038/s41598-021-88954-6>.
- Xie, S., Lin, F., Zhao, X., Gao, G., 2022. Enhanced lipid productivity coupled with carbon and nitrogen removal of the diatom *Skeletonema costatum* cultured in the high CO<sub>2</sub> level. *Algal Res.* 61, 102589.
- Xing, G.L., Yuan, H.L., Yang, J.S., Li, J.Y., Gao, Q.X., Li, W.L., Wang, E.T., 2018. Integrated analyses of transcriptome, proteome and fatty acid profilings of the oleaginous microalga *Auxenochlorella protothecoides* UTEX 2341 reveal differential reprogramming of fatty acid metabolism in response to low and high temperatures. *Algal Res.* 33 (May), 16–27. <https://doi.org/10.1016/j.algal.2018.04.028>.
- Yan, N., Fan, C., Chen, Y., Hu, Z., 2016. The potential for microalgae as bioreactors to produce pharmaceuticals. *Int. J. Mol. Sci.* 1–24 <https://doi.org/10.3390/ijms17060962>.
- Yang, M., Xie, X., Kong, F., Xie, K., Yu, S., Ma, J., 2022. Differences in Glycerolipid response of *Chlamydomonas reinhardtii* Starchless mutant to high light and nitrogen deprivation stress under three carbon supply regimes. *Front. Plant Sci.* 13 (May) <https://doi.org/10.3389/fpls.2022.860966>.
- Yoshimitsu, Y., Abe, J., Harayama, S., 2018. Cas9-guide RNA ribonucleoprotein-induced genome editing in the industrial green alga *Coccomyxa sp.* strain KJ 06 biological sciences 0604 genetics. *Biotechnol. Biofuels* 11 (1), 1–10. <https://doi.org/10.1186/s13068-018-1327-1>.
- Yu, Z., Zhang, Q., Peng, Z., Miao, X., 2019. Lipid droplets mediate salt stress tolerance in *Parachlorella kessleri*. *Plant Physiol.* 181 (October), 510–526. <https://doi.org/10.1104/pp.19.00666>.
- Yu, Z., Zhang, T., Zhu, Y., 2020. Whole-genome re-sequencing and transcriptome reveal cadmium tolerance related genes and pathways in *Chlamydomonas reinhardtii*. *Ecotoxicol. Environ. Saf.* 191 (January), 110231 <https://doi.org/10.1016/j.ecoenv.2020.110231>.
- Yuan, G., Xu, X., Zhang, W., Zhang, W., Cui, Y., Qin, S., Liu, T., 2019. Biolistic transformation of *Haematococcus pluvialis* with constructs based on the flanking sequences of its endogenous alpha tubulin gene. *Front. Microbiol.* 10 (JULY), 1–13. <https://doi.org/10.3389/fmicb.2019.01749>.
- Yusibov, V., Kushnir, N., Streatfield, S.J., 2016. Antibody production in plants and green algae. *Annu. Rev. Plant Biol.* 67, 669–701. <https://doi.org/10.1146/annurev-arplant-043015-111812>.
- Zeraatkar, A.K., Ahmadzadeh, H., Talebi, A.F., Moheimani, N.R., McHenry, M.P., 2016. Potential use of algae for heavy metal bioremediation, a critical review. *J. Environ. Manag.* 181, 817–831. <https://doi.org/10.1016/j.jenvman.2016.06.059>.
- Zhao, T., Li, G., Mi, S., Li, S., Hannon, G.J., Wang, X., 2007. A complex system of small RNAs in the unicellular green alga *Chlamydomonas reinhardtii*. *Genes Dev.* 21, 1190–1203 <https://doi.org/10.1101/gad.1543507>.
- Zheng, C., Aslam, M., Liu, X., Du, H., Xie, X., 2020. Impact of Pb on *Chlamydomonas reinhardtii* at physiological and transcriptional levels. *Front. Microbiol.* 11 (June), 1–14. <https://doi.org/10.3389/fmicb.2020.01443>.
- Zhu, B., Chen, G., Cao, X., Wei, D., 2017. Molecular characterization of CO<sub>2</sub> sequestration and assimilation in microalgae and its biotechnological applications. *Bioresour. Technol.* 244, 1207–1215. <https://doi.org/10.1016/j.biortech.2017.05.199>.
- Zhu, Q.L., Guo, S.N., Wen, F., Zhang, X.L., Wang, C.C., Si, L.F., Zheng, J.L., Liu, J., 2019. Transcriptional and physiological responses of *Dunaliella salina* to cadmium reveals time-dependent turnover of ribosome, photosystem, and ROS-scavenging pathways. *Aquat. Toxicol.* 207 (October 2018), 153–162. <https://doi.org/10.1016/j.aquatox.2018.12.007>.
- Zhu, Q., Zhang, M., Bao, J., Liu, J., 2021. Ecotoxicology and Environmental Safety of *Dunaliella salina*: Physiological, metabolomic, and transcriptomic analyses reveal the dynamic redox homeostasis upon extended exposure of *Dunaliella salina* GY-H13 cells to Cd. *Ecotoxicol. Environ. Saf.* 223, 112593 <https://doi.org/10.1016/j.ecoenv.2021.112593>.
- Zhu, Q., Zhang, M., Liu, B., Wen, F., Yang, Z., Ceballos, R.M., 2022. Transcriptome and metabolome profiling of a novel isolate *Chlorella sorokiniana* G32 (Chlorophyta) displaying enhanced starch accumulation at high growth rate under Mixotrophic condition. *Front. Microbiol.* 12 (January), 1–15. <https://doi.org/10.3389/fmicb.2021.760307>.
- Zienkiewicz, K., Du, Z.Y., Ma, W., Vollheyde, K., Benning, C., 2016. Stress-induced neutral lipid biosynthesis in microalgae — molecular, cellular and physiological insights. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* 1861 (9), 1269–1281. <https://doi.org/10.1016/j.bbalip.2016.02.008>.
- Zuñiga, C., Levering, J., Antoniewicz, M.R., Guarneri, M.T., Betenbaugh, M.J., Zengler, K., 2018. Predicting dynamic metabolic demands in the photosynthetic eukaryote *Chlorella vulgaris*. *Plant Physiol.* 176 (1), 450–462. <https://doi.org/10.1104/pp.17.00605>.