

# SEFS14



**14th Symposium for European  
Freshwater Sciences**

July 20-25, 2025, Bolu, Türkiye

# ABSTRACT BOOK

## **Plastic entrapment by riparian vegetation across ecological gradients in European rivers: first insights from the Biodiversa+ RIPARIANET Project**

Luca Gallitelli<sup>1</sup>, Giorgio Pace<sup>2,3</sup>, Maria Cristina Bruno<sup>4,5</sup>, Jose Barquin<sup>6</sup>, Giulia Cesarini<sup>1</sup>, Laura Concostrina Zubiri<sup>6</sup>, Micael Jonsson<sup>7</sup>, Stefano Larsen<sup>4,5</sup>, Monika Laux<sup>8</sup>, Ralf Schulz<sup>8</sup>, Massimiliano Scalici<sup>1,5</sup>

<sup>1</sup>University of Roma Tre, Department of Sciences, Laboratory of Ecology (Rome, Italy)

<sup>2</sup>Centre of Molecular and Environmental Biology (CBMA) / Aquatic Research Network (ARNET) Associate Laboratory, Department of Biology, University of Minho, Braga, Portugal.

<sup>3</sup>Institute of Science and Innovation for Bio-Sustainability (IB-S), University of Minho, Braga, Portugal.

<sup>4</sup>Research and Innovation Centre, Fondazione Edmund Mach, Via Mach 1, 38098 S. Michele all'Adige, Italy

<sup>5</sup>National Biodiversity Future Center (NBFC), Università di Palermo, Piazza Marina 61, 90133 Palermo, Italy

<sup>6</sup>HCantabria, Instituto de Hidráulica Ambiental, Universidad de Cantabria, Santander, Spain

<sup>7</sup>Department of Ecology and Environmental Science, Umeå University, Umeå, Sweden

<sup>8</sup>Institute for Environmental Sciences, RPTU – University of Kaiserslautern-Landau, Landau, Germany

**Presenting Author:** Luca Gallitelli

**Status:** Oral presentation

**Session Number & Date:** Advancing research on plastic pollution: Impacts, transport, and management, 10:45-11:00, Monday 21 July

### **Abstract**

Plastic litter accumulating in riverine riparian habitats is a threat of global concern. Macrolitter items, being highly visible (items > 0.5 cm) and impactful pollutants, pose significant threats to biodiversity and ecosystem functioning. Although research recently started to address plastic entrapment, large-scale studies and predictive models aiming at reducing biases and uncertainties in understanding plastic accumulation are still scarce. Given those gaps, this study investigates plastic entrapment by riparian vegetation at different ecological scales and gradients across European rivers. We focused on six river basins across Europe, covering the boreal (Sweden), continental (Germany), alpine (Trento, Italy), Mediterranean (Rome, Italy), and Atlantic (Northern Spain, Northern Portugal) climatic regions as part of the European project Biodiversa+ RIPARIANET. By surveying six European basins, we aim to unveil riverine macrolitter accumulation in riparian areas across biogeographic regions. We found that riparian vegetation acts as a sink for macrolitter, with the highest trapping value recorded in the Tiber River catchment (Italy) and the lowest in the Sävar River basin (Sweden). Among river basins, we highlighted a latitudinal gradient for plastic entrapment by vegetation, which increased from North to South. Among macrolitter items, we found that plastics was the most abundant litter type, followed by textile items. Additionally, we observed most macroplastics near the downstream zone of rivers rather than the upperstream zone.

Urbanization, land use, river discharge, river sinuosity, and vegetation structure are crucial predictors of macroplastic accumulation. Our findings shed light on how macroplastics accumulate in European riparian zones, emphasizing their ecological and societal implications and potentially supporting environmental managers in addressing macrolitter removal from the environment. Monitoring macroplastic accumulation is essential to understand the interactions between pollutants and ecosystems, enabling the development of effective conservation strategies. Given the potential impacts on biodiversity and ecosystem resilience, specific monitoring and clean-up activities should be prioritized to protect riparian ecosystems under future conditions.

