



Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) in Europe: an overview of management practices

Valeriu-Norocel Nicolescu¹ · William L. Mason² · Jean-Charles Bastien³ · Torsten Vor⁴ · Krasimira Petkova⁵ · Vilém Podrázský⁶ · Martina Đodan⁷ · Sanja Perić⁷ · Nicola La Porta⁸ · Robert Brus⁹ · Siniša Andrašev¹⁰ · Martin Slávik¹¹ · Juraj Modranský¹² · Michal Pástor^{11,12} · Károly Rédei¹³ · Branislav Cvjetkovic¹⁴ · Ahmet Sivacioglu¹⁵ · Vasyl Lavnyy¹⁶ · Cornelia Buzatu-Goanța¹⁷ · Gheorghe Mihăilescu^{1,18}

Received: 1 September 2022 / Accepted: 30 January 2023 / Published online: 22 March 2023
© The Author(s) 2023, corrected publication 2023

Abstract Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco), one of the most commercially important tree species in western North America and one of the most valuable timber trees worldwide, was introduced to Europe in 1827. It became a major species for afforestation in Western Europe after WWII, currently grows in 35 countries on over 0.83 million ha and is one of the most widespread non-native tree species across the continent. A lower sensitivity to drought makes Douglas-fir a potential alternative to the more drought-sensitive Norway spruce so its importance in Europe is expected to increase in the future. It is one of the

fastest growing conifer species cultivated in Europe, with the largest reported dimensions of 2.3 m in diameter and 67.5 m in height. Pure stands have high productivity (up to 20 m³ ha⁻¹a⁻¹) and production (over 1000 m³ ha⁻¹). The species is generally regenerated by planting (initial stocking density from less than 1000 seedlings ha⁻¹ to more than 4000 ha⁻¹), using seedlings of European provenance derived from seed orchards or certified seed stands. As the range of end-uses of its wood is very wide, the rotation period of Douglas-fir is highly variable and ranges between 40 and 120 years. When the production of large-sized, knot-free timber is targeted, thinnings are always coupled with pruning up to 6 m. There is an increasing interest in growing Douglas-fir in mixtures and managing stands through

The online version is available at <http://link.springer.com>.

Corresponding editor: Yu Lei.

✉ Valeriu-Norocel Nicolescu
vnnicolescu@gmail.com; vnnicolescu@unitbv.ro

¹ Faculty of Silviculture and Forest Engineering, Transylvania University of Brasov, Brasov, Romania

² Forest Research, Northern Research Station, Roslin, Midlothian, UK

³ INRAE Centre Val de Loire, Ardon, Orléans, France

⁴ Faculty of Forest Sciences and Forest Ecology, University Georg-August, Göttingen, Germany

⁵ University of Forestry, Sofia, Bulgaria

⁶ Department of Silviculture, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, Prague, Czech Republic

⁷ Croatian Forest Research Institute, Jastrebarsko, Croatia

⁸ Foundation Edmund Mach, San Michele all'Adige, Italy

⁹ Biotechnical Faculty, University of Ljubljana, Ljubljana, Slovenia

¹⁰ Institute of Lowland Forestry and Environment, Novi Sad, Serbia

¹¹ National Forest Centre, Forest Research Institute, Zvolen, Slovakia

¹² Faculty of Ecology and Environmental Sciences, Technical University in Zvolen, Zvolen, Slovakia

¹³ Faculty of Agricultural and Food Sciences and Environmental Management, University of Debrecen, Debrecen, Hungary

¹⁴ Faculty of Forestry, University of Banja Luka, Banja Luka, Bosnia and Herzegovina

¹⁵ Department of Silviculture, Faculty of Forestry, Kastamonu University, Kastamonu, Turkey

¹⁶ Department of Silviculture, Ukrainian National Forestry University, Lviv, Ukraine

¹⁷ Faculty of Horticulture and Forestry, Banat's University of Agricultural Sciences and Veterinary Medicine, Timisoara, Romania

¹⁸ Vâlcea County Branch, National Forest Administration-ROMSILVA, Râmnicu-Vâlcea, Romania

close-to-nature silviculture, but the species' intermediate shade tolerance means that it is best managed through group selection or shelterwood systems.

Keywords Douglas-fir · Ecological requirements · Growth and yield · Timber · Climate change

Introduction

Since the arrival of black locust (*Robinia pseudoacacia* L.) in 1601, a range of non-native tree species, particularly from North America and Asia, have been introduced to Europe to increase the attractiveness of landscapes, to augment the productivity of native forests, and to improve the profitability of forest management. Currently, 145 non-native or exotic tree species can be found growing in European forests and together cover an area of 8.54 million ha, or 4.0% of the continent's forested area (Brus et al. 2019). One of these species, Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) is one of the most commercially important tree species in western North America and one of the most important and valuable timber species worldwide (Hermann and Lavender 1999). The native range of Douglas-fir stretches over 3400 km from south to north (latitude 19° to 55° N.), the largest latitudinal spread of any conifer of western North America. It also extends over 1600 km from east to west, covering 14.3 million hectares in the U.S.A. and 4.5 million hectares in Canada (Hermann and Lavender 1990; Fletcher and Samuel 2010). Two geographic varieties are recognised: the coastal variety or green Douglas-fir (*P. menziesii* var. *menziesii*), and the interior variety (*P. menziesii* var. *glauca* (Beissn.) Franco) also called Rocky Mountain or blue Douglas-fir.

The species was introduced to Europe by David Douglas in 1827 when the first trees were planted at Scone Palace in Perthshire, Scotland (Haralamb 1967). This was followed by introductions to many European countries, with the first plantations occurring between 1851 and 1900 (van Loo and Dobrowolska 2019a). The coastal variety of Douglas-fir has proved much better adapted to European conditions than the interior variety, having a higher growth rate, being resilient to frost and more resistant to fungal diseases (Bastien et al. 2013; Lavender and Hermann 2014; Petkova et al. 2014; Konnert 2016), so green Douglas-fir is preferred throughout the continent (Konnert and Bastien 2019). In Western Europe, Douglas-fir was widely used in afforestation programs after the Second World War, and such plantings were supported in national and regional subsidy schemes (Bastien et al. 2013; Spiecker 2019). Eighty per cent of these plantations were established in three countries: France (half of all European plantings), Germany and the UK (da Ronch et al. 2016). In France, the area of Douglas-fir was ca. 4000 ha

in 1937 (Haralamb 1967), 220,000 ha in 1984 (Bouchon 1984, in Hermann and Lavender 1999), 330,000 ha in 1993 (Bastien 1998) and 411,000 ± 31,000 ha today (<https://inventaire-forestier.ign.fr>). In Germany, Douglas-fir covered ca. 90,000 ha in the mid-1980's (Braun and Weissleder 1986, in Hermann and Lavender 1999), increasing to 134,000 ha in mid-1990s and 217 600 ha at present (BMEL 2014). Currently, Douglas-fir is grown in 35 European countries and covers over 0.83 million ha, being the second most widespread non-native conifer species in Europe after Sitka spruce (*Picea sitchensis* (Bong.) Carr) (van Loo and Dobrowolska 2019b). In the southern hemisphere, Douglas-fir has been planted in New Zealand (112,000 ha-the second most important introduced tree species in the country-Dungey et al. 2010), Chile (15,000 ha-Bastien et al. 2013), and in Argentina (7500 ha-Bastien et al. 2013).

Currently there is no comprehensive summary of the silviculture and management of Douglas-fir in Europe that can be used to indicate its future role in different countries. Our paper provides a short overview of key factors influencing the management of Douglas-fir and summarises European silvicultural experience with the species, covering aspects such as stand establishment, early silvicultural interventions such as release cutting and cleaning-respacing, commercial thinning and pruning, as well as the potential use of close-to-nature silvicultural regimes. The focus is upon the impacts upon timber production, since effects upon wider ecosystem services have been examined by others (e.g., Wohlgenuth et al. 2019, 2021; Thomas et al. 2022).

Douglas-fir characteristics influencing management

Site and climate

European experience indicates that Douglas-fir can grow on a range of soil types apart from heavy soils with pseudogley close to the surface, on dry or heavily waterlogged soils where rooting is restricted, leading to instability, and on calcareous soils. It grows best on deep, well-aerated, moderately acidic (pH 5–6), free draining soils of loamy-sand or sandy-loam texture which allow good root development (Rameau et al. 1989; Riou-Nivert 1996; Horgan et al. 2003; Perić et al. 2011; Spellmann et al. 2015; Novák et al. 2018; Eckhart et al. 2019; Savill 2019; Čater 2021).

In continental Europe, Douglas-fir grows best at low to middle altitudes (between 300 and 900 m asl) in oak-dominated and European beech-dominated zones. However, it also grows at lower (200 m asl) and higher elevations (up to 1300 m asl in Slovakia and 1600 m asl in France; Coopérative Forestière Bois Limousin 2016), but with lower yields. The best climatic conditions for growing Douglas-fir in

Europe are in moderate to moist temperate climates, with mean annual temperatures of 7–8 °C (Czech Republic: Mondek and Balaš 2019), 7–9 °C (Romania: Stănescu et al. 1997), 8–11 °C (France: Coopérative Forestière Bois Limousin 2016), 9–11 °C (Slovenia: Čater 2021; Smolnikar et al. 2021). The most important limiting factor for successful cultivation of the species in Europe is the minimum mean annual precipitation, with a critical lower threshold between 600 mm (Czech Republic: Mondek and Balaš 2019; Romania: Haralamb 1967; Slovakia: Slávik pers.comm.) and 750 mm (France: de Champs and Demarq 1996; Bastien 1998). A second limiting factor is the minimum annual precipitation during the growing season, defined as 350 mm (Germany: Spellmann et al. 2015) or 450 mm (Romania: Marcu and Liubimirescu 1979).

An important characteristic of Douglas-fir, in the context of expected climate changes, is a greater tolerance of summer drought than other species on a wide variety of sites and in different climatic conditions (Wohlgemuth et al. 2021), especially in areas with at least 700 mm precipitation and on soils with good water reserves (CRPF 1999; Nicolescu 2019). In western (Germany, Switzerland, Belgium), central (Czech Republic) and eastern (Romania) parts of Europe, Douglas-fir has proved less sensitive to drought than Norway spruce (*Picea abies* (L.) Karst), European silver fir (*Abies alba* Mill.), Scots pine (*Pinus sylvestris* L.), larch (*Larix* spp.), oaks (*Quercus* spp.), European beech (*Fagus sylvatica* L.) (Eilmann and Rigling 2010, 2012; Manise and Vincke 2014; Nicolescu 2019; Spiecker 2019). Thus, coastal Douglas-fir could be a replacement for Norway spruce, providing high productivity and resilience to climate change (Chakraborty et al. 2019; Covre Foltran et al. 2020; Frei et al. 2022). However, Douglas-fir stands in Europe have suffered reduced growth and dieback during extreme drought

conditions, especially at lower elevations (Bastien 2019). Thus, new growing areas may be at medium and high altitudes (Schüler and Chakraborty 2021), with optimum sites possibly moving to higher altitudes (over 1000 m asl) (Wohlgemuth et al. 2021). Similar concerns are reported from its native range where water deficit-related stress is predicted to increase in the western United States and consequently Douglas-fir growth will decrease (Restaino et al. 2016).

Growth performance in Europe

Douglas-fir grows to substantial heights, for instance in parts of its native range (e.g., Oregon, USA), a number of trees > 90 m height have been recorded (Sillett et al. 2018). In Europe, although most Douglas-firs are young compared to long-lived specimens in the native range (> 1000 years; Cline et al. 1980; in Weiskittel et al. 2012), impressive heights and diameters have been attained (Table 1).

Mean annual volume increment in even-aged stands in Europe peaks at between 50 and 80 years and diminishes gradually afterwards (Schütz et al. 2015). Increments recorded in different countries range from 10 to 20 m³ ha⁻¹a⁻¹ (e.g., 14.8 in France, 18.9 in Germany; Kohnle et al. 2019), with still higher increments reported (Thomas et al. 2022). These values are significantly higher compared to European native forest species. The average annual increment of Douglas-fir exceeds that of other conifers by 76% and 47% in France and Germany respectively. In its native range, coastal Douglas-fir produces over 13 m³ ha⁻¹a⁻¹ on average (Talbert and Marshall 2005), while in New Zealand, the average annual volume increment reaches 16 m³ ha⁻¹a⁻¹ (<http://www.nzffa.org.nz>). These growth rates lead to high

Table 1 Maximum heights and diameters of Douglas-fir in different European countries

Height, m	Diameter (cm)	Country	Source	Observations
30		Serbia	Ratkníč (1995)	50 years old
40	100	Hungary	Rédei pers.comm	
43.1	38.4	Italy	Scotti (2016)	50 years old
44	80	Ukraine	Lavnyy pers.comm	
51	102	Bulgaria	Petkova et al. (2017)	100 years old
49.7		The Netherlands	http://www.bomeninfo.nl/tall%20trees.htm	National record
57.9		Austria	Jasser (2008)	103 years old, national record
64	80	Czech Republic	Podrázský pers.comm	148 years old; tallest tree in the country
Over 66.5	> 160	Germany	https://ddg-web.de/rekordbaeume.html?VCardId=6735	
67	118	France	Angelier (2007)	124 years old; tallest tree in the country
67.0	150	Slovenia	Hostnik (2021)	
67.5	111	UK	https://www.treeregister.org/champion	Wales, planted in 1921
38.5	233	UK	https://www.treeregister.org/champion	Scotland, planted in 1846

standing volumes that can reach 1000 m³ ha⁻¹ or more at 80 years of age (Table 2).

Timber: quality, uses, prices

Douglas-fir timber is better quality than that of Norway spruce and European silver fir and similar to European larch (*Larix decidua* L.) (Stănescu 1979; Zeidler et al. 2022). It is moderately hard, stiff, moderately stable, fast to normal drying, resistant to fungi and insect attacks and moderately resistant to rot (Riou-Nivert 1989, 1996; de Champs and Demarq 1996; Molnár 2008; Bastien et al. 2013; Pollet et al. 2013; France Bois Forêt 2016; Kahl et al. 2017; Podrázský et al. 2019). The timber is difficult to impregnate with preservatives, insecticides and fungicides (Liese et al. 1982; Suzuki et al. 1995).

In Europe, the average density of Douglas-fir ranges between 360 kg m⁻³ and 583 kg m⁻³ at 12% moisture content (Table 3). In its native range, it is 480 kg m⁻³ (Interior North: Alden 1997) and 500 kg m⁻³ (Interior West: Alden 1997) with higher densities in coastal areas (540 kg m⁻³) (Remeš and Zeidler 2014).

The maximum ring width for Douglas-fir timber to be considered the best grade /highest strength class ranges from 4 mm (Poland, The Netherlands) and 6 mm (Austria, Belgium, France, Germany, Great Britain), or even 8 mm in special circumstances (France) (Henin et al. 2019). High growth rates in this species do not appear to result in a reduction in density and associated wood properties (Pollet et al. 2017). Therefore, timber from trees grown at wide spacing in plantations should not be rejected for general use (Melin and Riou-Nivert 1985; in Pollet et al. 2017). However, for structural uses, a ring width of 5 mm to guarantee the best

Table 2 Maximum standing volumes of Douglas-fir stands in different European countries

Total standing volume (m ³ ha ⁻¹)	Country	Source	Observations
600–800	Europe-wide	Bastien et al. (2013)	40 years of age; production depending on site potential
603	Slovenia	Smolnikar et al. (2021)	46 years old
606	Bosnia and Herzegovina	Ibrahimspahić et al. (2006)	46 years old, mean dbh 29.9 cm, mean height 24.1 m
670	Romania	Negulescu and Săvulescu (1957, 1965)	40 years old; Norway spruce standing volume at the same age 310 m ³ ha ⁻¹
678	Croatia	Klepac (1962)	70 years old, mean dbh 43.5 cm
700–750	Italy	La Marca et al. (2016)	45 years old
868	Hungary	Redei pers.comm	50 years old, mean dbh 38.5 cm, mean height 32.2 m
988	Bulgaria	Popov (1991)	80 years old, mean dbh 46 cm, mean height 33 m
1153	Slovenia	Čokl (1965)	62 years old
1166	Bulgaria	Popov (2006, 2009)	82 years old, mean dbh 43 cm, mean height 38 m
1233	Bulgaria	Popov (2006)	71 years old, mean height 33 m

Table 3 Average Douglas-fir density reported in European studies

Average density, kg m ⁻³	Country	Source
360–480	Ireland, Scotland	Gil-Moreno et al. (2019)
425	Slovenia	Možina (1960)
440–513	Bulgaria	Bluskova (2006)
440–600	Italy	La Porta pers.comm
488	Czech Republic	Remeš and Zeidler (2014)
500	Belgium, France, Germany, Great Britain, Italy, Poland	Henin et al. (2019)
	Slovakia	(Ťavoda 2007; Petráš and Mecko 2008)
	France	France Bois Forêt (2016)
533–575	Czech Republic	(Giagli et al. 2019; Podrázský et al. 2019; Zeidler et al. 2022)
540	France	http://www.frenchtimber.com/en/french-species/douglas-fir/
570	The Netherlands	Polman and Militz (1996)
583	Serbia	Šoškić et al. (2007)

mechanical properties should not be exceeded (Nepveu and Blachon 1989; in Pollet et al. 2017). Users of Douglas-fir require ring wood regularity, an aspect of equal importance to average ring width, as irregular ring wood is more prone to distortion (Henin et al. 2019).

Because of its excellent mechanical properties, combined with a relatively low density and dimensional stability, Douglas-fir is particularly well-suited to structural applications such as timber frames, floor and roof trusses, glue-laminated beams, and flooring. As a result, in Western Europe, the price of Douglas-fir timber is on average 25% higher than Norway spruce (Pulkrab et al. 2014). For example, in France in 2020, the average price of Douglas-fir timber (1.0 m³ standing volume) was 61 € m⁻³ compared with 36 € m⁻³ for Norway spruce of the same volume (https://www.forestiere-cdc.fr/sites/default/files/2021-06/prix-de-vente-des-bois-sur-pied-en-foret-privee-indicateur-2021_0.pdf). Similarly, in Belgium in 2021–2022, the price for timber 50 cm diameter was 90–110 € m⁻³ for Douglas-fir compared to 60–80 € m⁻³ for Norway spruce (www.experts-forestiers.be/Tableauprixbois.pdf). Prices for Douglas-fir logs for veneer often exceed 110 € m⁻³ and can reach as high as 180 and 250 € m⁻³ (Germany-<https://www.forstpraxis.de/douglasieschwarzwald-rekordmenge-rekorderloes/>; https://www.landeforsten.de/wp-content/uploads/2021/03/ergebnisse_submission_nadelholz_2021.pdf) while a record 527 € m⁻³ was registered in Slovenia in 2021 (Gozd in gozdarstvo 2021).

With the attributes of good growth, desirable wood properties, and resistance to drought, Douglas-fir is attractive for the forest industry in many European countries (Spiecker 2019; Covre Foltran et al. 2020; Forest Europe 2020; Frei et al. 2022).

Root system

Douglas-fir is a deep-rooting species, with a tap root down to 1.5 m or to 3.2 m (Köstler et al. 1968; in Thomas et al. 2022) in favourable soils. In such soils, Douglas-fir develops a tap root that grows rapidly to about 50% of its final length in 3 to 5 years, and to 90% in 6 to 8 years (Hermann and Lavender 1990). Main lateral roots develop during the first or second growing season as branches of the tap root grow obliquely into deeper soil layers and provide anchorage (Hermann and Lavender 1990; Mauer and Palátová 2012). On well draining soils, Douglas-fir has better anchorage than spruce (Nicoll et al. 2006) and is less vulnerable to wind (Paşcovschi and Purcelean 1954; de Champs 1988; Mauer and Palátová 2012). However, when grown on shallow or poorly drained soils, plate-like root systems develop. Under such conditions, the species is prone to windthrow (Haralamb 1967; Hermann and Lavender 1990; Hart 1994; Sychra and Mauer 2013; Savill 2019) and more prone than European silver fir but perhaps less than Norway spruce

(Haralamb 1967). Therefore, on exposed sites, the planting of Douglas-fir should be avoided as the trees can be uprooted or become badly deformed (Wouters and Lorent 2002; Savill 2019).

Shade tolerance

In its natural range, except in youth when it is reasonably shade tolerant, coastal Douglas-fir is intermediate in shade tolerance (Herman and Lavender 1990) or shade intolerant compared to associate species (Larson 2010). This characteristic is affected by site quality, with decreasing tolerance with increasing soil moisture (Carter and Klinka 1992). European studies generally consider the species as having ‘intermediate’ tolerance (Niinemets and Valladares 2006) with recruitment being more light-demanding than European silver fir and Norway spruce (Schütz et al. 2015), while mature trees are more shade tolerant than sessile oak (*Quercus petraea* (Matt.) Liebl.), pedunculate oak (*Quercus robur* L.) and Scots pine (Thomas et al. 2022). Seedlings require a minimum of 15% (Mason et al. 2004) or 20 to 40% open sky light intensity for establishment and growth (Mailly and Kimmins 1997; Drever and Lertzman 2003; Herrington 2006; all in Frei et al. 2022). Thus, Douglas-fir is sufficiently shade-tolerant to be planted beneath well-thinned canopies; underplanting is likely to be more successful under a sparse canopy (Stokes et al. 2021).

Regeneration ecology

Douglas-fir trees start to produce seed cones at 30–35 years (Şofletea and Curtu 2007; Savill 2019), with the best production occurring from 50–60 years. Mast years occur every 2–3 years (Romania: Stănescu 1979), 4–6 years (UK: Savill 2019), 7 years (Germany: Spellmann et al. 2015) or irregularly (one heavy and one medium crop every 7 years in Ireland; COFORD 2020). Seed production is often higher in years following a dry summer (Van Vredenburg and La Bastide 1968). The seeds are dispersed by wind for distances of 25–30 m (Bulgaria, Milenkova 2020) up to 50 to 100 m (Czech Republic: Podrázský pers. comm; France: Bous-said 2008; Germany: Stimm 2004; Lange et al. 2022) or even 100–150 m (Switzerland: Wohlgemuth et al. 2019). Consequently, in stands with a high proportion of Douglas-fir, natural regeneration is common e.g., Britain (Jones 1945), Romania (Dumitriu-Tătăranu 1960; Negulescu and Săvulescu 1957), Belgium (Boudru 1989), France (Riou-Nivert 1996; Bastien 1998; CRPF 1999), Czech Republic (Sychra and Mauer 2013), Bulgaria (Popov et al. 2018; Milenkova 2020), Slovakia (Slávik pers. com.), Italy (La Marca and Pozzi 2016). Successful seed germination can be limited by vegetation competition (Malcolm et al. 2001) and favoured by improved light conditions from either

silvicultural treatments reducing canopy cover (La Marca and Pozzi 2016), windthrow creating large gaps in the stand (Raida 2018) or the sparse cover of light-demanding species such as Scots pine, silver birch (*Betula pendula* L.) or European larch (Bindewald et al. 2021).

The abundant natural regeneration of Douglas-fir has caused debate about the invasive potential of the species, although recent reports have considered that it is not invasive in Europe (Brus et al. 2019). This categorization was supported by Eberhard and Hasenauer (2018), Bindewald et al. (2021) and Lange et al. (2022) based on the natural spread of the species being site-limited and easy to manage (e.g., Essl 2005; Ammer et al. 2016; Raida 2018; Wohlgemuth et al. 2019). However, in certain countries (e.g., Germany) there continues to be contrasting opinions about the species invasiveness (Thomas et al. 2022).

Douglas-fir vulnerabilities to abiotic and biotic disturbances

As noted, Douglas-fir can be vulnerable to windthrow, especially if grown on soils that result in shallow rooting. In Germany, the species is considered as prone to windthrow as Norway spruce (Albrecht et al. 2010, 2013), attributed to the heavy weight of the large crowns (Boudru 1989; Hart 1994). Dense crowns also favour snow retention (Dobrev 1962), resulting in stem breakage as a result of heavy loading (Govedar et al. 2003). In Denmark, field studies indicated that Douglas-fir was less susceptible to windthrow than Norway spruce (Decker 2018), in line with the results of the tree pulling experiments conducted in Britain between 1960 and 2000 (Nicoll et al. 2006). Wind risk models now available can be used to predict the vulnerability of even-aged Douglas-fir stands, as well as other conifer species (Hale et al. 2015).

Douglas-fir can be damaged by very low winter temperatures as well as early autumn and late spring frosts (Bosnia and Herzegovina: Pintarić 1990; Croatia: Vidaković and Franjić 2004; Czech Republic: Novák et al. 2018; Ireland: Horgan et al. 2003; COFORD 2020; Romania: Drăcea 1923; Paşcovschi and Purcelean 1954; Ionuţ 1956; Sweden: Malmqvist 2017), especially on low-lying sites with poor air drainage (Ireland: Horgan et al. 2003; UK: MacDonald et al. 1957; in Savill 2019). Fire is an important disturbance affecting the dynamics of stands in its native range where moderate severe fires allow thicker bark Douglas-fir to survive and regenerate at the expense of thin bark species of lower fire resistance (Hermann and Lavender 1990; Sillett et al. 2018).

Douglas-fir is currently less threatened by pests and pathogens in Europe than are the indigenous Norway spruce and Scots pine (Spellmann et al. 2015). Fungal pathogens *Phaeocryptopus* (*Adelopus*) *gaeumannii* Petr. (Swiss needle cast)

and *Rhabdocline pseudotsugae* Sidow (Rhabdocline needle cast) can cause yellowing of foliage, reduction in growth, and eventual drop of needles (Georgieva 2009; Podrázský et al. 2019; Roques et al. 2019; Horgan et al. 2003, in COFORD 2020). Douglas-fir may also show root rot caused by *Heterobasidion* spp, the damage being mostly associated with *H. annosum* (Fr.) Bref (Roques et al. 2019). However, it is less susceptible to this fungus than many other conifers (COFORD 2020). Important insect pests of Douglas-fir are the vapourer moth (*Orgyia antiqua* L.), which severely defoliated Douglas-fir stands in Poland in the mid-1970s, and the nun moth (*Lymantria monacha* L.), with local outbreaks in the 1980s and 1990s, especially in central France. The large pine weevil (*Hylobius abietis* L.) can cause severe damage in young plantations (Spellmann et al. 2015; Roques et al. 2018; Wallertz et al. 2014; in COFORD 2020). Roques et al. (2018) noted that 'compared to other exotic conifers introduced to Europe, Douglas-fir is still relatively free of biotic damage because its phylogenetic distance to native tree species is preventing rapid switches of most native plants. However, climate change and the worldwide movement of plants for planting is likely to accelerate the arrival of pests from the native range'.

In both its natural range and in Europe, Douglas-fir is widely considered prone to game damages (browsing, fraying, especially by roe deer, red deer and wild boar), affecting the young, smooth, soft bark, resin-rich, up to 1–2 m height (Rădulescu and Cazacu 1968; Hermann and Lavender 1990; Riou-Nivert 1996; Spellmann et al. 2015; Podrázský et al. 2019; Savill 2019) (Fig. 1).

When browsing pressure is high, fencing plantations is necessary, as recommended in Bulgaria, Czech Republic, Hungary, Romania, Slovakia, Slovenia, and the UK. The application of chemical repellents (e.g., TRICO) can be effective against hares, and red, roe, fallow and sika deer and is recommended in some countries (Czech Republic: Podrázský pers.comm.; France: CRPF 1999; Slovenia: Brus pers.comm.). Individual protection by treeshelters is not advised other than for small gaps (<0.2 ha), as they are expensive and can produce slender plants with limited root growth, leading to bent stems after storms or breakage under wet snow (Petersen 2016).

Management of Douglas-fir

Even-aged stands

In Europe, Douglas-fir is generally established by using seedlings from a range of sources: these include certified seed stands (over 2200 stands covering about 4800 ha in the EU) and seed orchards (69 orchards, with 390 ha) as well as seed imports from its natural range (Konnert and



Fig. 1 Wild boar damage on Douglas-fir (photo T. Vor)

Bastien 2019). All Douglas-fir provenances recommended for Europe are of the coastal variety and originate from the part of its native range between 40° and 50° N latitude, west of the Cascade Range and below 600 m altitude, e.g., northern Oregon and western Washington states in the USA as well as the Vancouver region in British Columbia, Canada (Isaac-Renton et al. 2014; Chakraborty et al. 2015; Konnert and Bastien 2019).

Plant production, spacing and early establishment

Planting stock of Douglas-fir is produced in bare-root and containerized nurseries in Europe and in its native range (Aldhous and Mason 1994). Production begins with seed pre-treatment involving stratification by chilling for three or more weeks at 3–5 °C and sowing either in October–November (Marcu and Liubimirescu 1979) or in late March (Podrázský et al. 2019; Savill 2019). Bare-root seedlings are generally produced as 2–3 year-old stock around 30–60 cm height and a minimum root collar diameter of 5 mm with a fibrous rooting system; undercutting is recommended to stimulate root production and improve root:shoot balance (Marcu and Liubimirescu 1979; Boudru 1992; Aldhous and Mason 1994; de Champs and Demarq 1996; CRPF 1999; Anonymous 1999; Horgan et al. 2003). Containerised seedlings are normally grown on a 12–24-month cycle, with watering

and shading regimes in greenhouses often adjusted to try and improve quality (Curtis et al. 2007; Turner and Mitchell 2003; Talbert and Marshall 2005).

Douglas-fir is mostly planted in spring (Austria and Germany: Eberhard et al. 2021; Czech Republic: Podrázský et al. 2019; Ireland: Horgan et al. 2003; Romania: Ionescu 1963, Marcu and Liubimirescu 1979; Sweden: Malmqvist et al. 2017). Survival rates of 70%–80% or more can be obtained under good conditions, although failure rates tend to be higher than with some other conifers (Kohnle et al. 2019). Investigations of physiological factors causing post-planting failures revealed that Douglas-fir seedlings were more sensitive to root desiccation and stresses caused by rough handling than many other conifer species (McKay and Milner 2000). Cold storage is used in some regions to extend planting seasons (Aldhous and Mason 1994; O'Reilly et al. 1999). Autumn planting is not recommended as Douglas-fir seedlings are often damaged by game species (red deer, roe deer) (Marcu and Liubimirescu 1979).

The initial stand density used in Douglas-fir plantations will affect the long-term stability of the trees as well as stand productivity and timber quality (Klädtke et al. 2012; Kohnle et al. 2021; both in Eberhard et al. 2021). A lower initial stand density improves tree stability and promotes social differentiation (several authors, in Eberhard et al. 2021). Historically, the initial density recommended for Douglas-fir plantations was high, but this was reduced sharply from the mid-20th century onwards (i.e., in France from 2500 to 3000 plants ha⁻¹ in 1950–1955 to 1100–1300 plants ha⁻¹ at the end of 1980's; Lanier 1986; de Champs 1988; Riou-Nivert 1989; Giraud and Champaux 1997; Bastien 1998). Currently, the initial stand density of Douglas-fir plantations in Europe ranges from 600 plants ha⁻¹ to over 4000 plants ha⁻¹ with most countries using somewhere between 1000 and 2000 stems ha⁻¹ (Table 4).

The initial densities of Douglas-fir plantations in Europe are similar to those in its native range (750–1500 plants ha⁻¹-Cafferata 1986, in Hermann and Lavender 1990; 1100 plants ha⁻¹-Talbert and Marshall 2005) and in New Zealand (1250–1650 plants ha⁻¹-<http://www.nzffa.org.nz>).

Planting in pits is recommended as being the most suitable for all planting stock, preventing the root system from deformation (Ionescu 1963; Mauer and Houšková 2014, in Podrázský et al. 2019). Alternatively, soil cultivation has improved survival and early growth, particularly in more northern countries where cold soil temperatures may limit new root growth (Sharpe et al. 1990; Wallertz and Malmqvist 2013). Early growth and rate of establishment is enhanced when competition from herbaceous and woody weeds is controlled (Rose and Rosner 2005).

Table 4 Initial density and spacing recommended for Douglas-fir plantations in a range of European countries

Initial density (plants ha ⁻¹)	Initial spacing (m)	Country	Observations	Source
600–800	4×4, 3×4	Belgium		Wouters and Lorent (2002)
600–800 (1000)	5×2.5, 4×4, 4×2.5	France	Adapted to small estates or dynamic forest owners	Riou-Nivert (2020)
1000–1200		France	Dynamic silviculture, with only one intervention before mean height of 12 m	Riou-Nivert (2020)
1000–1500		Italy	Only one thinning until rotation age	La Porta pers.comm
1000–2000		1. Europe-wide	1. Offer optimal balance between quality, stability, total volume growth and diameter growth rate	1. Kohnle et al. (2019)
		2. Germany	2. To avoid growth rings wider than 8 mm	2. Kohnle (2008)
		3. Germany		3. Klädtke et al. (2012)
1100–2500	2×2, 3×3	Serbia		Jovanović and Stojanović (1982)
1200–1500		France	First intervention when mean height reaches 16 m	Riou-Nivert (2020)
1500–2000		Austria	Sawtimber and veneer production	Schönauer (2008)
Max. 1600		Germany		Knook and Hanewinkel (2019)
1700–2500	2×2, 2.5×2, 3×2	Hungary		Rédei pers.comm
1700–3500	2.0–3.0×1.5–2.0	Bulgaria		Petkova pers.comm
2000–2500		Belgium		Perin et al. (2014)
		Italy	Two thinning until rotation age	La Porta pers.comm
2500	2×2	1. Romania	Fill the gaps of natural regenerations	1. MAPP (2000a)
		2. Ireland		2. Horgan et al. (2003)
		3. United Kingdom		3. Mason pers.comm
3000		Czech Republic		Podrázský et al. (2019)
4400	1.5×1.5	Romania	Plantations 50% Douglas-fir 50% broadleaves	MAPP (2000a)

Young stand management

In many countries where the initial stand density is often no more than 2000 plants ha⁻¹, management of young stands does not generally involve cleaning or respacing. However, these measures become necessary when pioneer, light-demanding and fast-growing species such as silver birch, trembling aspen (*Populus tremula* L.), and goat willow (*Salix capraea* L.) become naturally established and compete with young Douglas-fir trees (Liubimirescu 1973). In very dense natural regenerations of European beech (together with hornbeam *Carpinus betulus* L., sessile oak *Quercus petraea* (Matt.) Liebl., small-leaved lime *Tilia cordata* Mill.) where Douglas-fir is planted to fill-in gaps, there is a need to protect and promote the conifer as it grows slowly in the first 4–5 years and can be suppressed by native species (Liubimirescu 1973; Marcu and Liubimirescu 1979; Eberhard and Hasenauer 2018; Kohnle et al. 2019). Consequently, woody weeds should be removed from around Douglas-fir seedlings in the first 2–3 years (Rose and Rosner 2005), especially on rich soil, followed by the first cleaning when height is about 2 m and ca. 2000 trees ha⁻¹ remain. This intervention is

followed by a respacing when the average height is 5–7 m, and ca. 1000 trees ha⁻¹ remain (Podrázský et al. 2019). In mixed stands, Douglas-fir should be at least 20%–30% (Podrázský et al. 2019) of the remaining trees with undesirable species such as hornbeam or poorly formed Douglas-fir trees removed (Liubimirescu 1973).

In denser stands, such as in the Czech Republic and Romania, where the species is mixed with Norway spruce or European beech, the first intervention is carried out when the average height is 4–5 m and the density has been reduced to 2000 plants ha⁻¹, of which Douglas-fir is 20%–30%. The second respacing follows when average height reaches 10 m and the density is reduced to ca. 1500 trees ha⁻¹ (Podrázský et al. 2019). If a necessary cleaning or respacing is delayed, the trees can be slender and prone to toppling after release due to poorly developed crowns and roots (Eberhard et al. 2021).

Thinning and pruning

Thinning should start early, ideally before 20 years of age (de Champs and Demarq 1996; Riou-Nivert 1996, 2020;

Kohnle 2008). Delayed early thinning can result in stand instability and make Douglas-fir susceptible to windthrow (Wilson and Baker 2001). The first thinning should be heavy, removing 20%–25% of trees (Romania: Liubimirescu 1973) up to 30%–35% (France: CRPF 1999) to fully utilize the growth potential of the species. During subsequent stand development, 3–5 thinnings may be carried out depending on the rotation length (CRPF 1999); the thinning cycle varies: from 5 years (Belgium: Wouters and Lorent 2002; Romania: Liubimirescu 1973), up to 10 years (Serbia: Andrašev pers.comm.). The volume removed per thinning may be as high as 60 m³ ha⁻¹ or even higher in dense stands (Germany: Vor pers.comm.).

A range of thinning regimes have been proposed for Douglas-fir stands but most involve an initial line thinning to allow machine access (CRPF 1999; Horgan et al. 2003) followed by a mixture of selective thinning types. While thinning from below or intermediate (both from above and from below) was frequently practiced in the past (e.g., Czech Republic: Podrázský pers.comm; Romania: Liubimirescu 1973), recent decades have seen increased use of thinning from above in several countries, including Germany (Weise et al. 2001; Kohnle 2008; Kohnle et al. 2019). In some countries, before the second thinning when the mean diameter is about 15 cm, the (potential) final crop trees to be favoured during subsequent operations are selected and permanently marked. The criteria for selection are vigour, stem quality and spacing (Klädtker and Abetz 2010, in Kohnle et al. 2019). The final number of crop trees will vary with the intensity of silviculture being used, thus in Belgium it is around 150–200 trees ha⁻¹ in traditional silviculture, increasing to 250–300 trees ha⁻¹ with more intensive management (Wouters and Lorent 2002). It will also decrease as the target diameter increases (Kohnle et al. 2019).

In the last two decades, several regimes for even-aged Douglas-fir silviculture have been proposed which vary according to the initial stand density, the frequency of thinning, the incidence of pruning, and the target wood products. For example, in France, a range of options has been devised for private Douglas-fir forests (Riou-Nivert 2020). Thus, when an initial low density (600–800 plants ha⁻¹) is used, the objective is to produce large diameter stems (200 trees ha⁻¹ over 50 cm dbh) in 50–60 years with the wood destined for carpentry and timber frames. This regime involves 3–4 thinnings, starting when the mean height is 16–18 m and includes pruning up to 6 m. At the other extreme is the high-density option with an initial density of 1500–1600 trees ha⁻¹ aiming for 40–45 cm dbh trees with fine branches and narrow growth rings in 40–45 years. This includes only 1–2 thinnings and no pruning with the first thinning taking place when mean height is ca. 20 m.

Branching and consequent knot abundance and size are very important features affecting the grading of roundwood,

veneer and timber (Drewett 2015; Henin et al. 2019). There is consensus that self pruning of Douglas-fir is slow, even in dense stands (Fig. 2) (e.g., Hermann and Lavender 1990; Hubert and Courraud 1998; Lowell et al. 2014; Drewett 2015; Savill 2019), so unpruned Douglas-fir can be very knotty (Haralamb 1967; Savill 2019; Smolnikar et al. 2021).

Therefore, pruning is widely regarded as essential to increase log quality when targeting the production of highly valuable wood products, e.g., France (Hubert and Courraud 1998), Germany (Kohnle et al. 2019, 2021), the Czech Republic (Novák et al. 2018; Mondek and Balaš 2019) and Switzerland (Schütz et al. 2015). The selling price for well-pruned logs could easily be 80 euros more per m³ than for unpruned logs (Schütz et al. 2015). If the number of branches is affected by provenance choice (Fletcher and Samuel 2010), the diameter of branches depends on stand density (Drewett 2015; Smolnikar et al. 2021). As noted by Hein et al. (2008), even at densities exceeding 1200 trees ha⁻¹, artificial pruning may be necessary to produce high quality logs.

The main characteristics of successful pruning operations include only pruning the final crop trees where the production of high quality wood assortments is the goal (Marcu and Liubimirescu 1979). Depending on the country and



Fig. 2 Slow natural pruning of 50-year-old Douglas-fir in Slovenia (photo R. Brus)

silvicultural regime used, the final number of crop trees to prune per hectare is variable: i.e., in France from 100 trees (pruning height up to 8–10 m) to 200–400 trees (up to 6 m) (Hubert and Courraud 1998; CNPF 2016), in the Czech Republic 200 trees (Podrázský et al. 2019) or 200–360 trees in Romania (Liubimirescu 1973; Liubimirescu et al. 1977). Usually, the final pruning height is between 5 and 10 m (Eberhard et al. 2021) with two interventions (Horgan et al. 2003; Riou-Nivert 2020). In practice, pruning height is strongly correlated with the target diameter: 5 m (diameter 60 cm), 10 m (diameter 80 cm) and (seldom) 15 m (diameter 100 cm) (Kohnle 2008). The diameter of trees chosen for pruning should be between 10 and 15 cm as it is too early to select final crop trees < 10 cm, and no more than 15 cm because there will not be enough clearwood produced to increase the log value when the tree is harvested. In other words, pruning occurs before the trees reach one third of their final diameter (Nițescu and Achimescu 1979; Siepmann 1981; Schütz 1990; CRPF 1999).

The most frequently proposed pruning season is the end of winter-beginning of spring (Querengässer 1956, from IDT 1961; Haralamb 1967; Marcu and Liubimirescu 1979) to allow for the healing of pruning wounds. Pruning green branches up to 25%–30% of live crown in one operation and retaining a minimum green crown of 50% leads to only negligible losses in volume growth; the wounds heal quickly and the taper is improved (Lanier 1986; Boudru 1989; CRPF 1999; MAPPM 2000b). The branches to remove should be ≤ 3 cm in diameter (Wouters and Lorent 2002).

Rotation length

Douglas-fir stands are managed with rotation ages between 40 and 80 years (Table 5), although longer rotations are found in parts of Eastern Europe. In its native range in Canada, the rotation age for pure coastal Douglas-fir stands was 50–80 years four decades ago (Scott 1981; in Hermann and Lavender 1990), and currently 40–60 years in industrial forests in the U.S.A. (Lowell et al. 2014). In New Zealand, the rotation age of Douglas-fir plantations is 40–50 years (without thinning—www.douglasfir.co.nz/net/) but may be 60 years when the majority of trees are harvested for lumber (www.nzffa.org.nz/farm-forestry-model/).

Large pruned trees with diameters > 60 cm are valued for the production of solid furniture, decorative panels, peeled veneer as well as for exterior uses (CRPF 2010). Shorter rotations (less than 50 years) are generally not favoured as smaller logs have high proportions of juvenile wood and less favorable mechanical properties than mature wood (Barett and Kellog 1991; Lausberg et al. 1995; Vikram et al. 2011; Drewett 2015; all cited by Pollet et al. 2017). Long rotations in combination with large diameters are favorable management options to reduce the amount of juvenile wood and

provide a more homogenous wood material (Henin et al. 2019). Slower radial growth usually benefits technological wood properties (e.g., reduced proportion of juvenile wood, branch diameters) (Henin et al. 2019), but long rotation are associated with higher risks, i.e., storm damage increases progressively with stand height (CRPF 2010; Kohnle et al. 2021). When reaching rotation age, European stands of Douglas-fir have traditionally been harvested through clear-felling, followed by replanting on the site as in the Czech Republic (Podrázský pers.comm.), Croatia (Đodan pers.com.), Hungary (Rédei pers.comm.), Ukraine (Lavnyy pers.comm.), UK (Mason pers.comm.). However, there has been increasing interest in adopting alternative silvicultural approaches characteristic of closer-to-nature forestry (Larsen et al. 2022).

Mixtures

In the context of a wider use of biodiversity-friendly practices such as closer-to-nature-forestry as part of the EU Biodiversity Strategy for 2030 (European Commission 2020), the use of Douglas-fir in mixed stands with native tree species is of major importance (Kohnle et al. 2021). This mixture with greater biodiversity than pure stands can (1) increase resistance and resilience to biotic agents detrimental to stand health, and (2) link stands of the exotic Douglas-fir to the native forest community, an aspect of particular importance within the concept of close-to-nature forest management (Kohnle et al. 2019).

The patterns of planting Douglas-fir to create mixed stands vary across Europe. These include establishing groups in a matrix of other species as in the Czech Republic (Novák et al. 2018; Slodičák et al. 2014; both in Podrázský et al. 2019) and in Romania, when the species is used in filling gaps in natural broadleaf regeneration (MAPPM 2000a). Group planting to form mixtures is also advocated as means of avoiding any negative environmental impacts from Douglas-fir (Ammer and Utschick 2004; in Thomas et al. 2022). Planting in rows can also be used to create mixtures but there should be a maximum of 4 m between rows to avoid the development of thick branches (Kohnle 2008). Current Romanian technical norms also recommend planting Douglas-fir in rows 1.5 m apart when the species is used in mixture with European beech (MAPPM 2000a).

Douglas-fir has been grown in mixtures with a wide range of mostly native conifers and broadleaf species. These include Norway spruce (Bulgaria, the Czech Republic, Germany, Slovenia), Sitka spruce (Ireland, UK), European silver fir (the Czech Republic, France, Slovenia), Scots pine (Bulgaria, Germany), black pine (*Pinus nigra* J.F. Arnold) (Bulgaria), European larch (Bulgaria, UK), hybrid larch (*Larix x eurolepis*) (Ireland), Japanese larch (*Larix kaempferi* (Lamb.) Carr) (Ireland, UK), western hemlock (*Tsuga*

Table 5 Stand rotation ages and target diameters of Douglas-fir in different European countries

Country	Rotation age, years	Target diameter, cm	Source
France	1. 40 (in monocultures)	3. 40 (mean diameter)	1. van Loo and Dobrowolska (2019b)
	2. Minimum 40, maximum 60–80		2. Bastien (1998)
	3. 40–50 (initial planting density 1000–1200, 1200–1500 or 1500–1600 seedlings ha ⁻¹)		3. Riou-Nivert (2020)
	4. 40–80	5. Minimum 50 (mean diameter)	4. Riou-Nivert (1996)
	5. 50–60 (initial planting density 600–800 seedlings ha ⁻¹)		5. Riou-Nivert (2020)
	6. 50–70		6. CRPF (1999)
	7. 65–75		7. Giraud and Champaux (1997)
	8. 66–74		8. Angelier (2007)
		9. From 40–50 cm on, even 55–60 cm in stands of good quality	9. CRPF (2010)
United Kingdom	1. Ca. 50; except where a premium is paid for larger logs produced on longer rotation		1. Mason pers.comm
	2. 50–65 (absolute rotation age)		2. Savill (2019)
Romania	50 (absolute rotation age)		Bakoş (1968)
	75–80 (veneer, sawnwood)		Liubimirescu (1973)
Portugal	50–60		Fontes et al. (2003)
Ireland	50–65		Horgan et al. (2003)
Belgium	50–80		Wouters and Lorent (2002)
	70–80		Boudru (1989)
Croatia	60		Đodan pers.comm
Germany (Baden-Württemberg)	1. 60+ (in mixed stands)	2. Target diameter 50+ 3. Target diameter 70+ 4. 70–80 (when mean age is 80–90 years)	1. van Loo and Dobrowolska (2019b)
	2. 60–80		2. Spellmann et al. (2015)
	3. 80–120		3. Spellmann et al. (2015)
			4. Şimon et al. (2014)
Bosnia and Herzegovina	60–120, depending on site quality		Cvjetkovic pers.comm
Czech Republic	80+		Podrázský pers.comm
Serbia	80–100		Andrašev pers.comm
Ukraine	81–90		Lavnyy pers.comm
Bulgaria	100–120		Petkova pers.comm

heterophylla (Raf.) Sarg.) (Ireland, UK), European beech (Czech Republic, France, Germany, Romania, Serbia, Switzerland, Ireland), Italian alder (*Alnus cordata* Loisel.) (Italy) (Čokl 1965; Petkova 1989; Alexandrov et al. 2000; MAPP 2000a; Wilson and Cameron 2015; Petkova et al. 2017; Keane et al. 2018; COFORD 2020; Nicolescu et al. 2021; Royal Forestry Society 2021; Andrašev pers.comm.; Đodan pers.comm.; La Porta pers.comm.; Podrázský pers.comm.; Vor pers.comm.). In the Czech Republic, the share of the species in mixtures is about 20%–40%, evenly distributed over the area (Podrázský et al. 2019). A similar share of Douglas-fir (30% on average) was used in mixed plantations in Austria and Germany (Eberhard et al. 2021). In both Denmark (Larsen and Nielsen 2007) and in the UK (Haufe et al. 2021), forest development guidelines have been

proposed to help the development of mixtures of Douglas-fir with broadleaves and conifers.

European beech is considered particularly suitable for mixing with Douglas-fir (MAPP 2000a; Thomas et al. 2015; Ammer et al. 2016). In such mixtures, Douglas-fir is stabilized against climatic impacts while the climate sensitivity of European beech is increased (Thurm et al. 2016). Douglas-fir growth recovery after drought was shortened and extended for European beech (Thurm et al. 2016). In addition, the productivity of Douglas-fir-European beech stands regularly showed higher productivity compared with that predicted from the species performance in pure stands, largely attributed to the Douglas-fir (Thurm et al. 2016; Lu et al. 2018), with a share up to 33% (Thomas et al. 2015). Another species that may be used in a mixture with

Douglas-fir is Norway spruce. The proportion of Douglas-fir should be only 30% in such mixed stands, a compromise which might offer environmental and social adaptability and help to maintain the productivity of current Norway spruce sites (Fuchs et al. 2022). Douglas-fir may also be used as a nurse species since experiments in Ireland have shown that when it is grown in mixture with Sitka spruce on marginal sites, the latter species has higher productivity than if grown in pure stands (Keane et al. 2018). In its natural range, mixtures of Douglas-fir and red alder (*Alnus rubra* Bong.) resulted in long-term productivity gains for seven decades on nitrogen deficient soils (Binkley 2003).

Irregular silviculture

As Douglas-fir's ecological characteristics are considered 'well adapted to irregular silviculture' (Royal Forestry Society 2021), alternative systems to clear-cutting are being carried out in different European countries. These include: group shelterwood cuttings (Bulgaria: Petkova pers.comm.; France: Riou-Nivert 2020; Slovakia: Slávik pers.comm.); single-tree selection cutting (Bosnia and Herzegovina: Cvjetkovic pers.comm.; Germany: Vor pers.comm.; Slovakia: Slávik pers.comm.), group selection cutting (UK: Schütz and Pommerening 2013); and irregular shelterwood cuttings (France: Bastien 1998; Riou-Nivert 2020; UK: Malcolm et al. 2001; Mason et al. 2004). With irregular shelterwood cuttings, the recommended gap size for satisfactory natural regeneration and seedling growth is at least 0.1 ha, with a gap diameter: top height ratio of 1.5–2.0 (Malcolm et al. 2001).

As noted above, Douglas-fir can be regenerated naturally under the shelter of old, even-aged stands using silvicultural systems such as group shelterwood cutting; if the residual stand is removed (5 to 15 years between the seedling cut and the final cut as proposed in France, Riou-Nivert 2020), the newly established regeneration grows without any competition from above, and subsequent growth dynamics are expected to be similar to the regeneration process in its native range (Kohnle et al. 2019). However, insufficient opening of the canopy may result in poor rooting and crown development (Eberhard et al. 2021). The species also regenerates under canopy shelter with lengthy regeneration periods (over 40 years in Bulgaria—Petkova pers.comm.), more typical of shade-tolerant species (Kohnle et al. 2019). This is the pattern in uneven-aged stands using single-tree selection cutting, group-selection cutting or irregular high forest cutting harvesting individual trees reaching the target diameter. The application of single-tree selection cutting is possible only if the basal area is relatively low (about 27 m² ha⁻¹) (Schütz et al. 2015). Under these conditions, young trees, tolerant of shade, are protected against late frost and temperature extremes (Slávik pers.comm.; Vor pers.comm.).

In its natural range in North America, there is increasing adoption of 'ecological silviculture' in public and some private forests, an approach which attempts to follow the natural disturbance regime in Douglas-fir and western hemlock forests to ensure provision of a wide range of environmental services (Palik et al. 2021). Particular features of this approach include the allowance for an extended establishment period (up to 30 years), variable density thinning to develop stand heterogeneity, and harvesting regimes which ensure retention of deadwood and other structural legacies (Larsen et al. 2022). Several of these have yet to be explored in European silvicultural practices with Douglas-fir, especially the use of variable density thinning to create spatial variations within stands which positively affects the restoration of desired ecosystem features after periods of even-aged management (Puettmann et al. 2016).

Conclusion

This paper has outlined the qualities that have led to the use of Douglas-fir, considered as 'one of the most successful species introduced in the history of European forestry' (Podrázský et al. 2013), on a relatively large scale in Europe. It is a very productive species, with valuable timber for various important end-uses. It can form both pure and mixed stands, including mixtures with various native species such as European beech.

In addition, the species generally affects the environment less than Norway spruce or numerous pine species. It has a lower negative impact on the biodiversity of soil flora and has a more favorable role on soil formation especially when mixed with broadleaved species such as European beech (Thomas et al. 2022).

The important position of this species in European forestry is strengthened in the context of anticipated climate change, which favors a species better adapted to drought than Norway spruce, the most important conifer species in Europe. Nevertheless, climate change as well as globalization may have implications for the management of Douglas-fir, for example if biotic pests from its native range are introduced and become adapted to Douglas-fir in Europe. Under anticipated climate change, a major problem will be to identify provenances best adapted to future climatic conditions (Eilmann et al. 2013; Isaac-Renton et al. 2014, Hintsteiner et al. 2018; Konnert et al. 2018; Spiecker 2019; all in Smolnikar et al. 2021). This could require the use of new seed sources and provenances from its natural range, adapted to drier conditions such as those from southern Oregon and northern California (Konnert and Bastien 2019; Marchi and Coccozza 2021; Schüler and Chakraborty 2021).

However, public perception of Douglas-fir in European countries varies appreciably from widespread acceptance in western countries to negative views (even restrictions on its use) in eastern countries. In this context, we echo the very realistic conclusion by Spiecker (2019) and Spiecker and Schuler (2019), ‘... *the reputation of Douglas-fir and the question of the continued growth of this species in Europe is loaded with hope, prejudice, reservation, and scepticism. The current debates among numerous stakeholders often vary from enthusiastic to emotional, and can benefit from an evidence-based, sound scientific knowledge*’. We hope that this paper can help to inform such debates.

Acknowledgements This paper is an output from the European COST Action FP1403 ‘Non-native tree species for European forests-experiences, risks, and opportunities’ (NNEXT) active during 2014–2018. The authors also acknowledge helpful suggestions from the reviewers and the editor.

Declarations

Conflict of interest The authors have no competing interests to declare that are relevant to the content of this article.

Ethical approval The authors declare they followed the rules of good scientific practice when preparing this article.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Albrecht A, Kohnle U, Hanewinkel M, Bauhus J (2010) Storm damage of Douglas-fir and norway spruce in southwest Germany: stability of Douglas-fir and the impact of silviculture on the vulnerability of conifers. In: Spiecker H, Kohnle U, Makkonen-Spiecker K, von Teuffel K (eds) Opportunities and risks for douglas fir in a changing climate. Freiburger Forstliche Forschung, Freiburg, pp 25–27
- Albrecht A, Kohnle U, Hanewinkel M, Bauhus J (2013) Storm damage to Douglas-fir unexpectedly high compared to Norway spruce. *Ann Forest Sci* 70(2):195–207
- Alden HA (1997) Softwoods of North America. General technical report FPL-GTR-102, United States department of agriculture. Forest service, Forests products laboratory, Madison. p 151
- Aldhous JR, Mason WL (eds) (1994) Forestry nursery practice. Forestry commission bulletin 111. HMSO, London, p 268
- Alexandrov A, Genov K, Popov E (2000) On the introduction of some North-American species. In: Book of scientific reports, international scientific conference “75th anniversary of forestry education in Bulgaria”, Sofia, vol. I., pp 384–390, 15–16 June 2000. (in Bulgarian)
- Ammer C, Bolte A, Herberg A, Höltermann A, Krüß A, Krug A, Nehring S, Schmidt O, Spellmann H, Vor T (2016) Recommendations for the cultivation of introduced forest tree species. *Gem Papier Forstwiss Naturschutz* 48(5):170–171 ((in German))
- Angelier A (2007) French Douglas-fir forests. Office National des Forêts, Paris, p 296 (in French)
- Anonymous (1999) Silvicultural guidelines in relation to stand development types. Ministerium Ländlicher Raum, Landesforstverwaltung Baden-Württemberg, Freiburg, pp 11–12 (in German)
- Bakoş V (1968) Introduction. General issues on afforestations in our country. In: Bakoş V (coord): culture of fast-growing forest species. Editura Agrosilvică, Bucureşti, pp 7–56 (in Romanian)
- Bastien J-C (2019) Potential of Douglas-fir under climate change. In: Spiecker H, Lindner M, Schuler J (eds) Douglas-fir—an option for Europe. European Forest Institute, Joensuu, pp 40–45
- Bastien J-C, Sanchez L, Michaud D (2013) Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco). In: Pâques LE (ed) Forest tree breeding in Europe. Current state-of-the-art and perspectives. Springer, Dordrecht Heidelberg-New York-London, pp 325–372
- Bastien Y (1998) Douglas-fir silviculture. In: Silviculture of conifer species, Ecole Nationale du Génie Rural, des Eaux et des Forêts, Nancy, pp 10–19 (in French)
- Bindewald A, Miocic S, Wedleer A, Bauhus J (2021) Forest inventory-based assessment of the invasion risk of *Pseudotsuga menziesii* (Mirb.) Franco and *Quercus rubra* L. in Germany. *Eur J Forest Res* 140:883–899. <https://doi.org/10.1007/s10342-021-01373-0>
- Binkley D (2003) Seven decades of stand development in mixed and pure stands of conifers and nitrogen-fixing red alder. *Can J Forest Res* 33:2274–2279
- Bluskova D (2006) Studies on the structure and properties of juvenile timber of the main tree species. In: Habilitation thesis for the academic title of “Professor”. University of Forestry, Sofia, p 107 (in Bulgarian)
- BMEL (2014) Forests in Germany—data on the 3rd National forest inventory. Bundesministerium für Ernährung und Landwirtschaft, Berlin (in German)
- Boudru M (1989) Forest and silviculture: applied silviculture. Les presses agronomiques de Gembloux, Gembloux, p 248 (in French)
- Boudru M (1992) Forest and silviculture: afforestation and reforestation. Les presses agronomiques de Gembloux, Gembloux, p 348 (in French)
- Boussaid O (2008) Genetic quality of a natural regeneration of Douglas-fir, *Pseudotsuga menziesii* (Mirb.) Franco. Master’s thesis. Orléans University, Orléans, p 42 (in French)
- Brus R, Pötzelsberger E, Lapin K, Brundu G, Orazio C, Straigyte L, Hasenauer H (2019) Extent, distribution and origin of non-native tree species in Europe. *Scand J Forest Res* 34(7):533–544
- Carter RE, Klinka K (1992) Variation in shade tolerance of Douglas-fir, western hemlock, and western red cedar in coastal British Columbia. *For Ecol Manag* 55:87–105. [https://doi.org/10.1016/0378-1127\(92\)90094-P](https://doi.org/10.1016/0378-1127(92)90094-P)
- Čáter M (2021) Microsites influence the light response of young Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco). *Forests* 12:687. <https://doi.org/10.3390/f12060687>
- Chakraborty D, Wang T, Andre K, Konnerth M, Lexer MJ, Matulla C, Schueler S (2015) Selecting populations for non-analogous climate conditions using universal response functions: The case of Douglas-fir in central Europe. *PLoS ONE* 10(8):e0136357. <https://doi.org/10.1371/journal.pone.0136357>

- Chakraborty D, Matulla C, Andre K, Weissenbacher L, Schueler S (2019) Survival of Douglas-fir provenances in Austria: site-specific late and early frost events are more important than provenance origin. *Ann for Sci* 76(4):100. <https://doi.org/10.1007/s13595-019-0083-2>
- CNPF (2016) Pruning Douglas-fir: take the height. Centre National de la Propriété Forestière Bourgogne Franche Comté, Dijon, p 6 (in French)
- COFORD (2020) Sustainable development and conservation of forest genetic resources 2020–2030. COFORD, Dublin, p 60
- Čokl M (1965) Growth of green Douglas fir in Slovenia. *Zbornik* 4:139–185 (in Slovenian)
- Coopérative forestière bois Limousin (2016) Douglas-fir, a chance for France and for foresters. Ed. CFBL, Ussel Cedex, p 136 (in French)
- Covre Foltran E, Ammer C, Lamersdorf N (2020) Douglas fir and Norway spruce admixtures in beech forests along in Northern Germany—are soil nutrient conditions affected? *bioRxiv the Preprint Server for Biology*. <https://doi.org/10.1101/2020.09.25.313213>
- CRPF (1999) Douglas-fir. Centre Régional de la Propriété Forestière d'Ile-de-France et du Centre, Orléans, p 4 (in French)
- CRPF (2010) When harvesting Douglas-fir? Centre Régional de la Propriété Forestière de Bourgogne, Dijon, p 6 (in French)
- Curtis RO, DeBell DS, Miller RE, Newton M, St. Clair JB, Stein WI (2007) Silvicultural research and the evolution of forest practices in the Douglas-fir region. In: General Technical Report PNW-GTR-696, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, p 172
- da Ronch F, Caudullo G, de Rigo D (2016) *Pseudotsuga menziesii* in Europe: distribution, habitat, usage and threats. In: San-Miguel-Ayán J, de Rigo D, Caudullo G, Houston Durant T, Mauri A (eds) European atlas of forest tree species. Publication Office of the EU, Luxembourg, pp 146–147
- de Champs J, Demarq P (1996) Douglas-fir silviculture. Bulletin technique no. 31, Office National des Forêts, octobre: 35–42 (in French)
- de Champs J (1988) What silviculture for current Douglas-fir? *Afocel-Armef, Inf-Forêt* 1:65–80 (in French)
- Decker L (2018) Windthrow risk assessment of Douglas-fir stands in Denmark. In: A comparison study of Douglas-fir (*Pseudotsuga menziesii*) and Norway spruce (*Picea abies*). Master's thesis. University of Copenhagen, Faculty of Sciences, Copenhagen
- Dobrev D (1962) Afforestation with foreign fast-growing coniferous tree species. Sofia, Zemizdat, p 220 (in Bulgarian)
- Drăcea M (1923) Exotic species and their naturalization in our country. *Revista Pădurilor* 3:197–212 (in Romanian)
- Drewett TA (2015) The growth and quality of UK-grown Douglas-fir. In: Ph.D. Thesis. Edinburgh Napier University, Edinburgh, p 224
- Dumitriu-Tătăranu I (1960) Forest and ornamental trees and shrubs cultivated in the People's Republic of Romania. Editura Agro-Silvică, București, p 810
- Dungey HS, Low CB, Watt MS, Hood IA, Stone JK, Kimberley M (2010) Planted forests of Douglas-fir in New Zealand: challenges for the breeding programme with climate change. In: Spiecker H, Kohnle U, Makkonen-Spiecker K, von Teuffel K (eds) Opportunities and risks for Douglas fir in a changing climate. Freiburger Forstliche Forschung, Freiburg, p 48
- Eberhard B, Hasenauer H (2018) Modeling regeneration of Douglas fir forests in central Europe. *Austrian J for Sci/ Centralblatt Für Das gesamteForstwesen* 135(1):33–51
- Eberhard BR, Eckhart T, Hasenauer H (2021) Evaluating strategies for the management of Douglas-fir in central Europe. *Forests* 12(8):1040. <https://doi.org/10.3390/f12081040>
- Eckhart T, Pötzelsberger E, Koeck R, Thom D, Lair GJ, van Loo M, Hasenauer H (2019) Forest stand productivity derived from site conditions: an assessment of old Douglas-fir stands (*Pseudotsuga menziesii* (Mirb.) Franco var. *menziesii*) in Central Europe. *Ann Forest Sci* 76:19. <https://doi.org/10.1007/s13595-019-0805-3>
- Eilmann B, Rigling A (2010) Douglas-fir—a substitute species for Scots pine in dry inner-Alpine valleys? In: Spiecker H, Kohnle U, Makkonen-Spiecker K, von Teuffel K (eds) Opportunities and risks for Douglas fir in a changing climate. Freiburger Forstliche Forschung, Freiburg, p 10
- Eilmann B, Rigling A (2012) Tree-growth analyses to estimate tree species' drought tolerance. *Tree Physiol* 32(2):178–187
- Essl F (2005) Distribution, status and habitat preference of spontaneous Douglas fir stands (*Pseudotsuga menziesii*) in Austria. *Phyton (horn)* 45(1):117–143
- European Commission (2020) EU Biodiversity Strategy for 2030. Bringing nature back into our lives. European Commission, Brussels, p 22
- Forest Europe (2020) State of Europe's forests. In: Ministerial Conference on the Protection of Forests in Europe. Forest Europe, Zvolen, Liaison Unit Bratislava, p 392
- Fletcher AM, Samuel CJA (2010) Choice of Douglas fir seed sources for use in British forests. *Forestry Commission Bulletin* 129, Edinburgh, p 55
- Fontes L, Tomé M, Coelho MB, Wright H, Sales Luis H, Savill P (2003) Modelling dominant height growth of Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) in Portugal. *Forestry* 76(5):509–523
- France Bois Forêt (2016) Usage classes and longevity of wooden structures. http://franceboisforet.com/wp-content/uploads/2014/06/fiches-C4-LCELOB_4-pages.pdf. [Accessed 10 March 2022] (in French)
- Frei RR, Moser B, Wohlgenuth T (2022) Competitive ability of natural Douglas fir regeneration in central European close-to-nature forests. *For Ecol Manag* 503:119767. <https://doi.org/10.1016/j.foreco.2021.119767>
- Fuchs JM, Hittenbeck A, Brandl S, Schmidt M, Paul C (2022) Adaptation strategies for spruce forests - economic potential of bark beetle management and Douglas fir cultivation in future tree species portfolios. *For: Int J for Res* 95(2):229–246
- Georgieva M (2009) Diseases on Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) and their impact on the introduction of the species in Bulgaria. In: Dissertation. Forest Research Institute, Bulgarian Academy of Sciences, Sofia (in Bulgarian)
- Giagli K, Timko L, Gryc V, Vavrčik H (2019) Is the quality of the non-native Douglas-fir wood produced in the czech forests comparable to native softwoods? *Bio Resources* 14(2):2931–2945
- Gil-Moreno D, Ridley-Ellis D, Harte AM (2019) Timber grading potential of Douglas fir in the Republic of Ireland and the UK. *Int Wood Prod J* 10(2):1–6. <https://doi.org/10.1080/20426445.2019.1617984>
- Giraud P, de Champeaux A (1997) Douglas-fir in Haut Beaujolais, Charolais and Morvan. Experience of some forest owners. *Revue Forestière Française* 49(4):325–338 (in French)
- Govedar Z, Ballian D, Mikić T, Pintarić K (2003) Success of different provenances of Douglas fir (*Pseudotsuga menziesii* Mirb. Franco.) within the IUFRO program on the experimental plot CrnaLokva“ near Gradiška. *Forestry (belgrade)* 3–4:61–74 (in Serbian)
- Gozd in gozdarstvo (2021) Forest and forestry. <https://www.gozd-les.com/les/cene-lesa/dosezene-cene-licitaciji-lesa> [Accessed June 10, 2021]
- Hale SE, Gardiner BA, Peace A, Nicoll B, Taylor P, Pizzirani S (2015) Comparison and validation of three versions of a forest wind risk model. *Environ Model Softw* 68:27–41
- Haralamb A (1967) Culture of forest species, 3rd edn. Editura Agro-Silvică, București, p 755 (in Romanian)

- Hart C (1994) Practical forestry for the agent and surveyor, 3rd edn. Alan Sutton, Stroud, p 658
- Haufe J, Kerr G, Stokes V (2021) Forest Development Types: an essential tool for diversifying stands in Britain. *Q J for* 115(3):175–182
- Hein S, Weiskittel AR, Kohnle U (2008) Effect of wide spacing on tree growth, branch and sapwood properties of young Douglas-fir [*Pseudotsuga menziesii*(Mirb.) Franco] in south-western Germany. *Eur J Forest Res* 127(6):481–493
- Henin JM, Pollet C, Schmitt U, Blohm J-H, Koch G, Melcher E, Wellington J, Brüchert F, Kohnle U, Sauter UH (2019) Technological properties of Douglas-fir wood. In: Spiecker H, Lindner M, Schuler J (eds) Douglas-fir - an option for Europe. European Forest Institute, Joensuu, pp 89–97
- Hermann RK, Lavender DP (1990) *Pseudotsuga menziesii* (Mirb.) Franco Douglas fir. In: Burns RM, Honkala BH (Coord.) Silvics of North America. Volume 1, Conifers. USDA Forest Service, Washington, pp 527–540
- Hermann RK, Lavender DP (1999) Douglas-fir planted forests. In: Boyle JR, Winjum JK, Kavanagh K, Jensen EC (eds) Planted forests: contributions to the quest for sustainable societies. Kluwer Academic Publishers, Dordrecht, pp 53–70
- Horgan T, Keane M, McCarthy R, Lally M, Thompson D (2003) A Guide to Forest Tree species selection and silviculture in Ireland. In: O'Carroll J (Eds). COFORD, Dublin, p 258
- Hostnik R (2021) The Giants. In: The Tallest Measured Tree in Slovenia. Available online: Velikanke/The Giants – YouTube. Accessed June 10, 2021
- Hubert M, Courraud R (1998) Pruning and formative pruning of forest trees. Institut pour le développement forestier, Paris, p 303 (in French)
- Ibrahimspahić A, Ballian D, Gurda S (2006) The analysis of flourishing of six conifer species in the area of Gostovičkarijeka. *Works Faculty for Univ Sarajevo* 36(1):57–67
- IDT (1961) Douglas-fir. New achievements in the culture of fast-growing species. Institutul de Documentare tehnică, București, pp 80–88 (in Romanian)
- Ionuț V (1956) A monograph of Douglas-fir in the People's Republic of Romania. Editura Agro-Silvică de Stat, București, p 50
- Ionescu AI (1963) Aspects on the culture of Douglas-fir and eastern white pine. *Revista Pădurilor* 78(8):434–437 (in Romanian)
- Isaac-Renton MG, Roberts DR, Hamann A, Spiecker H (2014) Douglas-fir plantations in Europe: a retrospective test of assisted migration to address climate change. *Global Change Biol* 20:2607–2617
- Jasser C (2008) Douglas-fir in upper Austria: possibilities and limits. *BFW-Praxisinformation* 16:19–20 (in German)
- Jones EW (1945) The regeneration of Douglas-fir *Pseudotsuga taxifolia* Britt. in the new forest. *J Ecol* 33:44–56
- Jovanović B, Stojanović LJ (1982) Development of cultures of some species of conifers in the extreme habitat of beech (*Muscofagetum*) in eastern Serbia. *Šumarstvo* 1:3–11
- Kahl T, Arnstadt T, Baber K, Bässler C, Bauhus J, Borcken W, Buscot F, Floren A, Heibl C, Hessenmöller D, Hofrichter M, Hoppe B, Kellner H, Krüger D, Linsenmair KE, Matzner E, Otto P, Purahong W, Seilwinder C, Schulze ED, Wende B, Weisser WW, Gossner MM (2017) Wood decay rates of 13 temperate tree species in relation to wood properties, enzyme activities and organismic diversities. *For Ecol Manag* 391:86–95
- Keane M, Mason B, Pfeifer A (2018) Species Mixtures in Irish Forests. In: A review. COFORD, department of agriculture food and the marine, Dublin, p 124
- Klädtke J, Kohnle U, Kublin E, Ehring A, Pretzsch H, Uhl E, Spellmann H, Weller A (2012) Growth and value production of Douglas-fir under varying stands densities. *Schweizerische Zeitschrift Für Forstwesen* 163(3):96–104. <https://doi.org/10.3188/szf.2012.0096>(in German)
- Klepac D (1962) A contribution to the knowledge of Douglas-fir and Eastern white pine increment. *Šumarski List* 86(1–2):10–31
- Knook J, Hanewinkel M (2019) Economics of growing Douglas-fir. In: Spiecker H, Lindner M, Schuler J (eds) Douglas-fir-an option for Europe. European Forest Institute, Joensuu, pp 99–104
- Kohnle U (2008) Growing Douglas-fir in South-West of Germany: factors of silvicultural success. *BFW-Praxisinformation* 16:12–13 (in German)
- Kohnle U, Klädtke J, Chopard B (2019) Management of Douglas-fir. In: Spiecker H, Lindner M, Schuler J (eds) Douglas-fir-an option for Europe. European Forest Institute, Joensuu, pp 73–83
- Kohnle U, Klädtke J, Chopard B (2021) Silviculture with Douglas fir in Germany and France. *Schweiz Z Forstwes* 172(2):66–75 (in German)
- Konnert M (2016) *Pseudotsuga Menziesii*. *Studia for Slovenica* 151:30–35
- Konnert M, Bastien JCh (2019) Genecology of Douglas-fir and tree improvement strategies. In: Spiecker H, Lindner M, Schuler J (eds) Douglas-fir - an option for Europe. European Forest Institute, Joensuu, pp 45–56
- La Marca O, Pozzi D (2016) From plantation silviculture to the one by natural renewal. In: Polistampa, Firenze (Eds) Realities and perspectives in the cultivation of Douglas-fir in Italy. *Attidella Accademia dei Georgofili* 8 (13): 43–64
- La Marca O, Mercurio R, Nocentini L (2016) Management and restoration of Italian afforestations under global change. *Reforesta* 2:50–59. <https://doi.org/10.21750/REFOR.2.06.21>
- Lange F, Ammer C, Leitinger G, Seliger A, Zerbe S (2022) Is Douglas fir [*Pseudotsuga menziesii* (Mirb.) Franco] invasive in central Europe? A case study from South-West Germany. *Front for Glob Change* 5:844580. <https://doi.org/10.3389/ffgc.2022.844580>
- Lanier L (1986) Handbook of Silviculture. ENGREF, Nancy, p 468 (in French)
- Larsen JB, Nielsen AB (2007) Nature-based Forest management - Where are we going? Elaborating forest development types in and with practice. *For Ecol Manag* 238:107–117
- Larsen JB, Angelstam P, Bauhus J, Carvalho JF, Diaci J, Dobrowolska D, Gazda A, Gustafsson L, Krumm F, Knoke T, Konczal A, Kuuluvainen T, Mason B, Motta R, Pötzelsberger E, Rigling A, Schuck A (2022) Closer-to-nature forest management. From science to policy 12. European Forest Institute, Joensuu, p 54. <https://doi.org/10.36333/fs12>
- Larson B (2010) The dynamics of Douglas-fir stands. In: Spiecker H, Kohnle U, Makkonen-Spiecker K, von Teuffel K (eds) Opportunities and risks for Douglas fir in a changing climate. Freiburger Forstliche Forschung, Freiburg, p IX–X
- Lavender DP, Hermann RK (2014) Douglas-fir: the genus *pseudotsuga*. Oregon forest research laboratory, Oregon state university, Corvallis, p 352
- Liese W, Moser E, Willeitner H (1982) Treatability of Douglas-fir sapwood from trees grown in Germany. *Holz Als Roh-Und Werkstoff* 40(9):321–325 (in German)
- Liubimirescu A (1973) On the need and application of tending operations in young Douglas-fir stands. *Revista Pădurilor* 88(4):209–211 (in Romanian)
- Liubimirescu A, Belea Gh, Jorda I (1977) Research on artificial pruning in young Douglas-fir stands. *Revista Pădurilor* 92(4):203–208 (in Romanian)
- Lowell EC, Maguire DA, Briggs DG, Turnblom EC, Jayawickrama KJS, Bryce J (2014) Effects of silviculture and genetics on branch/knot attributes of Coastal Pacific Northwest Douglas-fir and implications for wood quality—a synthesis. *Forests* 5:1717–1736

- Lu H, Condés S, del Rio M, Goudiaby V, den Ouden J, Mohren GMJ, Schelhaas MJ, de Waal R, Sterck FJ (2018) Species and soil effects on overyielding of tree species mixtures in the Netherlands. *For Ecol Manag* 409:105–118
- Macdonald J, Wood RF, Edwards MV, Aldhous JR (1957) Exotic forest trees in great Britain. In: *Forestry commission bulletin* 30, HMSO, London, p 167
- Malcolm DC, Mason WL, Clarke GC (2001) The transformation of conifer forests in Britain – regeneration, gap size and silvicultural systems. *For Ecol Manag* 157:7–23
- Malmqvist C, Wallertz K, Lindström A (2017) Storability and freezing tolerance of Douglas fir and Norway spruce seedlings grown in mid-Sweden. *Scand J Forest Res* 32(1):30–38. <https://doi.org/10.1080/02827581.2016.1183704>
- Malmqvist C (2017) Planting and survival of Douglas fir (*Pseudotsuga menziesii* (Mirb. Franco) in Sweden: Questions of seedling storability, site preparation, bud burst timing and freezing tolerance. Doctoral dissertation. Department of forestry and wood technology, Linnaeus University, Växjö, p 42
- Manise T, Vincke C (2014) Impacts of climate and site water deficits on the radial growth of beech, oak, spruce and Douglas-fir in Wallonia. *ForêtWallonne* 129:48–57 (in French)
- MAPP (2000a) Technical norms on species compositions, schemes and technologies for forest regeneration and afforestation of degraded lands 1. Ministerul Apelor, Pădurilor și Protecției Mediului, București, p 272 (in Romanian)
- MAPP (2000b) Technical norms for tending the stands 2. Ministerul Apelor, Pădurilor și Protecției Mediului, București, p 164 (in Romanian)
- Marchi M, Coccozza C (2021) Probabilistic provenance detection and management pathways for *Pseudotsugamenziesii* (Mirb.) Franco in Italy using climatic analogues. *Plants* 10(2):215. <https://doi.org/10.3390/plants10020215>
- Marcu Gh, Liubimirescu A (1979) Recommendations on zoning and culture of Douglas-fir in our country. *Revista Pădurilor* 94(2):74–80 (in Romanian)
- Mason WL, Edwards C, Hale SE (2004) Survival and early seedling growth of conifers with different shade tolerance in a Sitka spruce spacing trial and relationship to understorey light climate. *Silva Fennica* 38(4):357–370
- Mauer O, Palátová E (2012) Root system development in Douglas fir (*Pseudotsuga menziesii* [Mirb.] Franco) on fertile sites. *J for Sci* 58(9):400–409
- McKay HM, Milner AD (2000) Species and seasonal variability in the sensitivity of seedling conifer roots to drying and rough handling. *Forestry* 73:259–270
- Milenkova A (2020) Results of coniferous tree species afforestation in Osogovo, Southwest Bulgaria. Dissertation. University of Sofia, Sofia, p 162 (in Bulgarian)
- Molnár S (2008) Quality and utilization of wood of tree species managed by plantation – like growing technology. In: Führer E, Rédei K, Tóth B (eds) *Plantation-like timber production 2*. Agroinform Kiadó, Budapest, pp 153–190 (in Hungarian)
- Mondek J, Baláš M (2019) Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) and its role in the Czech forests. *J for Sci* 65(2):41–50
- Možina I (1960) Variations of gravity, summerwood, and strength of Douglas-fir (*Pseudotsuga taxifolia* Britton.). In: *Zbornik za kmetijstvo in gozdarstvo* 7, Ljubljana, p 136 (in Slovenian)
- Negulescu E, Săvulescu AI (1957) Dendrology. Editura Agro-Silvică de Stat, București, p 457 (in Romanian)
- Negulescu EG, Săvulescu AI (1965) Dendrology, 2nd edn. Editura Agro-Silvică de Stat, București, p 511 (in Romanian)
- Nicolescu VN (2019) Natural range, site requirements and shade tolerance. In: Spiecker H, Lindner M, Schuler J (eds) *Douglas-fir – an option for Europe*. European Forest Institute, Joensuu, pp 33–39
- Nicolescu VN, Mihăilescu Gh, Andronic I, Siladi RM, Tampa R, Beleşescu C, Bogdan RG (2021) A stand with Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) and Sitka spruce (*Picea sitchensis* (Bong.) Carr.) in Obștea Săliște Forest District (Vâlcea County). *Bucovina Forestieră* 21(1):73–81 (in Romanian)
- Nicoll BC, Gardiner BA, Rayner B, Peace AJ (2006) Anchorage of coniferous trees in relation to species, soil type, and rooting depth. *Can J Forest Res* 36:1871–1883
- Niinemets U, Valladares F (2006) Tolerance to shade, drought, and waterlogging of temperate northern hemisphere trees and shrub. *Ecol Monogr* 76(4):521–547
- Nițescu C, Achimescu C (1979) Technique of forest cultures. Tending operations. Editura Ceres, București, p 255 (in Romanian)
- Novák J, Kacálek D, Podrázský V, Šimerda L (eds) (2018) *Douglas-fir use in the forestry of the Czech Republic*. Lesnická práce, Praha, p 214 (in Czech)
- O'Reilly C, McCarthy N, Keane M, Harper CP, Gardiner JJ (1999) The physiological status of Douglas fir seedlings and the field performance of freshly lifted and cold stored stock. *Ann Sci for* 56:297–306
- Palik B, D'Amato AW, Franklin JF, Johnson KW (2021) *Ecological silviculture: foundations and applications*. Waveland Press, Illinois, p 43
- Pașcovschi S, Purcelean Ș (1954) Douglas-fir *Pseudotsuga taxifolia* Britt. (= *P. douglasii* Carr.). Technical guidelines for the culture of woody exotic species. In: Pașcovschi S, Purcelean Ș, Spârchez Z, Ocskay S, Beldie AI, Rădulescu S, Cocalcu T (eds) *Culture of woody exotic species*. Editura Agro-Silvică de Stat, București, pp 3–9 (in Romanian)
- Perić S, Tijardović M, Jazbec A (2011) Results of research on Douglas fir provenances in ecologically different continental parts of Croatia. *Šumarski List* 135(13):190–201 (in Croatian)
- Perin J, de Thier O, Claessens H, Lejeune Ph, Hébert J (2014) New harmonized productivity curves for Douglas-fir, Norway spruce and larches in Wallonia. *Forêt Wallonne* 129:26–41 (in French)
- Petersen R (2016) Single tree shelters and Douglas fir – does that work? *AFZ-DerWald* 71(24):17–20 (in German)
- Petkova K, Georgieva M, Uzunov M (2014) Investigation of Douglas-fir provenance test in North-Western Bulgaria at the age of 24 years. *J for Sci* 60(7):288–296
- Petkova K, Popov E, Tsvetkov I (2017) Country report. Bulgaria. In: Hasenauer H, Gazda A, Konner M, Mohren G, Pötzelsberger E, Spiecker H, van Loo M (eds) *Non-native tree species for European forests: Experiences, risks and opportunities*. COST Action FP1403 NNEXT Country Reports, Joint Volume. University of Natural Resources and Life Sciences Vienna, pp 40–60
- Petkova K (1989) Studies on establishing Douglas-fir (*Pseudotsuga menziesii* Mirb. (Franco) plantations in some regions of the Balkan Mountains. Dissertation. Higher Institute of Forestry, Sofia, p 190 (in Bulgarian)
- Petráš R, Mecko J (2008) Growth and production potential of Douglas-fir in Slovakia. In: *Growing of non-native tree species*. Česká lesnická společnost, Kroměříž, pp 59–64 (in Czech)
- Pintarić K (1990) Study of the growth of IUFRO Douglas fir from different provenances in two experimental areas in FR Bosnia and Herzegovina. *Forestry and Wood Processing* 7–12 XLIV: 113–125
- Podrázský V, Čermák R, Zahradník D, Kouba J (2013) Production of Douglas-Fir in the Czech Republic based on national forest inventory data. *J for Sci* 59(10):398–404
- Podrázský V, Mondek J, Zeidler A, Resnerová K, Borůvka V, Prknová H (2019) Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco). In: Podrázský V, Prknová H (eds) *Silvicultural, production and*

- environmental potential of the main introduced tree species in the Czech Republic. *Česka zemědělská univerzita v Praze, Lesnická práce*, Praha, pp 16–53
- Pollet C, Henin JM, Hébert J, Jourez B (2013) Effect of growth rate on the natural durability of Douglas-fir in western Europe. *Can J Forest Res* 43(10):891–896
- Pollet C, Henin JM, Hébert J, Jourez B (2017) Effect of growth rate on the physical and mechanical properties of Douglas-fir in western Europe. *Can J Forest Res* 47(8):1056–1065
- Polman JE, Militz H (1996) Wood quality of Douglas fir (*Pseudotsuga menziesii*(Mirb) Franco) from three stands in the Netherlands. *Ann Sci for* 53:1127–1136
- Popov E, Hinkov G, Velichkov I, Stefanova P (2018) Productivity of Douglas fir plantations in Bulgaria and the necessity for its utilisation. *Nauka Za Gorata* 2:3–23 **(in Bulgarian)**
- Popov E (1991) Study on the results from the introduction of Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) in Bulgaria. Dissertation. Bulgarian Academy of Sciences, Forest Research Institute, Sofia, p 162 **(in Bulgarian)**
- Popov E (2006) Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franko) plantations in the region of Koprivshtitsa". In: Proceedings of the international scientific conference Science in globalization in 21st century. Vol I Plant studies, technological questions in Plant Studies, Genetics and selection, weeds, diseases and enemies. 1–2 June 2006, Stara Zagora Union of Scientist, Stara Zagora, pp 158–163 **(in Bulgarian)**
- Popov E (2009) Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) plantations in the region of Rila monastery. In: Dimov D, Pavlov D, Eneva S, Videva M, Delchev G (eds.) Proceedings of the international scientific conference Economics and Society development on the Base of Knowledge, 4–5 June 2009, Stara Zagora, Bulgaria. Vol I. Agricultural science. Plant Studies, pp 370–375 **(in Bulgarian)**
- Puettmann KJ, Ares A, Burton JI, Dodson EK (2016) Forest Restoration using variable density thinning: lessons from douglas-fir stands in western Oregon. *Forests* 7:310. <https://doi.org/10.3390/f7120310>
- Pulkrab K, Sloup M, Zeman M (2014) Economic impact of Douglas-fir (*Pseudotsuga menziesii* [Mirb.] Franco) production in the Czech Republic. *J for Sci* 60(7):297–306
- Rădulescu I, Cazacu C (1968) Douglas-fir culture in Banat. In: Bakoş V (coord) Culture of fast growing tree species. Editura Agrosilvică, Bucureşti, pp 159–184 **(in Romanian)**
- Raida D (2018) Douglas fir (*Pseudotsuga menziesii* (Mirb.) Franco) in Ravbarkomanda plantations after the 2014 ice break. In: Diploma work. Biotehniška fakulteta, Ljubljana, p 38 **(in Slovenian)**
- Rameau JC, Mansion D, Dumé G (1989) French Forest Flora. In: Plains and hills. IDF, Paris, ENGREF, Nancy, p 1785 **(in French)**
- Ratknić M (1995) Volume tables of Douglas fir and Weymouth pine trees in Serbia. *Zbornik Radova Instituta Za Šumarstvo* 38–39:152–160 **(in Serbian)**
- Remeš J, Zeidler A (2014) Production potential and wood quality of Douglas fir from selected sites in the Czech Republic. *Wood Research* 59(3):509–520
- Restaino CM, Peterson DL, Littell J (2016) Increased water deficit decreases Douglas fir growth throughout western US forests. *PNAS* 113(34):9557–9562
- Riou-Nivert PH (1989) Douglas-fir, wood qualities, pruning and silviculture. *Revue Forestière Française* XL I(5):387–410 **(in French)**
- Riou-Nivert PH (1996) Conifères. Volume I: knowledge and recognition. Institut pour le développement forestier, Paris, p 256 **(in French)**
- Riou-Nivert PH (2020) Diversified silvicultural routes for Douglas-fir. 40 years of experience in private forests. Centre national de la propriété forestière (CNPF), Paris, p 27 **(in French)**
- Roques A, Auger-Rozenberg MA, Capretti P, Sauvard D, La Porta N, Santini A (2019) Pests and diseases in the native and European range of Douglas-fir. In: Spiecker H, Lindner M, Schuler J (eds) Douglas-fir - an option for Europe. European Forest Institute, Joensuu, pp 63–72
- Rose R, Rosner L (2005) Eighth-year response of Douglas-fir seedlings to area of weed control and herbaceous versus woody weed control. *Ann Sci for* 62:481–492
- Royal forestry society (2021) Managing for resilience. 10 case studies. The royal forestry society, Banbury, p 40
- Savill P (2019) The silviculture of trees used in British forestry, 3rd edn. CAB International, Wallingford and Boston, p 400
- Schönauer H (2008) Silviculture with Douglas-Fir. BFW-Praxisinformation 16:9–11 **(in German)**
- Schüler S, Chakraborty D (2021) Limiting factors for Douglas-fir cultivation in Central Europe under climate change. *Schweiz Z Forstwes* 172(2):84–93 **(in German)**
- Schütz JPH (1990) Silviculture 1. Principles of forest education. Presses polytechniques et universitaires romandes, Lausanne **(in French)**
- Schütz JPH, Pommerening A (2013) Can Douglas fir (*Pseudotsuga menziesii*(Mirb.) Franco) sustainably grow in complex forest structures? *For Ecol Manag* 303:175–183
- Schütz JPH, Amman PL, Zingg A (2015) Optimizing the yield of Douglas-fir with an appropriate thinning regime. *Eur J Forest Res* 134:469–480
- Scotti R (2016) 10-year projection of the value of a plant: dendro-auxometric modeling. In: Polistampa, Firenze, (eds) Realities and perspectives in the cultivation of Douglas fir in Italy. *Atidella Accademia dei Georgofili* 8 (13), pp 65–86 **(in Italian)**
- Sharpe AL, Mason WL, Howes REJ (1990) Early forest performance of roughly handled Sitka spruce and Douglas fir of different plant types. *Scott for* 44(4):257–265
- Siepmann F (1981) Comparison of the pruning criteria of the year 1950 with the current ones. *Allgemeine Forstzeitschrift* 30:758–760 **(in German)**
- Sillett SC, van Pelt R, Freund JA, Campbell-Spickler J, Carroll AL, Kramer RD (2018) Development and dominance of Douglas-fir in North American rainforests. *For Ecol Manag* 429:93–114
- Şimon DC, Stroe IM, Ostafi MF (2014) Study trip to Baden-Württemberg (Germany), 2014 edition. *Revista pădurilor* 3–4:121–123 **(in Romanian)**
- Smolnikar P, Brus R, Jarni K (2021) Differences in growth and log quality of Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) provenances. *Forests* 12(3):287. <https://doi.org/10.3390/f12030287>
- Şofletea N, Curtu L (2007) Dendrology. Editura Universităţii Transilvania, Braşov, p 418 **(in Romanian)**
- Šoškić B, Stojičić D, Petrović D (2007) Wood density and strength properties of Douglas-fir, Weymouth pine and Austrian pine wood. *Forestry (belgrade)* 3–4:1–11 **(in Serbian)**
- Spellmann H, Weller A, Brang P, Michiels HG, Bolte A (2015) Douglas fir (*Pseudotsuga menziesii* (Mirb) Franco). In: Vor T, Spellmann H, Bolte A, Ammer C (eds) Potentials and risks of introduced tree species. Göttinger Forstwissenschaften, Göttingen, pp 187–217 **(in German)**
- Spiecker H (2019) Executive summary. In: Spiecker H, Lindner M, Schuler J (eds) Douglas-fir - an option for Europe. European Forest Institute, Joensuu, pp 13–16
- Spiecker H, Schuler J (2019) Introduction. In: Spiecker H, Lindner M, Schuler J (eds) Douglas-fir - an option for Europe. European Forest Institute, Joensuu, pp 17–19
- Stănescu V (1979) Dendrology. Editura Didactică şi Pedagogică, Bucureşti, p 470 **(in Romanian)**
- Stănescu V, Şofletea N, Popescu O (1997) Woody Forest Flora of Romania. Editura Ceres, Bucureşti, p 451 **(in Romanian)**

- Stimm B (2004) Exotics in Bavarian Forests – historical legacy or valuable enrichment? *Lwfaktuell* 45:4–6 (in German)
- Stokes V, Kerr G, Connolly T (2021) Underplanting is a practical silvicultural method for regenerating and diversifying conifer stands in Britain. *For Int J for Res* 94(2):219–231. <https://doi.org/10.1093/forestry/cpaa027>
- Suzuki K, Asaoka I, Tani S, Okada K, Hidaka T (1995) Effects of pretreatments for the amelioration of preservative impregnability using the oscillating pressure method (OPM). In: Document-the international research group on wood preservation, IRG Secretariat, Stockholm, Sweden, issue 95–40044
- Sychra D, Mauer O (2013) Prosperity of Douglas fir (*Pseudotsuga menziesii* [Mirb.] Franco) plantations in relation to the shelter. *J for Sci* 59(9):352–358
- Talbert C, Marshall D (2005) Plantation productivity in the Douglas-fir region under intensive silvicultural practices: Results from research and operations. *J Forest* 103(2):65–70
- Ťavoda P (2007) Ecological demands and the spread of Douglas-fir in Slovakia. *Ekológia environmentalistika*:194–202 (in Slovak)
- Thomas FM, Bögelein R, Werner W (2015) Interaction between Douglas fir and European beech: investigations in pure and mixed stands. *Forstarchiv* 86(4):83–91 (in German)
- Thomas FM, Rzepecki A, Werner W (2022) Non-native Douglas fir (*Pseudotsuga menziesii*) in Central Europe: ecology, performance and nature conservation. *For Ecol Manag*. <https://doi.org/10.1016/j.foreco.2021.119956>
- Thurm EA, Uhl E, Pretzsch H (2016) Mixtures reduces climate sensitivity of Douglas-fir stem growth. *For Ecol Manag* 376:205–220
- Turner J, Mitchell SJ (2003) The effect of short day treatments on containerized Douglas-fir morphology, physiology and phenology. *New for* 26:279–295
- van Loo M, Dobrowolska D (2019a) History of introducing Douglas-fir in Europe. In: Spiecker H, Lindner M, Schuler J (eds) *Douglas-fir-an option for Europe*. European Forest Institute, Joensuu, pp 21–25
- van Loo M, Dobrowolska D (2019b) Current situation. In: Spiecker H, Lindner M, Schuler J (eds) *Douglas-fir - an option for Europe*. European Forest Institute, Joensuu, pp 26–29
- van Vredenburg CJH, La Bastide JGA (1968) The influence of meteorological factors upon the cone crop of Douglas-fir in the Netherlands. *Silvae Genetica* 18:182–186
- Vidaković M, Franjić J (2004) *Gymnosperms*. 3rd edn. Faculty of Forestry, University of Zagreb, Croatian Forests Ltd., Croatian Academy of Forest Sciences, Zagreb, p 823 (in Croatian)
- Vor T, Spellmann H, Bolte A, Ammer C (2015) Potentials and risks of introduced tree species. *Universitätsverlag Göttinger Forstwissenschaften*, Göttingen, p 228 (in German)
- Wallertz K, Malmqvist C (2013) The effect of mechanical site preparation methods on the establishment of Norway spruce (*Picea abies* (L.) Karst.) and Douglas fir (*Pseudotsuga menziesii* (Mirb.) Franco) in southern Sweden. *Forestry* 86:71–78
- Weise U, Flöß M, Kenk G (2001) Management and economic yield of Douglas fir stands in Baden-Württemberg. *AFZ-Der Wald* 58:802–806 (in German)
- Weiskittel AR, Crookston NL, Rehfeldt GE (2012) Projected future suitable habitat and productivity of Douglas-fir in western North America. *Schweiz Z Forstwes* 163(3):70–78
- Wilson JS, Baker PJ (2001) Flexibility in forest management: managing uncertainty in Douglas-fir forests of the Pacific Northwest. *For Ecol Manag* 145:219–227
- Wilson SMCG, Cameron AD (2015) Alternative models for productive upland forestry Model 2: Sitka spruce mixtures with alternative conifers. *Scott for* 69:26–32
- Wohlgemuth T, Hafner J, Höltermann A, Moser B, Nehring S, Rigling A (2019) Impact of Douglas-fir on forests and open land habitats. In: Spiecker H, Lindner M, Schuler J (eds) *Douglas-fir - an option for Europe*. European Forest Institute, Joensuu, pp 57–62
- Wohlgemuth T, Moser B, Pötzelberger E, Rigling A, Gossner MM (2021) About the invasiveness of Douglas-fir and its impact on soil and biodiversity. *Schweiz Z Forstwes* 172(2):118–127 (in German)
- Wouters Ph de, Lorent V (2002) Douglas-fir *Pseudotsuga menziesii*. *Silva Belgica* 109 (5), Sylviculture-fiche technique no. 12: 1-9
- Zeidler A, Borůvka V, Černý J, Baláš M (2022) Douglas-fir outperforms most commercial European softwoods. *Ind Crops Prod* 181:114828. <https://doi.org/10.1016/j.indcrop.2022.114828>

Websites

- <https://ddg-web.de/rekordbaeume.html?VCardId=6735> (Retrieved March 9, 2022)
- <https://experts-forestiers.be/Tableauprixbois.pdf> (Retrieved March 11, 2022)
- <http://inventaire-forestier.ign.fr> (Retrieved March 5, 2022)
- <http://www.bomeninfo.nl/tall%20trees.htm> (Retrieved July 5, 2021)
- <http://www.douglasfir.co.nz/net/> (Retrieved September 22, 2021)
- <http://www.frenchtimber.com/en/french-species/douglas-fir/> (Retrieved March 10, 2022)
- <https://www.forstpraxis.de/douglasie-schwarzwald-rekordmenge-rekorderloes> <http://www.nzffa.org.nz/> (Retrieved June 15, 2021)
- https://www.landesforsten.de/wp-content/uploads/2021/03/ergebnisse_submission_nadelholz_2021.pdf (Retrieved May 5, 2021)
- <https://www.lwk-niedersachsen.de/index.cfm/portal/4/nav/292/article/30508.html> (Retrieved May 10, 2021)
- <http://www.nzffa.org.nz/farm-forestry-model/> (Retrieved September 22, 2021)
- <https://www.treeregister.org/champion> (Retrieved January 8, 2022)
- https://www.forestiere-cdc.fr/sites/default/files/2021-06/prix-vente-des-bois-sur-pied-en-foret-privee-indicateur-2021_0.pdf (Retrieved March 11, 2022)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.