Programme and Abstracts of Conference on "Ecological problems of tourist lakes"



20-23 June 2011, Tihany, Hungary

Balaton Limnological Research Institute of the Hungarian Academy of Sciences



Unexpected cyanobacterial dominance in a deep oligomesotrophic lake, Lake Stechlin, Germany: *Aphanizomenon flos-aquae*, an ecosystem engineer

Judit Padisák¹, Viktória Üveges¹, Kálmán Tapolczai¹, Géza Balázs Selmeczy¹, Nico Salmaso², Peter Kasprzak³, Rainer Koschel³, Lothar Krienitz³

¹University of Pannonia, Department of Limnology, Veszprém, Hungary

²IASMA Research and Innovation Centre, Environment and Natural Resources Area, Fondazione Edmund Mach, S. Michele all'Adige, TN, Italy

³Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Stechlin-Neuglobsow, Germany

Recent research detected an ongoing expansion of cyanobacteria in European lakes with moderate tropic states without parallel increase in trophic state indicators. A perennial Aphanizomenon flos-aquae bloom in Lake Stechlin, Germany in 2009-2010 represents such a case. Lake Stechlin is a deep (zmax: 69.5 m) glacial lake that used to be considered oligotrophic in the recent past of its history. Records on phytoplankton are available since 1994. The first filament of Aphanizomenon flos-aquae appeared in the lake in 2000 and then it developed minor peaks by every late summers. In 2009, however, an intensive growth started reaching a maximum (310 µg L-1) in August. After a decline in August a winter population started to develop with a maximum around 920 µg L⁻¹ in December-February that persisted almost as monoculture under thick ice and snow. This project was implemented through the CENTRAL EUROPE Programme (European Lakes Under Environmental Stressors, 2CE243P3) and the project Klimagetriebene Veränderungen der Biodiversität von Mikrobiota (TemBi) supported by the Leibniz Gemeinschaft (WGL).