



Sotto l'Alto Patronato del Presidente della Repubblica  
*Under the high patronage of the President of the Italian Republic*



**ATTI**  
**del Secondo Congresso Internazionale di Selvicoltura**  
**Progettare il futuro per il settore forestale**

Firenze, 26-29 Novembre 2014

**PROCEEDINGS**  
**of the Second International Congress of Silviculture**  
**Designing the future of the forestry sector**

Florence, 26-29 November 2014

**VOL. II**

**Accademia Italiana di Scienze Forestali**  
Firenze - 2015

Quanto esposto è di esclusiva proprietà scientifica e intellettuale degli Autori ed esclude ogni responsabilità del curatore e dell'Editore.

*Intellectual and scientific property is exclusively of the authors of each contribution and does not entail responsibility of the editor and the publisher*

A cura di / *Edited by*  
Orazio Ciancio

Con la collaborazione di / *In collaboration with*  
Alga Ciuti, *Accademia Italiana di Scienze Forestali*  
Chiara Lisa, *Accademia Italiana di Scienze Forestali*  
Caterina Morosi, *Accademia Italiana di Scienze Forestali*  
Francesco Paolo Piemontese, *Università degli Studi di Firenze*  
Giovanna Puccioni, *Accademia Italiana di Scienze Forestali*

Gli Atti sono stampati grazie al contributo della Regione Toscana.

*These Proceedings have been printed with the financial support of the Tuscany Region*

© 2015 Accademia Italiana di Scienze Forestali  
Piazza Edison 11  
50133 Firenze

ISBN 978-88-87553-21-5

## CLIMATE SIGNALS DERIVED FROM DAY-TO-DAY ANALYSIS: CLIMATE SENSITIVITY OF *PICEA ABIES* IN TRENTINO

Claudia Coccozza<sup>1</sup>, Caterina Palombo<sup>1</sup>, Monica Anichini<sup>2</sup>, Roberto Tognetti<sup>3</sup>, Alessio Giovannelli<sup>2</sup>, Nicola La Porta<sup>3,4</sup>, Giovanni Emiliani<sup>2</sup>

<sup>1</sup>Dipartimento di Bioscienze e Territorio, Università del Molise, Pesche, Italy; claudia.coccozza@unimol.it

<sup>2</sup>Laboratorio di Xilogenesi, Istituto per la Valorizzazione Legno e delle Specie Arboree (IVALSA), Consiglio Nazionale delle Ricerche, Sesto Fiorentino, Italy

<sup>3</sup>Dipartimento di Bioscienze e Territorio, Università del Molise; The EFI Project Centre on Mountain Forests (MOUNTFOR), Edmund Mach Foundation, San Michele all'Adige, Italy

<sup>4</sup>The EFI Project Centre on Mountain Forests (MOUNTFOR), Edmund Mach Foundation; Department of Sustainable Agro-Ecosystems and Bioresources, IASMA Research and Innovation Centre, Edmund Mach Foundation, San Michele all'Adige, Italy

Climate changes directly affect tree growth. Tree performance under changing conditions is maintained by xylem structure that determines the hydraulic and mechanical properties of a stem. The interest for the understanding of how trees grow during the climate change has led to the development of efficient methods for measuring and monitoring growth. Xylogenesis and annual radial increments of Norway spruce (*Picea abies* (L.) Karst.) in Trentino (Eastern Italian Alps) were monitored between 2010 and 2012 with automatic dendrometers and cell analysis. The analyses of intra-annual dynamics of wood formation were used to describe seasonal changes in xylem differentiation phases and to calculate the timing of cell development in Norway spruce. The investigation was conducted in two sites at different elevation, Savignano (650 m a.s.l.) and Lavazè (1800 m a.s.l.), to detect the effect of climate signals in the day-to-day dynamics of wood formation. Dendroclimatological analysis was performed to examine the relationship between the formation of tree rings of Norway spruce and climatic parameters in two contrasting sites. Climate-growth relationships were analyzed using correlation function analysis and moving correlation function, detecting relations between phenological phases of wood formation and seasonal patterns of temperature and precipitation. Effects of climatic variables on stem diameter variation and cell structure were examined, respectively, daily and each 15 days. The results were interpreted according to the dynamics of forest vegetation and synchronicity of cambial activity with meteorological parameters. Models of cambial and tree growth are required to improve the predictive power of ecosystem process models, in which tree growth is often an essential and complex component.

*Keywords:* cell structure, tree-ring, dendroclimatology, tree growth response, moving correlation analysis.

*Parole chiave:* struttura cellulare, anello annuale, dendroclimatologia, risposte di crescita, analisi di correlazione.

<http://dx.doi.org/10.4129/2cis-cc-cli>

### 1. Climate disturbances and effects on tree growth and survival

European forests are continuously exposed to increasing disturbances due to ongoing climatic changes. These changes impact the forest carbon budget and are suggested to contribute to the recently observed carbon sink saturation in European forests throughout the twentieth century (Seidl *et al.*, 2014). The exponential increase of vapor pressure deficit (VPD) with warming air temperature, at constant absolute humidity, may trigger changes in the sensitivity and vulnerability to disturbances, which in turn may increase the risk of tree mortality once the threshold of tree resilience is overcome (Fig. 1).

We generally think of gradual linear changes in climate and ecosystem status. However, abrupt climatic change

can lead to abrupt ecosystem change, while gradual climatic change may also trigger abrupt ecosystem change. Forests will, therefore, have to adapt to non-linear risk of erratic and extreme weather events, such as prolonged drought, heat waves, storms and floods (Lindner *et al.*, 2010). In particular, increasing droughts and warm temperatures are supposed to induce “physiological stress” (because of water and carbon imbalance), determining “forest mortality” under a changing climate (Allen *et al.*, 2010), contributing to abrupt ecosystem changes.

Tree mortality constitutes a major element of forest dynamics, influencing species distributions, energy fluxes, hydrological processes, and carbon fluxes (Adams *et al.*, 2009). However, the processes of tree death are poorly understood, since they often result from a complex of multiple abiotic and biotic environmental factors that

occur consecutively in time. Tree mortality can be described in terms of *growth-dependent* mortality factors, which affect tree vigor, sometimes for decades prior to death, also due to opportunistic pathogens of host chronic weakness, and *growth-independent* factors, which directly lead to tree death, often without being detectable in tree growth pattern. The critical importance of understanding climate sensitivities of forests is associated with the mechanisms that trigger plant mortality and drive vegetation change and their implications for assessments of climate change impacts and consequent land surface-atmospheric feedbacks (Anderegg *et al.*, 2013). For instance, drought has been frequently discussed as a trigger for forest decline (Eilmann *et al.*, 2011), causing directly mortality via hydraulic failure in extreme events (Bréda *et al.*, 2006), or indirectly as a persistent weakening of trees by reducing carbon storage (McDowell *et al.*, 2008; Deslauriers *et al.*, 2014). At an increase of drought stress corresponds a reduction of tree tolerance to stress factors (Rebetz and Dobbertin, 2004), as well as phyllophagous insects, pathogenic fungi, frost, air pollutants and mechanical injury, which may lead to a strongly decrease in tree vigor. Consequently, the impacts of regional tree die-off can produce change in forest growth conditions acting as a combination of abiotic stress (e.g., drought spells and heat waves) and amplifying biotic agents able to reduce ecosystem resilience (Van Leeuwen *et al.*, 2000). For instance, the reduction of habitat for wildlife and ecosystem potential to sequester carbon can induce the enhancing of opportunities for invasion by exotic species and formation of novel communities, with consequences in alterations of hydrologic cycle and temporal disruptions in ecosystem goods and services (Adams *et al.*, 2009). Water shortage, in particular, is likely to become a major factor limiting species distribution and establishment in the near future (IPCC, 2007), leading to increased forest decline and rapid decline-induced vegetation shifts (Eilmann *et al.*, 2011).

## 2. Tree adaptation detection

Emerging drought risks for trees have effects on the anatomical structure, as well as on the physical and chemical properties of wood. However, trees have evolved phenological, morphological and physiological adaptation systems to cope with drought conditions (Battipaglia *et al.*, 2010; Pineda-Garcia *et al.*, 2013), inducing specific patterns of cambial activity and changes in xylem-hydraulic conductivity and vulnerability to cavitation (Sperry, 2011).

Climate change has been also implicated in shifting phenological patterns of wood formation and bud burst, such as the timing and duration of the growing season across a wide variety of ecosystems (Rossi *et al.*, 2013). However, biological processes frequently follow more complex and non-linear response patterns, according to limiting factors that generate shifts and discontinuities (Rossi *et al.*, 2013). Thus, the shift of one phenological phase is associated with synchronous and comparable shifts of the following phases. However, small increases in the duration of xylogenesis could correspond to a

substantial increase in cell production. These findings suggest that changes of the length of the growing season could be substantial in terms of cell production and carbon uptake by trees, and consequently forest productivity (Rossi *et al.*, 2013). Modifications of the environmental conditions that affect plant phenology could potentially engender disproportionate responses in plant growth mechanisms (Fig. 2). Tree performance under changing conditions is maintained by the xylem structure, that determines the hydraulic functionality and mechanical properties of a stem. In this sense, tree ring anatomy and stem diameter monitoring can provide a mechanistic and functional understanding of xylem plastic responses, which can differ in times and fulfill differing functions that define climate-growth relationships (Fonti *et al.*, 2013).

The cause-effect relationships between xylogenesis, structure and function of the xylem can be linked through a concerted effort involving multiple disciplines, as functional ecology, wood anatomy, plant genetic, physiology and dendrochronology (Battipaglia *et al.*, 2014). The activation and cessation of plant growth during the seasonal course has been mainly related to temperature and water availability (Eilmann *et al.*, 2011; Coccozza *et al.*, 2012).

The complexity of the interaction between environmental conditions and growth dynamics is indicated by the varying degree of temporal correspondences between stem radius variations and specific climate variables (e.g., Zweifel *et al.*, 2010).

The irreversible radial growth and the reversible living-cell dehydration-rehydration processes are related to the depletion and replenishment of stem water stored in woody tissues, and they are detectable through stem diameter variations obtained from daily dendrometer measurements (Steppe *et al.*, 2006). Turgor pressure drives irreversible cell expansion and deposition of wall polymers, that define cellular enlargement and, consequently, radial growth, as water-dependent processes (Deslauriers *et al.*, 2007).

For instance in conifers, the tissue water-content-related stem diameter variations are largely determined by soil water availability and atmospheric VPD (Zweifel *et al.*, 2005). Whereas, phase duration, taken as an indicator of the reversible changes in stem radius, depends mainly on transpiration and sap flow (Steppe *et al.*, 2006).

## 3. A study case: high-resolution analysis of stem growth in *Picea abies* in Northern Italy

Significant interest in how trees grow has led to advances in several methods for measuring and monitoring stem diameter variation, each of them with their own advantages and disadvantages. Timing and dynamics of xylogenesis are crucial aspects for understanding the effects of climate changes on forest ecosystems because they represent the time “window” in which environmental factors directly affect growth (Rossi *et al.*, 2007). Dynamics of wood formation are similar in all species; the onset of cell production occurs in the first part of the growing season, while cambial activity, synchronized with photoperiod, reaches maximum rates

during summer solstice (Rossi *et al.*, 2007). Intra-annual high-resolution stem growth analysis is a first attempt to better understand mechanisms of tree growth processes in relation to the environment. A general hypothesis is that changes in cambium phenology with increasing temperature will be more consistent at higher elevations, with significant variation in rate and duration of the cell division, enlargement and lignification, than at mid-low elevation.

Although the complex interactions between environmental conditions and plant traits are non linear responses and dendrometric records remain often difficult to interpret, the sensitivity of signal recording in the present experiment enabled to determine synchronisation between plant signals and environmental variables.

The sensitivity of tree growth components (change from each measured value to the next) to environmental conditions was tested through high-resolution approaches capable to detect changing in time (dendrometers) and space (cambium anatomy). These tools gave us the possibility to detect (i) stem radius variations (irreversible and reversible radius variation), and (ii) the seasonal course of cambium activity, as a result of the ability of Norway spruce (*Picea abies* (L.) Karst.) to adapt to local conditions. We expected that the changing temperature affects the timing and capacity of cell production of cambium, allowing at lower elevation a longer growth season, and resulting in increased cell production and radial growth. In order to detect temperature-dependent growth strategies of Norway spruce, adult trees of similar age in two sites of Trentino, differing in altitudes, were chosen as model system for an integrated approach of monitoring stem diameter variation and cambial cell production. Tree growth at different altitudes was continuously monitored through dendrometers for three years. On the same trees, microcores were sampled bi-weekly throughout one growing season for examining cambial activity. The strong interest in monitoring forest health is well justified by the importance of forests that in Trentino cover more than 60% of land, characterizing economy, landscape and culture. Since changing disturbance regimes are increasingly challenging for sustainable forest management, they can influence the choices of close-to-nature management aiming to enhance forest ecosystem services, saving ecological specificity and environmental resilience.

Sampling sites were: Pomarolo (45° 56' 41" N; 11° 03' 28" E) and Lavazè (46° 21' 37" N; 11° 29' 42" E). The first one is representative of deciduous forest (*Quercetum pubescentis*) at low altitude (680 m a.s.l.), dominated by *Quercus pubescens*, *Quercus robur*, *Fraxinus ornus* and *Ostrya carpinifolia*, with secondary presence of *Castanea sativa*, *Pinus silvestris*, *Larix decidua* and *Picea abies*. The second, located at 1780 m a.s.l., is representative of the association *Picetum subalpinum* mainly formed by *Picea abies* with rare presence of *Pinus cembra* and *Larix decidua*. The climate in Pomarolo is mild continental with sub-Mediterranean influences; mean annual temperature is around 11 °C and mean annual precipitation is 1150 mm. This site is located on cambisol, in a limestone area. In Lavazè the climate is alpine-continental and the annual mean temperature is 4 °C, while the mean annual

precipitation are 1100 mm. The stand grows on podsollic soil above quartz porphyric rocks.

In both sites, a meteorological automatic station was placed in a nearby open area (WMO 1998). Precipitation, air temperature, relative humidity and global radiation were recorded several times per hour and all these values were hourly averaged. In order to follow the seasonal progression of cambial cell development and stem diameter variation, two methods, dendrometers and microcoring, were used. Stem radius variations were daily monitored from January 2010 to December 2012 using automatic point dendrometers (Label *et al.*, 2000) on five individual Norway spruce trees in each site. We posed attention in selecting healthy and vigorous trees within 25-30 m of distance to each other, with similar diameter ranging from 35-40 cm at breast height. The used dendrometers measure the linear displacement of a sensing rod pressed against the bark. The operating principle of the linear variable transducer (AB Electronics Ltd., Romford, Essex, UK) that responds to stem radius variation is described elsewhere (Giovannelli *et al.*, 2007; Coccozza *et al.*, 2009). Trees were monitored with these high-resolution automated dendrometers installed on the trunk at 130 cm from the soil surface. Raw data were recorded every 15 min, and hourly and daily averages were then calculated.

The sampling for the analysis of wood formation in three replicate plants of Norway spruce was performed from 16 April to 13 October 2010 every fifteen days, and repeated the 23 November for control. The sampling consisted in the collection of two microcores (1.8 mm in diameter, 15 mm in length) containing the bark, cambium and the last formed tree-rings from each of the selected stems using Trephor corer (Rossi *et al.*, 2006a). Immediately after the sampling, the microcores were put in Eppendorf vials with 75% ethanol and 25% acetic acid for 24 h and then stored in 70% alcohol solution to preserve forming cells from degradation. Transversal microsections 10-12 µm thick were cut by a sliding microtome, restraining the microcores with a special holder. The microsections were stained with Safranin and Astra-blue, dehydrated with graded series of ethanol, and fixed to microscope slides with Canada balsam. Ring formation was assessed by anatomical observations with an Olympus BX 40 light microscope at a magnification of 400x. Observations consisted in counting the number of cells in the cambial zone and tracheids in enlargement, wall thickening and mature phases along three radial files on each microsection (Rossi *et al.*, 2006b).

Both the methodologies stressed a distinct growth pattern throughout the season in each of the two sites, which was related to altitude and temperature. Correlations between parameters derived from dendrometers (maximum daily shrinkage and stem increment) and environmental variables highlighted the effects of climate on stem diameter fluctuations and radial growth patterns. The onset of radial growth and intra-annual variations of stem diameter were differently defined at each elevation. In trees at higher elevation (Lavazè), intra-annual variations in stem diameter revealed weaving peaks of radial increment, which may be the consequence of distinct stem rehydration/dehydration

cycles. Whereas, at lower elevation tree growth was more regular, showing a constantly increasing trend (Fig. 3).

The vegetative season, with the formation of enlargement cells, occurred starting from May (DOY 123) in Pomarolo and from June (between DOY 137 and 151) in Lavazè, while the last developing cells were detected in September (DOY 256) in Pomarolo and at the end of August (DOY 242) in Lavazè. The mean number of cells produced in 2009 was 163 in Pomarolo and 27 in Lavazè. Trees showed different timing of wood formation at the two altitudes, with a longer xylogenesis in Pomarolo ( $\approx 150$  d) than in Lavazè ( $\approx 70$  d). The onset of cambial activity was earlier and ended later in trees at lower elevation (Pomarolo), confirming that the start of cell division is a key factor determining the duration of xylogenesis and differences in tree ring widths.

#### 4. Conclusion and perspectives

Under global warming, energy-limited forests, like those at higher elevation (Lavazè), can decrease productivity and more easily reach and overcome the threshold of mortality because of increasing water deficit. Again, it is not clear whether plasticity in xylogenesis or disproportionate increases in xylem cell production might occur in low-elevation trees (water-limited forests), with lengthening of the period of cell division. Research gaps and key questions need to be filled and answered for predicting how trees will respond to projected climate change.

The development of mechanistic models can provide a more complete picture of forest resilience to environmental disturbance (Fig. 4).

Indeed, at the regional level, a mechanistic understanding of the connections between the hydrological and carbon cycles in different species is needed to elucidate the impacts of climate change on tree growth or tree mortality. As an example, Scots pine under drought stress has been found to build a more effective water-conducting system (e.g., larger tracheids) in a shorter period of wood formation, at the cost of a probably higher vulnerability to cavitation (larger tracheids with thinner

cell walls), but without losing the capability to recover (Eilmann *et al.*, 2011). The shortening of the growth period in trees under increasing dry conditions would indicate that the period in which wood formation takes place could become much shorter, under the projected climate scenario for the Alpine environment, than the potential and actual phenological growth period.

Although several tree-ring studies have been carried out in *P. abies*, how its radial growth at intra-annual resolution is influenced by weather conditions has scarcely been explored. In this study, technology advancements have contributed to the ability to collect information at high spatial and temporal resolution and to store large amounts of data (Battipaglia *et al.*, 2014).

Attempting to isolate a growth signal from large amounts of data has proven challenging, though feasible. Many researchers have used daily weather variables to explain variability of the “isolated growth” (Deslauriers *et al.*, 2003). Further work involving simple physiological and functional models will also expand our understanding of tree water deficits, providing insight into the efficiency of species-specific response to drought conditions (Zweifel *et al.*, 2005). In this sense, cambium growth models, which inherently require a thorough understanding of influences on cambium activity, may provide a more complete basis for data interpretation, when using tree rings as an archive of environmental information (Breitenmoser *et al.*, 2013).

#### Acknowledgements

This work was supported by the project SOFIE-2 (Reg. Delib. N° 3012-2007) funded by the *Provincia Autonoma di Trento*, Italy. The authors would like to thank Mauro Confalonieri, Giorgio Zattoni, Bruno Crosignani of the Forest Service of the Autonomous Province of Trento for the collaboration to provide and to maintain the access to both sites along the years of the work and Cristina Salvadori of the Fondazione Edmund Mach for the kind suggestions for the choice of trees.




Figure 1. Scheme of the ecological factor drivers of stress resilience and tree mortality.




Figure 2. In a global warming scenario, it is questioned whether trees will adapt and adjust structurally and functionally in order to maintain growth performance or if they will reduce productivity and eventually overcome the threshold of mortality.




Figure 3. Radial growth of Norway spruce detected trough dendrometers in Lavazè (high elevation) and Pomarolo (low elevation) in the 2010 growing season.




Figure 4. Potential trajectories of abiotic constraints with global warming, as amplified by biotic agents, and consequent damages to tree functionality and forest ecosystem.

## RIASSUNTO

### Segnali del clima dall'analisi giorno per giorno: sensibilità al clima di *Picea abies* in Trentino

I cambiamenti climatici influenzano direttamente l'accrescimento delle piante. Le performance degli alberi in risposta ai cambiamenti climatici sono definite dalla struttura dello xilema che determina le proprietà idrauliche e meccaniche del fusto. L'interesse per i processi che regolano la crescita degli alberi ha definito diversi metodi di misura e monitoraggio. Le dinamiche della xilogenesi e di accrescimento del fusto di abete rosso (*Picea abies* (L.) Karst.) in Trentino sono state monitorate tra il 2010 e il 2012 mediante l'uso di dendrometri automatici e analisi cellulare. Le dinamiche annuali della formazione del legno sono state utilizzate per descrivere le variazioni stagionali delle fasi di differenziazione dello xilema e per calcolare lo sviluppo cellulare dell'abete rosso. Lo studio è stato condotto in due siti differenti per altitudine, Savignano (650 m s.l.m.) e Lavazè (1800 m s.l.m.), per identificare il segnale climatico, giorno per giorno, nella dinamica di formazione del legno. La relazione clima-crescita è stata studiata al fine di definire la possibile azione delle variabili climatiche sulla formazione del legno dell'abete rosso ed è stata analizzata mediante funzione di correlazione, rilevando relazioni tra la formazione del legno e gli andamenti stagionali di temperature e precipitazioni. Gli effetti delle variabili climatiche sulla variazione del diametro del fusto e sulla struttura cellulare sono state analizzate, rispettivamente, a scala giornaliera. I risultati sono stati interpretati seguendo le dinamiche forestali e la sincronicità dell'attività cambiale. Lo sviluppo di modelli di attività cambiale e di crescita degli alberi è necessario per migliorare la modellistica dei processi di ecosistema, dove una componente essenziale e complessa è proprio la crescita dell'albero.

## BIBLIOGRAPHY

- Adams H.D., Guardiola-Claramonte M., Barron-Gafford G.A., Villegas J.C., Breshears D.D., Zou C.B., Troch P.A., Huxman T.E., 2009 – *Temperature sensitivity of drought-induced tree mortality portends increased regional die-off under global-change-type drought*. PNAS, 106: 7063-7066.  
<http://dx.doi.org/10.1073/pnas.0901438106>
- Allen C.D., Macalady A.K., Chenchouni H., Bachelet D., McDowell N., Vennetier M., Kitzberger T., Rigling A., Breshears D.D., Hogg E.H., Gonzalez P., Fensham R., Zhang Z., Castro J., Demidova N., Lim J.H., Allard G., Running S.W., Semerci A., Cobb N., 2010 – *A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests*. Forest Ecology and Management, 259: 660-684.  
<http://dx.doi.org/10.1016/j.foreco.2009.09.001>
- Anderegg W.R.L., Kane J.M., Anderegg L.D.L., 2013 – *Consequences of widespread tree mortality triggered by drought and temperature stress*. Nature Climate Change, 3: 30-36.  
<http://dx.doi.org/10.1038/nclimate1635>
- Battipaglia G., De Micco V., Brand W.A., Linke P., Aronne G., Saurer M., Cherubini P., 2010 – *Variations of vessel diameter and delta 13C in false rings of Arbutus unedo L. reflect different environmental conditions*. New Phytologist, 188: 1099-1112.  
<http://dx.doi.org/10.1111/j.1469-8137.2010.03443.x>
- Battipaglia G., De Micco V., Sass-Klaassen U., Tognetti R., Mäkelä A., 2014 – *Editorial. Special issue: WSE symposium: Wood growth under environmental changes: the need for a multidisciplinary approach*. Tree Physiology, 34: 787-791.  
<http://dx.doi.org/10.1093/treephys/tpu076>
- Breda N., Huc R., Granier A., Dreyer E., 2006 – *Temperate forest trees and stands under severe drought: a review of ecophysiological responses, adaptation processes and long-term consequences*. Annals of Forest Science, Springer Verlag (Germany), 63: 625-644.  
<http://dx.doi.org/10.1051/forest:2006042>
- Breitenmoser P., Bronnimann S., Frank D., 2013 – *Forward modelling of tree-ring width and comparison with a global network of tree-ring chronologies*. Climate of the Past Discussions, 9: 4065-4098. <http://dx.doi.org/10.5194/cpd-9-4065-2013>
- Cocozza C., Lasserre B., Giovannelli A., Castro G., Fragnelli G., Tognetti R., 2009 – *Low temperature induces different cold sensitivity in two poplar clones (Populus × canadensis Monch 'I-214' and P. deltoides Marsh. 'Dvina')*. Journal of Experimental Botany, 60: 3655-3664.  
<http://dx.doi.org/10.1093/jxb/erp212>
- Cocozza C., Giovannelli A., Lasserre B., Cantini C., Lombardi F., Tognetti R., 2012 – *A novel mathematical procedure to interpret the stem radius variation in olive trees*. Agricultural and Forest Meteorology, 161: 80-93.  
<http://dx.doi.org/10.1016/j.agrformet.2012.03.016>
- Deslauriers A., Morin H., Urbinati C., Carrer M., 2003 – *Daily weather response of balsam fir (Abies balsamea (L.) Mill.) stem radius increment from dendrometer analysis in the boreal forests of Québec (Canada)*. Trees 17: 477-484.  
<http://dx.doi.org/10.1007/s00468-003-0260-4>
- Deslauriers A., Anfodillo T., Rossi S., Carraro V., 2007 – *Using simple causal modeling to understand how water and temperature affect daily stem radial variation in trees*. Tree Physiology, 27: 1125-1136.  
<http://dx.doi.org/10.1093/treephys/27.8.1125>
- Deslauriers A., Beaulieu M., Balducci L., Giovannelli A., Gagnon M.J., Rossi S., 2014 – *Impact of warming and drought on carbon balance related to wood formation in black spruce*. Annals of Botany, 114: 335-345. <http://dx.doi.org/10.1093/aob/mcu111>
- Eilmann B., Zweifel R., Buchmann N., Graf Pannatier E., Rigling A., 2011 – *Drought alters timing, quantity, and quality of wood formation in Scots pine*. Journal of Experimental Botany, 62: 2763-2771. <http://dx.doi.org/10.1093/jxb/erq443>
- Fonti P., Bryukhanova M.V., Mygland V.S., Kirilyanov A.V., Naumova O.V., Vaganov E.A., 2013 – *Tempe-*



- temperature-induced responses of xylem structure of Larix sibirica* (Pinaceae) from Russian Altay. *American Journal of Botany* 100: 1-12.  
<http://dx.doi.org/10.3732/ajb.1200484>
- Giovanelli A., Deslauriers A., Fragnelli G., Scaletti L., Castro G., Rossi S., Crivellaro A., 2007 – *Evaluation of drought response of two poplar clones (Populus x canadensis Monch 'I-214' and P. deltoides Marsh 'Dvina') through high resolution analysis of stem growth*. *Journal of Experimental Botany*, 58: 2673-2683.  
<http://dx.doi.org/10.1093/jxb/erm117>
- IPCC, 2007 – *Climate Change (2007): the physical science basis*. Contribution of working Group I to the fourth assessment report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge.
- Label P., Beritognolo I., Burtin P., Dehon L., Couée I., Breton C., Charpentier J.P., Jay-Allemand C., 2000 – *Cambial activity and xylem differentiation in walnut Juglans spp.* In: Savidge R.A., Bennett J.R., Napier R., eds. *Cell and molecular biology of wood formation*. Oxford: BIOS Scientific Publishers, 209-221.
- Lindner M., Maroschek M., Netherer S., Kremer A., Barbati A., Garcia-Gonzalo J., Seidl R., Delzon S., Corona P., Kolstro M., Lexer M.J., Marchetti M., 2010 – *Climate change impacts, adaptive capacity, and vulnerability of European forest ecosystems*. *Forest Ecology and Management*, 259: 698-709.  
<http://dx.doi.org/10.1016/j.foreco.2009.09.023>
- McDowell N.G., Pockman W.T., Allen C.D., Breshears D.D., Cobb N., Kolb T., Plaut J., Sperry J., West A., Williams D.G., Yezzer E.A., 2008 – *Mechanisms of plant survival and mortality during drought: why do some plants survive while others succumb to drought?* *New Phytologist*, 178: 719-739.  
<http://dx.doi.org/10.1111/j.1469-8137.2008.02436.x>
- Pineda-Garcia F., Paz H., Meinzer F.C., 2013 – *Drought resistance in early and late secondary successional species from a tropical dry forest: the interplay between xylem resistance to embolism, sapwood water storage and leaf shedding*. *Plant Cell and Environment*, 36: 405-418.  
<http://dx.doi.org/10.1111/j.1365-3040.2012.02582.x>
- Rebetez M., Dobbertin M., 2004 – *Climate change may already threaten Scots pine stands in the Swiss Alps*. *Theoretical and Applied Climatology*, 79: 1-9.  
<http://dx.doi.org/10.1007/s00704-004-0058-3>
- Rossi S., Anfodillo T., Menardi R., 2006a – *Trephor: a new tool for sampling micro-cores from tree stems*. *IAWA Journal*, 27: 89-97.  
<http://dx.doi.org/10.1163/22941932-90000139>
- Rossi S., Deslauriers A., Anfodillo T., 2006b – *Assessment of cambial activity and xylogenesis by micro-sampling tree species: An example at the Alpine timberline*. *IAWA Journal*, 27: 383-394.  
<http://dx.doi.org/10.1163/22941932-90000161>
- Rossi S., Deslauriers A., Anfodillo T., Carraro V., Carrer M., Urbinati C., Menardi R., Fontanella F., 2007 – *Effetti della temperatura e del fotoperiodo sulla xylogenesi al limite superiore del bosco*. *Italia Forestale e Montana*, 2: 81-97.  
<http://dx.doi.org/10.4129/IFM.2007.2.02>
- Rossi S., Anfodillo T., Čufar K., Cuny H.E., Deslauriers A., Fonti P., Frank D., Gričar J., Gruber A., King G.M., Krause C., Morin H., Oberhuber W., Prislan P., Rathgeber C.B.K., 2013 – *A meta-analysis of cambium-phenology and growth: linear and non-linear patterns in conifers of the northern hemisphere*. *Annals of Botany*, 1-10.
- Seidl R., Schelhaas M.J., Rammer W., Verkerk P.J., 2014 – *Increasing forest disturbances in Europe and their impact on carbon storage*. *Nature Climate Change*, 4: 806-810. <http://dx.doi.org/10.1038/nclimate2393>
- Sperry J.S., 2011 – *Hydraulics of vascular water transport*. In: *Mechanical integration of plant cells and plants*. Wojtaszek P., ed. Berlin: Springer, pp. 303-327.  
[http://dx.doi.org/10.1007/978-3-642-19091-9\\_12](http://dx.doi.org/10.1007/978-3-642-19091-9_12)
- Steppe K., De Pauw D.J.W., Lemeur R., Vanrolleghem P.A., 2006 – *A mathematical model linking tree sap flow dynamics to daily stem diameter fluctuations and radial stem growth*. *Tree Physiology*, 26: 257-273.  
<http://dx.doi.org/10.1093/treephys/26.3.257>
- Van Leeuwen E.P., Hendriks K.C.M.A., Klap J.M., DeVries W., De Jong E., Erisman W., 2000 – *Effects of environmental stress on forest crown condition in Europe. Part II: Estimation of stress induced by meteorology and air pollutants*. *Water, Air and Soil Pollution*, 119: 335-362.  
<http://dx.doi.org/10.1023/A:1005182226292>
- Zweifel R., Zimmermann L., Newbery D.M., 2005 – *Modeling tree water deficit from microclimate: an approach to quantifying drought stress*. *Tree Physiology*, 25: 147-156.  
<http://dx.doi.org/10.1093/treephys/25.2.147>
- Zweifel R., Eugster W., Etzold S., Dobbertin M., Buchmann N., Häslér R., 2010 – *Link between continuous stem radius changes and net ecosystem productivity of a subalpine Norway spruce forest in the Swiss Alps*. *New Phytologist*, 187: 819-830.  
<http://dx.doi.org/10.1111/j.1469-8137.2010.030>

**I PIANI DI ASSESTAMENTO FORESTALE QUALI STRUMENTI  
DI ATTUAZIONE DELLE MISURE DI TUTELA E CONSERVAZIONE  
DEI SITI RETE NATURA 2000 BASILICATA:  
LA ZSC “ABETINA DI LAURENZANA” (PZ)**

**Salvatore Cipollaro<sup>1</sup>, Maria Pompili<sup>2</sup>, Antonella Logiurato<sup>2</sup>, Antonio Racana<sup>1</sup>,  
Vito Orlando<sup>2</sup>, Gino Panzardi<sup>2</sup>, Antonio Bellotti<sup>1</sup>, Rocco Ferrari<sup>3</sup>**

<sup>1</sup>Regione Basilicata, Dipartimento Politiche Agricole e Forestali, Ufficio Foreste e Tutela del Territorio, Potenza; salvatore.cipollaro@regione.basilicata.it

<sup>2</sup>Regione Basilicata, Dipartimento Ambiente e Territorio, Infrastrutture, OO.PP. e Trasporti Ufficio Parchi, Biodiversità e Tutela della Natura, Potenza

<sup>3</sup>Dottore Forestale, libero professionista

La Regione Basilicata, pianifica le risorse silvo-pastorali attraverso la redazione dei PAF (Piani di Assestamento Forestale). Il Comune di Laurenzana (Pz) ha provveduto a redigere il PAF per il decennio 2014-2023, per una superficie pari a 1.306,55 ettari, ricadendo per intero all'interno del Parco Nazionale dell'Appennino Lucano Val d'Agri e Lagonegrese e per 328,43 ettari nel sito RN2000 Basilicata IT9210005 (Abetina di Laurenzana). La ZSC (Zona Speciale di Conservazione) è caratterizzata dalla presenza di 2 Habitat forestali di interesse comunitario, il 91M0 e 9220. Con il Decreto del 16 settembre 2013 del Ministero dell'Ambiente e della Tutela del Territorio e del Mare, 20 SIC della Basilicata, grazie all'adozione delle MTC (Misure di Tutela e Conservazione) da parte della Regione, si sono evoluti in ZSC. Attraverso i PAF, i comuni applicano le MTC nell'ambito dei territori di loro pertinenza ricadenti nelle ZSC. Il presente contributo serve ad illustrare l'inserimento contestualizzato e pertinente di alcune MTC, sia generali che sito-specifiche, all'interno del PAF di Laurenzana, ed in particolare: 1) adozione di un adeguato strumento di pianificazione forestale; 2) mappatura degli esemplari monumentali di specie forestali; 3) censimento, mappatura e monitoraggio dei nuclei di rinnovazione naturale di abete bianco nel SIC; 4) censimento, mappatura e monitoraggio delle cenosi forestali con caratteri di boschi vetusti; 5) sostituzione delle specie forestali esotiche. Il piano prevede, inoltre, l'applicazione di percorsi adeguatamente individuati con le MTC, per rendere fruibile l'area dal punto di vista turistico. Il PAF amplia la sua valenza pianificatoria, rispondendo ad esigenze ambientali in senso lato e permettendo una gestione attiva della ZSC in questione.

*Parole chiave:* piani di assestamento forestale, misure di tutela e conservazione, ZSC.

*Keywords:* forest management plans, protection and conservation measures, SAC.

<http://dx.doi.org/10.4129/2cis-sc-pia>

## **1. Introduzione**

La Strategia Nazionale per la Biodiversità identifica 13 aree di lavoro tra le quali la n. 4 (foreste) e analizza minacce, obiettivi e priorità di intervento per la risorsa forestale, in particolare l'integrazione della tutela della biodiversità nella pianificazione forestale e di settore finalizzata ad una gestione sostenibile (MATTM, 2011). Le foreste italiane sono le più ricche al livello europeo, ospitando 117 specie differenti che corrispondono ai 2/3 del patrimonio floristico arboreo europeo. In Italia 10 categorie forestali, delle 14 ritenute più rappresentative della variabilità ecologica forestale del continente europeo (EEA). La legge n. 42 del 10 novembre 1998 della Regione Basilicata e s.s.m.i. "Norme in materia Forestale" e la D.G.R. n. 613/2008 "Linee guida per la redazione dei Piani di Assestamento Forestale" disciplinano la pianificazione forestale a livello regionale ed

hanno permesso di raggiungere risultati di un certo rilievo nel settore attraverso la stesura di Piani di Assestamento Forestale riguardanti una ragguardevole percentuale del territorio boschivo lucano. Il PAF può però divenire anche uno strumento gestionale di ampio respiro attraverso l'inclusione al suo interno di ulteriori elementi gestionali come le Misure di Tutela e Conservazione (MTC) dei Siti Rete Natura 2000. Un esempio concreto riguarda il PAF del Comune di Laurenzana in cui ricade per intero l'area protetta "Abetina di Laurenzana" che, con Decreto Ministeriale (MATTM) del 16 Settembre 2013, è stata designata Zona Speciale di Conservazione, evoluzione dei SIC forniti di strumento pianificatorio (Articolo 6 Direttiva Habitat). Ciò è stato possibile successivamente grazie all'adozione delle MTC da parte della Regione Basilicata con due D.G.R., la 951/2012 e la 30/2013 La ZSC in questione ricade interamente nella Zona 1 del Parco Nazionale dell'Appennino