



---

7° *Congresso della Società Italiana  
di Biologia Evoluzionistica*

Abstracts book

Rome, 28-31 August 2017  
*Department of Sciences, Roma Tre University, viale G.Marconi 446*

---

## *Organising Committee*

GABRIELE GENTILE (Coord.)

*Roma Tor Vergata*

GIULIANA ALLEGREUCCI

*Roma Tor Vergata*

MARCO ALBERTO BOLOGNA

*Roma Tre*

MONICA CAROSI

*Roma Tre*

DONATELLA CESARONI

*Roma Tor Vergata*

ALESSIO DE BIASE

*Roma Sapienza*

ANDREA DI GIULIO

*Roma Tre*

EMILIANO MANCINI

*Roma Tre*

MARCO OLIVERIO

*Roma Sapienza*

LEONARDO VIGNOLI

*Roma Tre*

## *Scientific Committee*

MARCO PASSAMONTI (Coord.)

*Bologna*

GIULIANA ALLEGREUCCI

*Roma Tor Vergata*

GIORGIO BERTORELLE

*Ferrara*

MAURIZIO CASIRAGHI

*Milano Bicocca*

PAOLO CIUCCI

*Roma Sapienza*

GIUSEPPE FUSCO

*Padova*

GABRIELE GENTILE

*Roma Tor Vergata*

SILVIA GHIROTTO

*Ferrara*

ALESSANDRO MINELLI

*Padova*

TELMO PIEVANI

*Padova*

ANDREA PILASTRO

*Padova*

OMAR ROTA STABELLI

*Edmund Mach Foundation (TN)*

## *Organising Secretariat*

CRISTINA COMPAGNO

*Management of Tourism and*

*Biodiversity - Elly Travel srl*

[info@sibe2017.uniroma2.it](mailto:info@sibe2017.uniroma2.it)

## *Sponsors*



PARCO  
ARCHEOLOGICO  
DELL'APPIA  
ANTICA



## Dating the impossible: the origin and divergence of the largest infection on Earth

Francesco Drago<sup>1</sup>, Matthias Scholz<sup>2</sup>, Nicola Segata<sup>3</sup>, Omar Rota-Stabelli<sup>2</sup>

<sup>1</sup> Università degli Studi di Padova,

<sup>2</sup> Fondazione Edmund Mach,

<sup>3</sup> CIBIO, Università degli Studi di Trento

*Wolbachia* are common intracellular bacteria of many arthropod and nematode species; they are responsible for likely the most widespread and evolutionary significant infection on Earth. *Wolbachia* are generally acquired by maternal transmission, but they often move across species boundaries, impairing the use of host divergences as independent calibration to infer their evolutionary history. Here we present a first systematic effort to estimate the origin and the divergence of the *Wolbachia* infection by coupling the molecular clock analysis of a genomic and an MLST data sets with three types of disputed calibration priors: a root prior for splits within Proteobacteria, the recently proposed *Nomada* host-symbiont co-divergence, and an adjusted mutation rate gathered from *Drosophila melanogaster* and *Nasonia* species. We first perform a hierarchical based model comparison on the MLST data set to select the best clock, replacement, and tree priors; we then used the best prior combination to calibrate both the genomic and the MLST data sets using different combinations of calibrations. Although divergence estimates are characterized by a high degree of uncertainty and strongly depend on the use of certain calibrations, most results point towards a pre-Cambrian origin of *Wolbachia* (its split from *Ehrlichia*). Diversification of *Wolbachia* supergroups is more recent, from the late Paleozoic to the Mesozoic: this is compatible with a long-fuse model of evolution, and suggests a co-radiation with holometabolous insects. Our results provide a comprehensive first effort to date *Wolbachia* evolution: results are model, prior, and data set dependent and reveal the difficulties embedded in the molecular dating of such elusive bacteria characterized by the absence of reliable calibration priors and a clear evolutionary history.