



EVOLUZIONE  
2017

---

*7° Congresso della Società Italiana  
di Biologia Evoluzionistica*

Abstracts book

Rome, 28-31 August 2017

*Department of Sciences, Roma Tre University, viale G.Marconi 446*

---

### *Organising Committee*

GABRIELE GENTILE (Coord.)  
*Roma Tor Vergata*  
GIULIANA ALLEGROCCI  
*Roma Tor Vergata*  
MARCO ALBERTO BOLOGNA  
*Roma Tre*  
MONICA CAROSI  
*Roma Tre*  
DONATELLA CESARONI  
*Roma Tor Vergata*  
ALESSIO DE BIASE  
*Roma Sapienza*  
ANDREA DI GIULIO  
*Roma Tre*  
EMILIANO MANCINI  
*Roma Tre*  
MARCO OLIVERIO  
*Roma Sapienza*  
LEONARDO VIGNOLI  
*Roma Tre*

### *Organising Secretariat*

CRISTINA COMPAGNO  
*Management of Tourism and  
Biodiversity - Elly Travel srl*  
[info@sibe2017.uniroma2.it](mailto:info@sibe2017.uniroma2.it)

### *Sponsors*



### *Scientific Committee*

MARCO PASSAMONTI (Coord.)  
*Bologna*  
GIULIANA ALLEGROCCI  
*Roma Tor Vergata*  
GIORGIO BERTORELLE  
*Ferrara*  
MAURIZIO CASIRAGHI  
*Milano Bicocca*  
PAOLO CIUCCI  
*Roma Sapienza*  
GIUSEPPE FUSCO  
*Padova*  
GABRIELE GENTILE  
*Roma Tor Vergata*  
SILVIA GHIROTTI  
*Ferrara*  
ALESSANDRO MINELLI  
*Padova*  
TELMO PIEVANI  
*Padova*  
ANDREA PILASTRO  
*Padova*  
OMAR ROTA STABELLI  
*Edmund Mach Foundation (TN)*

## Multiple sources of incongruence in *Wolbachia* phylogeny

Francesco Drago<sup>1</sup>, Omar Rota-Stabelli<sup>2</sup>

<sup>1</sup>Università degli Studi di Padova,

<sup>2</sup>Fondazione Edmund Mach

*Wolbachia* are maternally inherited intracellular endosymbionts bacteria, commonly found in the reproductive tissues of arthropods and nematodes. Because of its ability to manipulate host's reproduction and influence the transmissibility of host's pathogens, *Wolbachia* has substantial implications on the host species evolution and is a promising biological tool to control diseases and invasive pests. Various aspects of the biology of *Wolbachia* are still widely unexplored, particularly its biological interactions, spread and acquisition: this knowledge is an important prerequisite to properly characterize the evolution of this taxon. To further understand the biodiversity and the evolution of *Wolbachia* we compare the phylogenetic signals of two independent data sets: a gene-rich phylogenomic (GENOME) and a taxon-rich multilocus sequence typing (MLST) alignment. Our MLST data set provides a topology consistent with previous MLST phylogenies, but in conflict with the GENOME data set for the position of wZoo and wCte, infecting respectively termite and cat flea. These incongruences have been explored using various approaches to estimate intragenic recombination, evaluating strain misassignment and multiple *Wolbachia* infections on the same host species, and performing topology comparison tests. Results indicate that these discrepancies are due, at least partially, to undetected co-infections, resulting in different *Wolbachia* strains being sampled in the GENOME and the MLST dataset. Coalescent aware phylogenies further show among genes incongruences for wZoo in the GENOME data set: this is compatible with either horizontal transfer, undetected co-infections, or poor gene sampling for certain *Wolbachia* strains. Overall, our results suggest that discrepancies between our MLST and GENOME dataset are not due to stochastic (poor signal in MLST) or systematic errors (signal misinterpretation), but are caused by the complex epidemiological pattern of *Wolbachia* which may have resulted in different strains being sampled in different specimens and/ or have resulted in chimeric genome assemblies.