



IOBC-WPRS



# **Future IPM 3.0 towards a sustainable agriculture**

**IOBC-WPRS general assembly  
Meeting of the WGs Integrated protection in viticulture,  
Induced resistance in plants against insects and diseases and  
Multitrophic interactions in soil**

**15-20 October 2017, Riva del Garda, Italy**



Future IPM 3.0

## **BOOK OF ABSTRACTS**



## Protein-based products as resistance inducers: disease control and mechanisms of action

Martina Cappelletti, Michele Perazzoli, Andrea Nesler, Livio Antonielli, Gerardo Puopolo, Oscar Giovannini, Ilaria Pertot

First, second, third, fifth, sixth and seventh authors: Department of Sustainable Ecosystems and Bioresources, Research and Innovation Centre, Fondazione Edmund Mach, 38010 San Michele all'Adige, Italy; first author: Department of Agrifood, Environmental and Animal Sciences, University of Udine, 33100 Udine, Italy; third author: Bi-PA - Biological Products for Agriculture, B-1840 Londerzeel, Belgium; fourth author: Bioresources Unit, Department of Health and Environment, Austrian Institute of Technology, 3430 Tulln and der Donau, Austria  
E-mail address: [martina.cappelletti@fmach.it](mailto:martina.cappelletti@fmach.it)

### Highlights

- Leaf treatments with a protein derivative represent a sustainable strategy in plant protection, because they induce grapevine resistance, and change the structure of leaf microbial communities on grapevine
- Plant-protein hydrolysates reduce powdery mildew severity, and their biocontrol activity is affected by the protein source, degree of hydrolysis and peptide composition

### Introduction

Grapevine (*Vitis vinifera*) is one of the major fruit crops in the world, and downy mildew (caused by the oomycete *Plasmopara viticola*) is a serious disease that requires frequent fungicide applications. Increasing concerns about the negative impacts of pesticides on human health and the environment encourage the development of harmless alternatives to synthetic chemicals, such as resistance inducers (Delaunoy et al., 2014). Proteins and peptides represent a wide category of plant elicitors (Albert, 2013), and the protein derivative called Nutrient Broth (NB) showed a high efficacy in controlling powdery mildew under field conditions (Nesler et al., 2015). This study aimed to dissect the mechanisms of action of NB against grapevine downy mildew caused by the oomycete *P. viticola* and to develop low-cost protein hydrolysates from agro-industrial by-products.

### Material and methods

Grapevine plants (Pinot noir ENTAV115) grown under greenhouse conditions or *in vitro* (Nesler et al., 2015) were kept untreated (UNT) or treated with water (H<sub>2</sub>O), 3.0 g/l NB (Nesler et al., 2015), or with a commercial product based on laminarins (LAM, 0.75 ml/l Vacciplant, Belchim Crop Protection). RNA extraction and quantitative real-time PCR reactions were carried out for the amplification of pathogenesis-related genes (*PR-1*, *PR-2*, and *PR-4*), osmotins (*OSM-1* and *OSM-2*) and chitinase (*CHIT-3*) (Nesler et al., 2015). Collection of phyllosphere microorganisms, DNA extraction and amplification of bacterial (V6-V8 of the 16S rRNA) and fungal (ITS3-ITS4 of the internal transcribed spacer, ITS) fragments were performed as described by Cappelletti et al. (2016).

Soybean, rapeseed and guar meals were subjected to enzymatic (Alcalase or Flavourzyme at 1% or 50% E/S) or chemical (6N sulfuric acid, H<sub>2</sub>SO<sub>4</sub>; condition A: 121°C, 15 min, condition B: 100°C, 8 h) hydrolysis (Cappelletti et al., 2017). Courgettes (*Cucurbita pepo*) and powdery mildew caused by *Podosphaera xanthii* were selected as easy-to-handle study pathosystem. Courgette plants (cv Nero Milano) grown in greenhouse (Nesler et al., 2015) were sprayed with protein hydrolysates (1



g/l), water (H<sub>2</sub>O) or non-hydrolysed protein sources (N-H), and for the acid hydrolysis with a potassium sulfate (K<sub>2</sub>SO<sub>4</sub>) solution. The identification of peptides and amino acids was performed by an external service company (ISB Srl, Italy).

## Results and discussion

The preventive foliar application of NB reduced downy mildew severity as compared with control plants (UNT and H<sub>2</sub>O-treated), and the efficacy was higher in NB- than in LAM-treated plants. The expression levels of *PR-1*, *PR-2*, *PR-4*, *OSM-1*, *OSM-2* and *CHIT-3* genes were upregulated by NB before *P. viticola* inoculation, demonstrating the induction of grapevine resistance. Although the expression level of *CHIT-3*, *OSM-1*, *OSM-2* and *PR-4* was higher in LAM- as compared with NB-treated plants, LAM showed lower efficacy than NB against downy mildew, suggesting that multiple mechanisms of action are involved in the biocontrol activity of NB.

Indeed, NB changed the structure of phyllosphere bacterial and fungal populations as compared with control plants (UNT and H<sub>2</sub>O-treated), and these modifications were affected by the composition of the originally residing microbiome. The NB treatment increased the proportion of some genera (e.g. *Exiguobacterium*, *Pseudomonas*, *Serratia*, *Lysobacter*) that potentially include biocontrol strains, suggesting that these changes may contribute to disease control. Furthermore, experiments using *in vitro* grown plants, in the absence of phyllosphere microorganisms, showed that the NB reduced downy mildew symptoms as compared with H<sub>2</sub>O-treated plants, and induced the expression of *PR-2*, *PR-4*, *CHIT-3*, *OSM-1* and *OSM-2* before *P. viticola* inoculation. In conclusion, NB reduced downy mildew symptoms mainly by the induction of defence mechanisms in grapevine, and changed proportions of some microbial taxa linked to the biological control of plant pathogens, possibly providing a partial contribution to the control of downy mildew and to the activation of defence signalling pathways.

In order to develop cheaper and environmental-friendly protein-based products to control grapevine diseases, courgette powdery mildew was used as preliminary model pathosystem. Protein hydrolysates obtained by agro-industrial by-products were obtained, and guar hydrolysates significantly reduced powdery mildew symptoms. Particularly, two specific hydrolysis methods led to the formation of bioactive products (guar enzymatic hydrolysate Alcalase 50% and guar acid hydrolysate condition B). The biocontrol activity of hydrolysates was affected by the original protein source, the method and the degree of hydrolysis, namely the percentage of cleaved peptide bonds. The composition in free amino acids and peptide fragment could regulate plant responses to the pathogen infection. However, the use of strong acids during the hydrolysis causes an increase of salinity (K<sub>2</sub>SO<sub>4</sub>) of protein hydrolysates, which contributes to the disease control. The foliar application of low-cost protein hydrolysates represents an innovative approach to control crop diseases, and further studies are required to fully clarify their mechanisms of action and the effects on phyllosphere microorganisms.

## Acknowledgements

This project has received funding from the European Union's Seventh Framework Programme for research, technological development and demonstration under grant agreement no. 265865 (project PURE, theme FP7-KBBE-2010-4) and no. 324416 (project INNOVA, theme FP7-PEOPLE-2012-IAPP), and from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 722642 (project INTERFUTURE).



---

## References

- Delaunois B et al. 2014. Elicitors as alternative strategy to pesticides in grapevine? Current knowledge on their mode of action from controlled conditions to vineyard. *Environmental Science and Pollution Research International* 21: 4837-4846.
- Albert M 2013. Peptides as triggers of plant defence. *Journal of Experimental Botany* 64: 5269-5279.
- Nesler A et al. 2015. A complex protein hydrolysate acts as biogenic elicitor of plant resistance against powdery mildew. *Frontiers in Plant Science* 6: 715.
- Cappelletti M et al. 2016. Leaf treatments with a protein-based resistance inducer partially modify microbial communities of grapevine. *Frontiers in Plant Science* 7: 1053.
- Cappelletti M et al. 2017. Development and assessment of plant protein hydrolysates as biopesticides against zucchini powdery mildew. 69<sup>th</sup> International Symposium on Crop Protection - Session: Phytopathology I, Gent, Belgium, May 2017.