

Modelling spatio-temporal dynamics of snow depth and large herbivore's winter habitat selection


Emanuele Cordano¹, Emanuele Eccel², Federico Ossi², Julius Bright Bross^{3,4}, Wibke Peters⁵, Francesca Cagnacci^{2,3}

¹ Rendena100 Sole Proprietorship (www.rendena100.eu), Tione di Trento, TN, Italy; ² Research and Innovation Centre, Fondazione Edmund Mach, San Michele all'Adige, TN, Italy; ³ Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA; ⁴ Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, Oxford, United Kingdom; ⁵ Department of Biodiversity, Conservation and Wildlife Management, Bavarian State Institute of Forestry, Freising, Germany




Foreword

Winter snowy conditions and the wildlife management strategies play a critical role in governing the spatial distribution of ungulates in temperate ecosystems. Snow depth severely restricts roe deer distribution due to the limitations resulting from their small body mass (18-49 kg: Lister and Sumner (1998)) and short legs (50-60 cm: Holand and Linnell (1998)). Large roe deer living in snowy areas, employ a partial migration strategy, with all individuals overwintering in ranges characterized by less extreme snow conditions. Consequently, snow occurrence constitutes a proxy for animal location and movements. Due to the recently observed strong variability of climate and snow conditions in alpine areas, a **high-resolution and topography-accounted spatio-temporal modelling of snow cover is required.**



Hydrological modelling


GEOtop 2.0 Hydrological Model (Endrizzi et al. (2014), www.geotop.org) is a physically-based *spatially distributed* model solving water and energy balance differential equation, **producing snow depth area maps from meteorological data by taking into account snow melting process.**




Input weather forcings (25-year long):

- Observed time series in the weather stations (www.meteotrentino.it) (1 scenario);
- Downscaled data by regional climate model projections (Rockel, Will, and Hense (2008), courtesy of Emanuel Eccel and CMCC) (IPCC scenarios RCP4.5 and RCP 8.5) (2 scenarios).

Snow mean depth and snow cover are aggregated from daily to seasonal scale (NDJFMA)(Cordano (2019)). RCP4.5 and RCP8.5 illustrated scenarios are created with weather forcings having patterns similar to both 2013-2014 and 2014-2015 observed seasons, but modified accordingly to the projections.




Higher variability on snow occurs at medium and low elevations, i.e. *snow depth* > 5 cm, in the decade 2006-2015 (latest year of the analysis) than in the past decade 1996-2005. Future climate projections confirm this behaviour.



Ecological modelling

Roe deer in presence of snow strongly try to select zones with canopy cover and avoided places with high snow depths. Snow modelling results have been coupled with ecological campaign periods (1999-2002: (a); 2012-2015: (b))(see prediction plots of relative probability of use of forested habitats during, *Bright Bross et al., Article in preparation*)



Concluding remarks

1. Climate and snow variability is a challenge for researching on habitat selection by roe deer (*Capreolus capreolus*) in an alpine area;
2. Physically-based modelling allows to estimate snow depth in a gridded spatio-temporal coverage accounting for topographical details from terrestrial weather observations;
3. In the study area snow depth variability increases with climate change projections.

References

- Cordano, Emanuele. 2019. *Geotopbricks: An R Plug-in for the Distributed Hydrological Model Geotop*. <https://CRAN.R-project.org/package=geotopbricks>.
- Endrizzi, S., S. Gruber, M. Dall'Amico, and R. Rigon. 2014. "GEOtop 2.0: Simulating the Combined Energy and Water Balance at and Below the Land Surface Accounting for Soil Freezing, Snow Cover and Terrain Effects." *Geoscientific Model Development* 7 (6): 2831-57. <https://doi.org/10.5194/gmd-7-2831-2014>.
- Holand, A. Myrsterud, Ø., and J. D. C. Linnell. 1998. "Roe Deer in Northern Environments: Physiology and Behaviour." Edited by R. Andersen, P. Duncan, and J. D. C. Linnell. Oslo, Norway: Scandinavian University Press.
- Lister, P. Grubb, A. M., and S. R. M. Sumner. 1998. "Taxonomy, Morphology and Evolution of European Roe Deer." Edited by R. Andersen, P. Duncan, and J. D. C. Linnell. Oslo, Norway: Scandinavian University Press.
- Rockel, Burkhard, Andreas Will, and Andreas Hense. 2008. "The Regional Climate Model Cosmo-Clim (Clim)." *Meteorologische Zeitschrift* 17 (4): 347-48. <https://doi.org/10.1127/0941-2948/2008/0309>.